

## **Bioactivity characteristics of trypsin immobilized by affinity interaction and covalent conjugation**

Dae Ho Jang and Eun Kyu Lee<sup>\*,1</sup>

Bioprocessing Research Lab., Dept. of Chemical Engineering,

<sup>1</sup>Micro Biochip Center, Hanyang University, Ansan, Korea 425-791

Tel: 031-400-4072, FAX: 031-408-3779, [eklee@hanyang.ac.kr](mailto:eklee@hanyang.ac.kr)

For technical applications enzymes are usually immobilized to a large variety of supports <sup>1)</sup>. Also, non-denaturing immobilization of proteins is an important issue in protein chip applications <sup>2)</sup>. Protein functionality usually decreases when it is immobilized on solid surface because the tertiary structures of a protein could be distorted by immobilization. In order to solve this problem, many researchers have focused on using 'spacer arms' and exploiting 'oriented' immobilization <sup>1-4)</sup>.

We investigated the effects of immobilization chemistry on the yield of immobilization and the enzymatic activity of the immobilized enzymes, using trypsin as a model protein and aldehyde-decorated macroporous polymer beads (Toyopearl AF Formyl 650M, Tosho co., Japan) as a model matrix. Trypsin was immobilized by two methods; covalent conjugation by reductive amination (at pH 10.0) and affinity interaction via streptavidin-biotin system. The amount of covalently immobilized trypsin was 3.2 mg/ml-gel. It was about three times higher than that of the affinity method. However, when the specific activity of the immobilized trypsin was evaluated by its ability to digest BAPNA (400 Da), human insulin (5,870 Da), and BSA (66,000 Da), the covalent and affinity immobilized trypsin showed ca. 37% and 50%, respectively, of that of the soluble enzyme (on the BAPNA substrate). While the activity of the covalently immobilized enzyme decreased as the molecular size of the substrate increased, the affinity immobilized trypsin showed higher activity on insulin and BSA. This result seemed to indicate the streptavidin-biotin system allowed more steric flexibility in its interaction with a substrate molecule.

To confirm this, we studied the flexibility of immobilized trypsin using quartz

crystal microbalance-dissipation (QCM-D). SAMs (self-assembled monolayers) were formed on the Q-sensor surface by aminoalkanethiols, and glutaraldehyde was attached to the SAMs. Trypsin was immobilized on the modified Q-sensors in two ways: reductive amination (at pH 10.0) and the streptavidin-biotin system. The dissipation shift of the affinity immobilized trypsin was  $0.5 \times 10^{-6}$ , whereas that of the covalently attached enzyme was almost zero. This indicated the streptavidin-biotin system allowed higher flexibility than the covalent conjugation. These results suggested that the bioactivity of the immobilized enzyme could be strongly dependent on its molecular flexibility.

#### References

1. Meryem, N., Klaus, M. and Hans, B., Immobilization of trypsin on polyester fleece via different spacers (2001), *Enzyme and Microbial Technology*, 29, 567-574.
2. Heng, Z. and Michael, S., Protein chip technology (2003), *Current Opinion in Chemical Biology*, 7, 55-63.
3. Jaroslava, T., Oriented immobilization of biologically active proteins as a tool for revealing protein interactions and function (1999), *Journal of Chromatography B*, 722, 11-31.
4. Suh, C. W., Park, S. H., Park, S. G. and Lee, E. K., Covalent immobilization and solid-phase refolding of enterokinase for fusion protein cleavage (2005), *Process Biochemistry*, 40, 1755-1762.