

[IM07] Molecular Hydrogen and Silicon Monoxide Imaging of the NGC 1333 IRAS 4A Outflow

Minho Choi¹, Klaus W. Hodapp², Masahiko Hayashi³, Kentaro Motohara^{3,4},
Soojong Pak^{1,5}, and Tae-Soo Pyo³

¹*Korea Astronomy and Space Science Institute, Hwaam 61-1, Yuseong, Daejeon 305-348*, ²*Institute for Astronomy, University of Hawaii, 640 North A'ohoku Place, Hilo, HI 96720, USA*, ³*Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'ohoku Place, Hilo, HI 96720, USA*, ⁴*Institute of Astronomy, University of Tokyo, Mitaka, Tokyo 181-0015, Japan*, ⁵*Department of Astronomy and Space Science, Kyung Hee University, Seocheon, Giheung, Yongin, Gyeonggi 446-701*

The NGC 1333 region was observed in the near-IR H₂ 1-0 S(1) line and the 7 mm SiO line. The northeast-southwest bipolar outflow driven by IRAS 4A was studied by combining the H₂ and the SiO maps. The southwestern outflow lobe curves smoothly, and the position angle increases with the distance from the driving source. The base and the outer tip of the northeastern outflow lobe are located at positions opposite to the corresponding parts of the southwestern lobe. This point-symmetry suggests that the outflow axis may be drifting or rotating clockwise in the plane of the sky and that the cause of the axis drift may be intrinsic to the outflow engine. The axis drift model provides a good explanation for the large deflection angle of the northeastern outflow.

[IM08] Distribution of molecular clouds in the Outer Galaxy

Miju Kang^{1,2}, Youngung Lee²

¹*Chungnam National University*, ²*Korea Astronomy Space Science Institute*

We present an analysis of the molecular gas distribution in the second quadrant of the Galactic Plane, using the ¹²CO J=1-0 Outer Galaxy Survey of FCRAO(Five College Radio Astronomy Observatory). We estimate kinematic distances toward all the identified molecular clouds except for those with negative velocities, which could be all local clouds, using a flat rotation curve, and the Clemens' rotation curve. The spiral arm in the Outer Galaxy will be delineated based on the distribution status. Comparison will be discussed with other distance determination in this direction. Several physical properties of the molecular clouds on the arm and those between the arms will be discussed.