

**E-E2-29****Enhanced  $\alpha$ -tocopherol content and free radical scavenging activity of *Codonopsis lanceolata* transformed with  $\gamma$ -TMT gene**

**Bimal Kumar Ghimire<sup>1</sup>, Sun Seong Kwon<sup>1</sup>, Hyun Young Kil<sup>1</sup>, Eun Won Seo<sup>1</sup>, Myong Jo Kim<sup>2</sup>, Amal Kumar Ghimeray, Eun Jeong Goh<sup>1</sup> and Chang Yeon Yu<sup>1</sup>†**

<sup>1</sup>Bioherb Research Institute, Kangwon National University, Chunchon 200-701, Korea

<sup>2</sup>Bio-resources Technology, Kangwon National University, Chunchon 200-701, Korea

*Codonopsis lanceolata* is an important medicinal plant that synthesis tocopherol, which is an antioxidant and a phytochemical associated with reduced cancer risk, coronary heart disease, and psychoneurosis.  $\gamma$ -tocopherol is the most abundant form of tocopherol found in the *Codonopsis lanceolata*, whereas,  $\alpha$ -tocopherol, which is the most biologically active form of vitamin E, exist in low amount. Therefore, to shift the tocopherol composition, over expression of the  $\gamma$ -TMT gene successfully established in *Codonopsis lanceolata*. HPLC analysis of leaf and roots of T<sub>1</sub> transgenic lines revealed an increased  $\alpha$ -tocopherol content by 6.52 and 2.60 fold respectively over the controls plants. The antioxidant activity of methanolic crude extracts of transgenic and non transgenic control plants was investigated using 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging method. Free radical scavenging activity was evaluated using  $\alpha$ -tocopherol, Butylated Hydroxyanisole (BHA) and Butylated Hydroxy toluene (BHT) as standard antioxidants. In general, the transgenic *Codonopsis lanceolata* showed strong DPPH radical scavenging activity over the non transformed control plant. Thus, over expression of  $\gamma$ -TMT gene resulted in increased  $\alpha$ -tocopherol content with an enhanced antioxidant activity.

† Corresponding author : (E-mail) [cyyu@kangwon.ac.kr](mailto:cyyu@kangwon.ac.kr) (Tel) +82-33-250-6411

**E-E2-30****Inhibitory effect mechanism of *Prunus persica* (Fructus) extract on melanin synthesis and photoaging**

**Hyen Joo Park<sup>1</sup>, Kwang-Kyun Park<sup>2</sup>, Jae-Kwan Hwang<sup>3</sup>, Won-Yoon Chung<sup>2</sup> and Sang Kook Lee<sup>1\*</sup>**

<sup>1</sup>College of Pharmacy, Ewha Womans University, Seoul 120-750

<sup>2</sup>College of Dentistry, Yonsei University, Seoul 120-752

<sup>3</sup>College of Life Science, Yonsei University, Seoul 120-749

In our previous study, *Prunus persica* (Fructus) extract (PFE) was found to be the inhibitory effects on melanogenesis with employing tyrosinase activity and melanin synthesis in melan-a cells. The BuOH-soluble fraction, especially, inhibited melanin formation in a dose-dependent manner without cytotoxic effects. In this study, the gene expression levels of tyrosinase and tyrosinase-related protein-1, -2 (TRP-1, -2) were analyzed by a real-time reverse transcriptase-polymerase chain reaction (RT-PCR). Exposure of mouse melan-a cells to PFE and the BuOH-soluble fraction of the extract markedly reduced the mRNA expression of tyrosinase and TRP-1 & -2.

In addition, antiaging effects of PFE were performed using HaCaT human keratinocytes and human skin fibroblast cells. PFE attenuated cell damage induced by hypoxanthine-xanthine oxidase in a dose-dependent manner indicating that PFE has the potential of the scavenging effect of reactive oxygen species (ROS) in human skin cell. Moreover, PFE significantly suppressed UVA-induced ROS production determined by the oxidation of DCFH to highly fluorescent 2',7'-dichlorodihydro-fluorescein (DCF) using FACS analysis. Additional study revealed that UVA irradiation of HaCaT cells increased the gelatinolytic activities of MMP-2, -9, which was significantly suppressed by the treatment with PFE at the test concentration of 5 mg/ml. PFE also showed the suppressive effect of the mRNA expression of MMP-9.

These results suggest that PFE might be applicable as natural ingredients for skin-whitening and antiaging agents.

\*corresponding author: Tel. 02-3277-3023, e-mail: [sklee@ewha.ac.kr](mailto:sklee@ewha.ac.kr)