

FA01

Beyond Micromagnetic Modeling of Spin Dynamics in Confined Magnetic Elements

Sang-Koog Kim*

Research Center for Spin Dynamics and Spin-Wave Devices, Seoul National University, Seoul 151-744, Republic of Korea
 Nanospintronics Laboratory, Department of Materials Science and Engineering, College of Engineering,
 Seoul National University, Seoul 151-744, Republic of Korea

*Corresponding author: sangkoog@snu.ac.kr, Phone: +82 2 880 5854, Fax: +82 2 885 1457

Magnetization dynamics in bulk and geometrically confined micrometer or nano-size magnetic elements is of considerably growing interest in the research field of time-variable magnetic phenomena. Individual spin motions excited in magnetic nanostructures are well described by the Landau-Lifshitz-Gilbert (LLG) equation of motion. The numerical micromagnetic calculation of various model systems of restricted geometry [1-8] based on the LLG equation is likely to be a useful bridge connecting the gap between theories and experiments on ultrafast magnetization dynamics. During the past five years, our research group have conducted micromagnetic simulations on a variety of patterned thin films [1-8] and have obtained interesting results in the area of vortex dynamics, spin waves, and domain wall (DW) motions. In this talk, our recent simulation results on the vortex motions [1], interactions between vortex and antivortex [7,8], vortex-core reversals [1,2,3], and their related phenomena [2,3,4] will be presented. This talk covers the exact mechanism of vortex-core reversals [1], spin-wave radiation [2,3,4], the behaviors of those spin waves traveling in nanowire-waveguides [2,3], and a complex behavior of the periodic transformation of the internal structures of moving DWs in magnetic nanostripes [5].

Ki-Suk Lee, Sangkook Choi, Jun-Young Lee, and K. Y. Guslienko are acknowledged for their research works at ReC-SDSW. This work was supported by Creative Research Initiatives (ReC-SDSW) of MOST/KOSEF.

REFERENCES

- [1] K.-S. Lee, K. Y. Guslienko, J.-Y. Lee, and S.-K. Kim (unpublished).
- [2] S. Choi, K.-S. Lee, K. Y. Guslienko, and S.-K. Kim, *Phys. Rev. Lett.* **98**, 087205 (2007).
- [3] S. Choi, K.-S. Lee, and S.-K. Kim, *Appl. Phys. Lett.* **89**, 062501 (2006).
- [4] K.-S. Lee, S. K. Choi, and S.-K. Kim, *Appl. Phys. Lett.* **87**, 192502 (2005).
- [5] J.-Y. Lee, K.-S. Lee, S. Choi, K. Y. Guslienko, and S.-K. Kim (unpublished).
- [6] S. Choi, S.-K. Kim, V. E. Demidov, and S. O. Demokritov, *Appl. Phys. Lett.* **90**, 083114 (2007).
- [7] S.-K. Kim, K.-S. Lee, B.-W. Kang, K.-J. Lee, and J. B. Kontricht, *Appl. Phys. Lett.* **86**, 052504 (2005).
- [8] K.-S. Lee, B.-W. Kang, Y.-S. Yu, and S.-K. Kim, *Appl. Phys. Lett.* **85**, 1568 (2004).

FA02

Vortex Core Dynamics Imaged by Time-Resolved Scanning Transmission X-Ray Microscopy

**H. Stoll*¹, B. Van Waeyenberge², A. Puzic¹, K.W. Chou¹, D. Dolgos¹, M. Curcio¹,
 A. Vansteenkiste², I. Neudecker³, G. Woltersdorf³, C.H. Back³, and G. Schütz¹**

¹Max Planck Institute for Metals Research, Heisenbergstr. 3, 70569 Stuttgart, Germany

²Department of Subatomic and Radiation Physics, Ghent University, Proeftuinstraat 86, 9000 Gent, Belgium

³Department of Physics, University of Regensburg, Universitätsstr. 31, 93053 Regensburg, Germany

*Corresponding author: stoll@mf.mpg.de, Phone: +49 711 689 1848, Fax: +49 711 689 1952

Fast magnetization dynamics of ferromagnetic structures on a sub-micron length scale are currently attracting substantial scientific interest for both, technological and fundamental reasons [1]. The implementation of stroboscopic measurement techniques into a scanning transmission X-ray microscope (STXM, ALS, Beamline 11.0.2) appears to be a suitable method for such studies, as a time resolution of 50-70 ps combined with a lateral resolution of 20-40 nm can be achieved.

The vortex core plays a key role in the dynamics of micron and sub-micron sized ferromagnetic structures containing a single vortex [2]. New ways were developed to switch the out-of-plane core polarization. The reversal of the vortex core by excitation with short bursts of an in-plane alternating field was discovered recently [3]. Dynamic switching of the vortex core polarization was even possible with a single period magnetic field burst as low as 1.5 mT (Fig. 1), in contrast to (static) perpendicular fields of half a Tesla which were needed so far.




Fig. 1: Experimental results of switching the vortex core polarization [3].

REFERENCES

- [1] I. Tudosa, C. Stamm, A.B. Kashuba et al., *Nature* **428**, 831 (2004).
- [2] J. Millat and A. Thiaville, *Science* **298**, 555 (2002).
- [3] B. Van Waeyenberge, A. Puzic, H. Stoll et al., *Nature* **444**, 461 (2006)