

Results: 전체적으로는 평균연령은 34.8 ± 3.9 세, 채취된 난자 수는 11.31 ± 7.7 개, 양질의 난자 수는 9.75 ± 6.71 개, 이식배아수는 3.84 ± 1.59 개, 이식당 임신율은 35%였으며 216주기에서 획득 난자의 수가 3개 미만이었다. 획득 난자 수가 3개 미만인 주기에서 40세 미만 (155주기)과 40세 이상 (61주기)을 비교한 결과 채취된 난자 수와 이식 배아 수에서는 차이가 없었으나 (2.1 ± 0.8 Vs. 1.8 ± 0.9 , 1.6 ± 0.7 Vs. 1.5 ± 0.7 , $p=ns$) 임상적 임신율은 19.4%와 6.6%로써 통계적으로 유의하게 ($p=0.02$) 차이가 있었다. 또한 minimal stimulation 및 FSHrec/GnRH antagonist의 배란유도에 따른 임신율도 양군 모두 통계적으로 차이가 없었다.

Conclusion: 결론적으로 난소의 기능이 저하된 환자의 체외수정 및 배아이식술은 40세 미만에서는 배란유도 방법에 관계없이 적절한 임신율을 기대할 수 있으나 40세 이상의 환자에서는 임신율이 저조하여 배란유도 방법을 포함한 다양한 보완 및 대체적 방법을 개발하는 것이 필요하다고 하겠다.

P-40 Comparison of Clinical Outcomes between Conventional in vitro Fertilization (IVF) and Intracytoplasmic Sperm Injection (ICSI) in ART Cycles with Low Number of Retrieved Oocytes

Sun-Hee Lee¹, Haengseok Song¹, Myo Kyung Kim¹, Dong-Wook Park¹, Mi-Ra Shin¹, Jin Young Kim², Mi Kyoung Koong², In Ok Song², Jin Hyun Jun¹

¹Laboratory of Reproductive Biology and Infertility, ²Department of Obstetrics and Gynecology, Cheil General Hospital, Kwandong University College of Medicine

Objectives: ICSI was initially developed to treat male infertility for fertilization and has significantly improved success rates of ART with male infertility. Recently its application has been expanded to the cases of unexplained infertility, poor fertilization and low egg number without comprehensible evidence to support its use. The aim of this study was to investigate whether, in cases of low oocyte number with normal semen parameters, the efficacy of ICSI is better than that of conventional IVF with respect to the rates of fertilization and pregnancy outcomes.

Methods: In total 288 cycles, number of retrieved oocytes was under five and semen analysis was normal parameters - CSI was performed in 229 cycles and conventional IVF was performed in 59 cycles. All female partners were less than 38 years old. The rates of matured oocyte, fertilization, good embryo, clinical pregnancy, implantation, and delivery were compared. Total fertilization failure was evaluated to examine whether there is an increased risk of cancellation rate of embryo transfer in conventional IVF. Chi square test and t-test were used for statistical analysis. P values < 0.05 were considered significant.

Results: There were no significant differences between two groups in mean age of female partner, oocyte maturation rate, fertilization rate, good embryo rate, and mean number of transferred embryos. Cancellation rate of embryo transfer due to total fertilization failure was similar between two groups (9.1% for IVF vs. 11.3% for ICSI). Consequently, there were no differences in implantation and pregnancy outcomes. In fact, clinical pregnancy and implantation rates were slightly higher in conventional IVF group (28.0% and 18.6%) than that of ICSI group (23.9% and 16.8%), respectively.

Conclusion: Our results suggest that conventional IVF is as efficient as ICSI in all aspects of clinical outcomes of ART cycles with low number of retrieved oocytes and normal semen parameters.