

P83

### Comparison on the Activities According to Bees' Numbers of *Osmia cornifrons* and *O. pedicornis* Released in the Apple Orchards

Sang-beom Lee, Kyung-Yong Lee, In-Gun Park, Hyung-Joo Yoon, Sun-Joo Jang,  
Jin-Geun Lee<sup>1</sup> and Byung-Hoon Im<sup>1</sup>

Department of Agricultural Biology, The National Institute of Agricultural Science & Technology, RDA,  
Suwon 441-100, Korea

<sup>1</sup>Kyungbuk Provincial Agricultural Research & Extension Service, Sangjoo, Kyungbuk, 742-040, Korea

This study was examined the pollinating activity and effect according to bees' numbers of *Osmia cornifrons* and *O. pedicornis* released in the apple orchard. The activities of out-going and in-coming bees of 750 heads and 1,000 heads groups released were more effective than those of 500 heads groups released. Also, the foraging activity and the pollen collecting activity of 750 heads and 1,000 heads groups released were more effective than those of 500 heads groups released. The rate of egg laid by *O. cornifrons* in the bamboo-tube was more higher than that by *O. pedicornis*. The normal fruit rate was showed *O. cornifrons* of 1,000 heads group released ; 41.3%, *O. cornifrons* of 750 heads group released, *O. pedicornis* of 750 heads and 1,000 heads groups released; 32~33%, and *O. cornifrons* and *O. pedicornis* of 500 heads groups released ; 21~27% in those order. Therefore, for producing of high quality fruit, *O. cornifrons* and *O. pedicornis* must be released over 750 heads groups during the apple blossoming season in apple orchard.

**Key words:** *Osmia cornifrons*, *Osmia pedicornis*, bee, apple orchard, pollinating activity

P84

### Transgene Expression Stability in *CMVP0-CP* Introduced Chili Peppers

Lee Sung Nam, Kwon Ji Yeon<sup>2</sup>, Choi Sun Hee, Harn Chee Hark<sup>1</sup>,  
Ryu Ki Hyun<sup>2\*</sup> and Kim Hyun Hee\*

Department of Life Science, Sahmyook University, Seoul 130-650, Korea

<sup>1</sup>Biotechnology Institute, Nongwoo Bio Co. Ltd., Yeosu 469-885, Korea

<sup>2</sup>Plant Virus GenBank, Seoul Women's University, Seoul 139-774, Korea

Many genetically modified crops were developed but the genetic instability of the introduced gene came to be a big problem to be solved for application of the transgenic plants. The existence and expression stability of transgene in *CMVP0-CP* introduced transgenic chili peppers was investigated with two lines, H15 and B20. Genomic DNAs, total RNAs and total proteins extracted from young leaves of non-transgenic and transgenic plants were analyzed by PCR, RT-PCR, and Western blot techniques. The transgenes were confirmed to exist stably during two generations of T3 and T4, and transcription and translation of the gene were detected constantly in T3 and T4 plants. It seems that the existence and expression stability of the introduced gene in *CMVP0-CP* transgenic chili peppers keeps up during the generations.

**Key words:** Instability, transgene, *CMVP0-CP*, chili pepper, PCR, RT-PCR, Western blot, stability