

관절와 상완 관절의 불안정증의 병태 생리와 분류 및 치료 방침의 결정

전남대학교 의과대학 정형외과학교실

김 명 선

Factors maintaining the Shoulder Joint Stability

1. Static stability factors

- 1) Articular conformity
- 2) Articular version
- 3) Glenoid labrum
- 4) Negative intraarticular pressure
- 5) Adhesion-cohesion
- 6) Capsuloligamentous structures
- 7) Rotator cuff

2. Dynamic stability factors

- 1) Rotator cuff
- 2) Long head of biceps brachii
- 3) Scapular rotators
- 4) Proprioception

Pathophysiology of Glenohumeral (GH) Instability

1. Articular version abnormality

- 1) Excessive glenoid retroversion
 - : In most cases, eccentric articular surface wear
 - : Infrequently, a variant of glenoid dysplasia
 - : contributing factor to post. instability

2. Bankart lesion

- 1) Detachment of the capsulolabral complex from the glenoid rim & scapular neck

- 2) Bankart variant: ALPSA (Anterior Labroligamentous Periosteal Sleeve Avulsion) lesion
- 3) First time patients c traumatic ant. shoulder dislocation: 97 % isolated Bankart lesion without intracapsular injury (Tayler & Arciero, Am J Sports Med, 1997)
- 4) Recurrent complete dislocation requires an additional pathoanatomic components (capsular plastic deformation or stretch)

3. Hill-Sachs lesion

- 1) Ant. dislocation: >80%, Ant. subluxation: 25%
- 2) Atraumatic instability that did not respond to conservative therapy: 60 %
(Werner et al, Arthroscopy, 2004)
- 3) The small Hill-Sachs lesion : not major contributor to recurrent ant. instability
- 4) >30% Hill-Sachs lesion: Recurrent ant. instability, even with capsular repair (by Rowe et al)

4. Glenoid bony loss

- 1) Repetitive overhead activities may load GH joint asymmetrically & lead to posterior glenoid erosion
(Gupta & Lee, JSES, 2005)
- 2) 78% chronic ant. GH instability patients: osseous lesion of glenoid (bony Bankart or wear related to repeated instability)
(Edwards et al, Arthroscopy, 2003)
- 3) > 25% bony loss of glenoid surface: bony reconstruction (Bigliani et al, Am J Sports Med, 1998)
- 4) Bone loss to convert normal glenoid to 'Inverted pear configuration'
: Particular risk for Redislocation after surgery
: recommend a coracoid process transfer (Latarjet procedure)
(Burkhart & DeBeer, Arthroscopy, 2000)
- 5) Overall aim of any reconstructive procedure directed at larger defects
: to deepen the socket & support the capsule

5. Capsular injury

- 1) Intrasubstance injury
: 55% of ant. dislocation → demonstrated capsular rupture by arthrography (by Reeves)
: 15% of ant. dislocation → labral detachment & ant. capsular rupture (Symeonides, JBJS(Br), 1972)
: Experimental Stress-Strain data at failure of IGHLIC (Bigliani et al, J Orthop Res, 1992)
→ before failure, significant plastic deformation occurred
→ may be acquired through submaximal trauma (single or repetitive) without causing rupture or detachment
: When ant. shear force overcomes capsular tensile strength or rotator cuff cannot effectively contract
→ Ligament may fail on ultrastructural level
: Joint capsule of instability patient

→ the amount of stable & reducible cross-links (which is abundant in remodeling tissue) ↑

(Rodeo et al, *Am J Sports Med*, 1998)

→ Histologic changes, (denuded synovial layers, subsynovial edema, cellularity ↑, vascularity ↑)

(McFarland et al, *Am J Sports Med*, 2002)

: Age-related attrition of rotator cuff tissue is greater than in capsular tissue

→ such that ant. dislocation commonly results in rotator cuff tear, potentially leading to capsular injury in older patients

2) HAGL (Humeral Avulsion of the Glenohumeral Ligament) lesion

(by Wolf et al, *Arthroscopy*, 1995)

: Traumatic ant. instability who show no signs of Bankart lesion

: Thickened, rolled edge of capsular defect

: Typically found in the inf. pouch of the shoulder below the level of subscapularis muscle

: Associated GH abnormalities : most common form; Rotator cuff tear (> 90%; subscapularis tear)

: Bony HAGL (BHAGL) lesion (Bach et al, *JBJS(Br)*, 1988)

→ bony avulsion of humeral neck

: HAGL lesion should be repaired anatomically

3) Repetitive injury

: Overhead athletes (pitchers, throwers, swimmers, volleyball players, tennis players, etc)

→ relating to instability (repetitive stresses → lead to microtrauma)

: The cumulative effect of repetitive subfailure strain causes irreversible stretching of IGHL

→ shoulder instability

(Pollock et al, *JSES*, 2000)

: Repetitive rotational motion of the GH joint may also contribute to instability

6. Capsular laxity

1) Intrinsic capsular laxity

: The degree of laxity varies among individuals

: It is unclear if constitutional laxity is a risk factor for clinical shoulder instability

: Asymptomatic shoulder can exhibit a range of rotational or translational motion comparable to that of symptomatic instability shoulder.

: Asymptomatic subluxation or even dislocation may occur in 'Normal shoulder' at the time of anesthesia

2) Inherited Disorders of Collagen

: relatively rare

: most associated collagen disorder: Ehlers-Danlos syndrome (EDS)

→ Laxity ↑, problem with wound healing, vascular anomalies

Classification of Glenohumeral Instability

1. Classification according to 5 factors

1) Direction (방향)

- (1) Unidirectional: Anterior, Posterior, Inferior
- (2) Bidirectional: Anteroinferior, Posteroinferior
- (3) Multidirectional

2) Degree (정도)

- (1) Dislocation (탈구)
- (2) Subluxation (아탈구)

3) Mechanism (발생 기전)

- (1) Traumatic (외상성)
- (2) Atraumatic (비외상성)
- (3) Acquired (repetitive microtrauma, 후천성)
- (4) Congenital (선천성)
- (5) Neuromuscular (근신경성): Erb' palsy, Cerebral palsy, Seizures

4) Frequency (빈도)

- (1) Acute (급성)
- (2) Chronic (만성): Recurrent, Fixed
- (3) Habitual

5) Voluntariness (수의 여부)

- (1) Involuntary (불수의성)
- (2) Voluntary (수의성)

2. Classification by Thomas & Matsen (JBJS(Am), 1989)

1) TUBS (Traumatic, Unidirectional, Bankart, Surgery)

- : Patients with traumatic etiology
- : usually have unidirectional instability
- : often have obvious Bankart lesion
- : frequently require surgery when the instability is recurrent

2) AMBRII (Atraumatic, Multidirectional, Bilateral, Rehabilitation, Inferior capsular shift, Interval closure)

- : Patients with atraumatic etiology
- : often have multidirectional laxity

- : that is frequently bilateral
- : and usually responds to a rehabilitation program
- : However, should surgery be performed, the surgeon must pay particular attention to performing an inferior capsular shift & closing the rotator interval

3) Classification by Rockwood (Orthop Trans, 1979)

- (1) Type 1: Traumatic subluxation without previous dislocation
- (2) Type 2: Traumatic subluxation after a previous dislocation
- (3) Type 3A: Voluntary subluxation in patients with psychiatric problems
- (4) Type 3B: Voluntary subluxation in patients without psychiatric problems
- (5) Type 4: Atraumatic involuntary subluxation

4) Classification by Gerber & Nyffeler (CORR, 2002)

(1) Static instability (Class A)

- ㄱ. Static Superior Subluxation (Class A1)
- ㄴ. Static Anterior Subluxation (Class A2)
- ㄷ. Static Posterior Subluxation (Class A3)
- ㄹ. Static Inferior Subluxation (Class A4)

(2) Dynamic instability (Class B)

- ㄱ. Chronic, Locked Dislocation of the shoulder (Class B1)
- ㄴ. Unidirectional Instability without Hyperlaxity (Class B2)
- ㄷ. Unidirectional Instability with Hyperlaxity (Class B3)
- ㄹ. Multidirectional Instability without Hyperlaxity (Class B4)
- ㅁ. Multidirectional Instability with Hyperlaxity (Class B5)
- ㅂ. Unidirectional or Multidirectional Instability with Voluntary Reduction (Voluntary Instability) (Class B6)

(3) Voluntary dislocations (Class C)

5) Classification of Posterior instability

(1) Posterior dislocation

- ㄱ. Acute posterior dislocation
- ㄴ. Chronic (locked) posterior dislocation

(2) Recurrent posterior subluxation

- ㄱ. Volitional
- ㄴ. Psychogenic
 - : learned

(3) Dysplastic

- : Glenoid retroversion
- : Humeral head retrotorsion

(4) Acquired

- : Soft tissue deficiency
- : Bony deficiency
- : Scapulothoracic dysfunction

Decision making in the Treatment of Glenohumeral(GH) Instability

- Decision making should be individualized depending on
Type of instability, Pathophysiology, Age, Sex, Patient Activity, Patients demand, Associated injury, etc.
- Treatment Option
 - : Nonoperative Tx: Immobilization c Rehabilitation & Exercise
 - : Surgical Tx: Arthroscopy or Open surgery
- Consideration in Decision making of Treatment choice
 - : Frequency (1st episode vs. Recurrent)
 - : Age (Young vs. Old)
 - : Pathology (Soft tissue vs. Bone < Glenoid or Humeral bone defect)
 - : Duration (Acute vs. Chronic)
 - : Force (Traumatic vs. Atraumatic)
- Good Candidates for Nonoperative Treatment
 - : Children
 - : Voluntary Instability patient
 - : Posterior GH Instability patient
 - : Those requiring supranormal ROM (Baseball pitcher, Gymnast, Dancer, etc)

1. Anterior Instability

- 1) Risk factors of Recurrence after linital Anterior Dislocation
 - (1) Younger age: < 20 years: 90% recurrence, > 40 years: 10~15% recurrence
 - (2) Men > Women
 - (3) Athletes > Nonathletes
 - (4) Activity & Occupation: the Military, Overhead worker, Manual labor
 - (5) Severity of Original Trauma
 - the more easily the dislocation occurred initially, the more easily it recurs
 - (6) Effect of Fracture
 - lower recurrence when Frist time dislocation is associated with Greater tuberosity fracture
- 2) Indications of Surgical Treatment
 - (1) Irreducible, Open, Recurrent dislocation,
 - (2) Failed nonoperative treatment,
 - (3) Glenoid bone loss: > 25%,
 - (4) Large engaging Hill-Sachs lesion

- 3) Relative Contraindications of Surgical Treatment
 - (1) Recurrent instability associated with uncontrolled epilepsy
 - (2) Inferior subluxation with Stroke or Deltoid insufficiency
 - (3) Multidirectional instability with Voluntary instability

- 4) Relative Indications of A/S Repair
 - (1) Traumatic Unidirectional Injury
 - (2) Noncontact sport athletes
 - (3) Thick Mobile Bankart lesion
 - (4) Little or No discernible Capsular Laxity
 - (5) Glenoid bone loss: < 25%
 - (6) Non-engaging Hill-Sachs lesion

- 5) Relative Indications of Open Repair
 - (1) HAGL(Humeral Avulsion of the Glenohumeral Ligaments) lesion
 - (2) Capsular ruptures
 - (3) Previous Failed Open or A/S repair
 - (4) Prior Failed thermal capsulorrhaphy
 - (5) Significant Glenoid or Humeral bone loss
 - (6) Irreducible chronic dislocation

- 6) Recommended Treatment of Bankart & Hill–Sachs Lesions
 - (1) Arthroscopic repair with or without Remplissage procedures
: Small or moderate size lesions,
 - (2) Open bone grafting techniques
: Significant glenoid bone loss (> 25%)
: Large engaging Hill-Sachs deformity of the humerus

Type of Lesion	Recommended Treatment
Large acute bony Bankart	Arthroscopic or open internal fixation
Small or medium-size acute bony Bankart (< 25% of inferior glenoid diameter)	Arthroscopic Bankart repair using suture anchors
Large chronic bony Bankart	Modified Latarjet reconstruction, preceded by arthroscopic treatment of concomitant pathology
Large Hill-Sachs (normal glenoid)	Arthroscopic Bankart repair and remplissage of Hill-Sachs defect
Borderline large bone loss on either side	Arthroscopic Bankart repair and remplissage of Hill-Sachs defect

2. Posterior Instability

: Relatively uncommon debilitating condition

: Careful evaluation, Vigorous nonoperative management, Carefully selected surgical approaches

(Robinson & Aderinto, JBJS(Am), 2005)

1) Several aspects complicating the Treatment choice for Recurrent Posterior Instability

(1) Most often atraumatic (vs. Anterior Instability)

(2) Pathogenesis

: Multifactoral

: Complex

: Less well understood than Anterior Instability

(3) Results of Posterior repair

: at least 10 times worse than the average for Anterior repair

(4) Recurrent Posterior Instability

: functional limitation & pain can be minimal

: is not a definite indication for surgery & stressed the need for careful patient selection before surgical reconstruction

(Tibone et al, Am J Sports Med, 1981)

2) Nonoperative Tx can be well managed by

(1) Patient Education

(2) Muscle strengthening

(3) Neuromuscular retraining

3) Indications of Surgical Stabilization

- Recurrent Involuntary Posterior subluxation or Dislocation

: in spite of concerned effort at a Well-Structured Rehabilitation program

4) Essential Consideration for Correct Surgery

(1) identify All directions of Instability

(2) identify Any Anatomic Predisposing factors

: Glenoid or humeral defect

: Abnormal Glenoid version

: RC tears

: Neurologic injuries

: Generalized ligamentous laxity

3. Multidirectional Instability (MDI, =Atraumatic Instability)

- Treatment goal

: Restoration of shoulder function by increasing the effectiveness of Concavity Compression (Dynamic compression of humeral head into glenoid concavity)

1) Characteristics of Pathophysiology

- (1) Loss of proper neuromuscular control of humeroscapular positioning
 - : cannot be restored surgically
 - : requires prolonged adherence to well-constructed reconditioning program
- (2) Dysfunction of Concavity Compression effect
- (3) There is no single lesion that can be repaired
- (4) Capsular laxity might not be the essential lesion
- (5) Ability of surgery alone to cure Atraumatic Instability is limited
 - : not easy to restore the problem of Neuromuscular control or Relative glenoid flattening

2) Presurgical Consideration in the Treatment of MDI

- (1) The patient must have Major functional problems that are clearly related to instability
- (2) The patient must clearly understand that Good strength & Proper kinematic technique are the primary stabilizing factors for the shoulder
- (3) The patient must conscientiously participate in a Strengthening & training program after surgery
- (4) The surgery will compromise ROM in the hope of gaining stability
- (5) Postoperative rehabilitation after Capsular shifting procedure is protracted
- (6) In spite of the best operative & postoperative management, the success in Re-establishing normal shoulder function is substantially less than that of Traumatic Instability
- (7) The shoulder must have an identified mechanical problem for which surgery is the answer

REFERENCES

1. Bassett RW, Browne AO, Moray BF and An KN: Glenohumeral muscle force and moment mechanics in a position of shoulder instability. *J Biomech*, 23:405-415, 1990.
2. Bowen MK, Deng XH, Warner JP, et al: The effect of joint compression on stability of the glenohumeral joint. *Trans Orthop Res Soc*, 17:289, 1992.
3. Burkhart SS, De Beer JF, Barth JR, Cresswell T, Roberts C, Richards DP: Results of modified Latarjet reconstruction in patients with anteroinferior instability and significant bone loss. *Arthroscopy*, 23: 1033-1041, 2007.
4. Calandra J, Baker C and Uribe J: The incidence of Hill-sachs lesions in initial anterior shoulder dislocations. *Arthroscopy*, 5:254-257, 1989.
5. Chen AL, Hunt SA, Hawkins RJ, Zuckerman JD: Management of bone loss associated with recurrent anterior glenohumeral instability. *Am J SportsMed*, 33:912-925, 2005.
6. Chuang TY, Adams CR, Burkhart SS: Use of preoperative three dimensional computed tomography to quantify glenoid bone loss in shoulder instability. *Arthroscopy*, 24:376-382, 2008.
7. Clark J, Sidles J and Masten F: The relationship of glenohumeral joint capsule to the rotator cuff. *Clin Orthop*, 254:29-34, 1990.

8. Cole BJ, Rios CG, Mazzocca AD and Warner JP: Anatomy, biomechanics, and pathophysiology of glenohumeral instability. In: Iannotti JP and Williams GR ed. Disorders of the shoulder, 2nd ed. Philadelphia, Lippincott Williams and Wilkins:281-312, 2007.
9. Craig E: The posterior mechanism of acute anterior shoulder dislocation. Clin Orthop, 190:212-216, 1984.
10. Doucas WC and Speer KP: Anatomy, pathophysiology, and biomechanics of shoulder instability. Orthop Clin N Am, 32:381-391, 2001.
11. Emery R and Mullaji A: Glenohumeral joint instability in normal adolescents. Incidence and significance. J Bone Joint Surg, 73B:406-408, 1991.
12. Flatow E, Miller S and Neer CI: Chronic anterior dislocation of the shoulder. J shoulder Elbow Surg, 2:2-10, 1993.
13. Gerber C and Nyffeler RW: Classification of glenohumeral joint instability. Clin Orthop, 400:65-76, 2002.
14. Itoi E, Hatakeyama Y, Kido T, et al: A new method of immobilization after traumatic anterior dislocation of the shoulder: A preliminary study. J Shoulder Elbow Surg, 12:413-415, 2003.
15. Jensen KL and Rockwood CA: Glenohumeral instability: Classification, clinical assessment, and imaging. In: Norris TR ed. Orthopaedic Knowledge Update. 2nd ed. Rosemont, AAOS:65-70, 2002.
16. Lafosse L, Lejeune E, Bouchard A, Kakuda C, Gobezie R, Kochhar T: The arthroscopic Latarjet procedure for the treatment of anterior shoulder instability. Arthroscopy, 23:1242, 2007.
17. Larrain MV, Montenegro HJ, Mauas DM, Collazo CC, Pavon F: Arthroscopic management of traumatic anterior shoulder instability in collision athletes: Analysis of 204 cases with a 4- to 9-year follow-up and results with the suture anchor technique. Arthroscopy, 22: 1283-1289, 2006.
18. Latarjet M: Treatment of recurrent dislocation of the shoulder. Lyon Chir, 49:994-997, 1954.
19. Lephart S, Waner J, Borsa P, et al: Proprioception of the shoulder. Joint in healthy, unstable, and surgically repaired shoulders. J shoulder Elbow Surg, 3: 371-380, 1994.
20. Lippitt S, Harris S, Harryman DT, et al: In vivo quantification of the laxity of normal and unstable glenohumeral joints. J Shoulder Elbow Surg, 3:215-223, 1994.
21. Lippitt S, Vanderhooft J, Harris S, et al: Glenohumeral stability from concavity-compression. A quantitative analysis. J Shoulder Elbow Surg, 2:27-35, 1993
22. Matsen FA III, Chebli C, Lippitt S: Principles for the evaluation and management of shoulder instability. J Bone Joint Surg Am, 88:648-659, 2006.
23. Matsen FL, Thomas S and Rockwood CJ: Anterior glenohumeral instability. In Rockwood CJ and Matsen FL, eds. The shoulder. Philadelphia. WB Saunders, 526-622, 1990.
24. Mologne TS, Provencher MT, Menzel KA, Vachon TA, Dewing CB: Arthroscopic stabilization in patients with an inverted pear glenoid: Results in patients with bone loss of the anterior glenoid. Am J Sports Med, 35:1276-1283, 2007.
25. O'Brien S, Arnoczky S, Warren R et al: Developmental anatomy of the shoulder and anatomy of the glenohumeral joint. In Rockwood CJ and Matsen FL, eds. The shoulder. Philadelphia, WB Saunders, 1-33, 1990.
26. O'Brien SJ, Neves MC, Arnoczky SP, et al: The anatomy and histology of the inferior glenohumeral ligament complex of the shoulder. Am J Sports Med, 18:449-456, 1990.
27. O'Driscoll S and Evans D: Contralateral shoulder instability following anterior repair. An epidemiological investigation. J Bone Joint Surg, 73B: 941-946, 1991.
28. Phillips BB: Recurrent dislocations. In: Canale ST ed. Campbell's operative orthopaedics. 10th ed. St.

- Louis, Mosby-yearbook Inc:2377-2448, 2003.
29. Reeves B: Arthrography in acute dislocation of the shoulder. *J Bone Joint Surg*, 48B: 182, 1968.
 30. Reeves B: Experiments on tensile strength of the anterior capsular structures of the shoulder in man. *J Bone Joint Surg*, 50B: 858-865, 1968.
 31. Rhee YG: The shoulder: diagnosis and treatment. 1st ed. Seoul, Young-chang medical books:65-153, 2003.
 32. Rowe C and Sakellarides H: Factors related to recurrences of the anterior dislocation of the shoulder. *Clin Orthop*, 20:40-48, 1961.
 33. Rowe C, Pierce D and Clark J: Voluntary dislocation of the shoulder. A preliminary report on clinical, eletromyographic, and psychiatric study of twenty-six patients. *J Bone Joint Surg*. 55A:445-460, 1973.
 34. Spencer EE Jr, Brems JJ: A simple technique for management of locked posterior shoulder dislocations: Report of two cases. *J Shoulder Elbow Surg*, 14:650-652, 2005.
 35. Sugaya H, Kon Y, Tsuchiya A: Arthroscopic repair of glenoid fractures using suture anchors. *Arthroscopy*, 21:635, 2005.
 36. Sugaya H, Moriishi J, Kanisawa I, Tsuchiya A: Arthroscopic osseous Bankart repair for chronic recurrent traumatic anterior glenohumeral instability. *J Bone Joint Surg Am*, 87:1752-1760, 2005.
 37. Turkel S, Panio M, Marshall, J et al: Stabilizing mechanisms preventing anterior dislocation of the glenohumeral joint. *J Bone Joint Surg*, 63A:1208-1217, 1981.
 38. Warner J, Deng X, Warren R, et al: Superior-inferior translation in the intact and vented glenohumeral joint. *J Shoulder Elbow Surg*, 2:99-125, 1993.
 39. Warner JJ, Gill TJ, O' Hollerhan JD, Pathare N, Millett PJ: Anatomical glenoid reconstruction for recurrent anterior glenohumeral instability with glenoid deficiency using an autogenous tricortical iliac crest bone graft. *Am J Sports Med*, 34: 205-212, 2006.
 40. Warner JJ, Micheli LJ, Arslanian LE, et al: Scapulothoracic motion in normal shoulders and shouders with glenohumeral instability and impingement syndrome. A study using Moire tomographic analysis. *Clin Orthop*, 285:191-199, 1992.
 41. Warren, Kornblatt I and Marchand R: Static factors affecting posterior shoulder stability. *Orthop Trans*, 8:89, 1984.
 42. Weulker N, Brewes F and Sperveslage C: Passive glenohumeral joint stabilization. A biomechanical study. *J Shoulder Elbow Surg*, 3:129-134, 1994.
 43. Wolf E, Cheng J and Dickson K: Humeral avulsion of glenohumeral ligaments as cause of anterior shoulder instability. *Arthroscopy*, 11:600-607,1995.