

[초CD-01] Recent progress in dark energy research

Chan-Gyung Park

*Division of Science Education and Institute of Fusion Science,
Chonbuk National University*

Astronomical observations strongly suggest that the expansion rate of our universe is currently under acceleration. The nature of the so-called dark energy causing the acceleration is unknown, and it is one of the fundamental mysteries in the present day theoretical cosmology. Here we briefly review the current state of cosmic dark energy research in both theoretical and observational sides. Constraints on dynamical dark energy models (e.g., w -fluid, quintessence, and modified gravity) with recent observational data from type Ia supernovae, cosmic microwave background radiation, and large-scale structures in the universe indicate a preferred direction toward the simplest Λ CDM world model. We also discuss some issues regarding the early dark energy model and the spherical collapse of matter in the presence of dark energy.

[구CD-02] 21 cm signal from highly clustered Population III and Population II objects at high redshift

Kyungjin Ahn¹, Hao Xu², Michael Norman², Marcelo Alvarez³, John Wise⁴,
*¹Chosun University, ²University of California at San Diego, ³Canadian Institute for
Theoretical Astrophysics, ⁴Georgia Institute of Technology*

We present a prediction for 21cm differential brightness temperature (dT_b) from a set of strongly clustered sources of Population III and II objects at high redshift, from a suite of numerical simulations of their formation and radiative processes. These objects are located inside a highly biased density environment ("Rarepeak"), which is a rare, high-density peak which extends to ~ 7 comoving Mpc. We study the impact on the resulting 21 cm signal from their ultraviolet and X-ray properties. The boost of emission ($dT_b > 0$) by high-density environment, moderate leakage of X-ray photons, and strong absorption due to Lyman-alpha pumping contrive to make Rarepeak a discernible, spatially-extended (sky angle $\sim 10'$) object around $z \sim 15$, which is found to be detectable as a single object by Square Kilometre Array (SKA) with integration time of $\sim [600-2000]$ hours. We also examine detectability of many such peaks through SKA precursors.