

galactic star formation models are challenged. To make quantitative analysis of the ISM in the galaxy, we apply 2D cross-correlation technique to the multi-wavelength data for the first time. By cross-correlating different tracers of star formation, dust and gas phases in IC 10 in a two dimensional way, we discuss the gas properties and star formation history of the galaxy.

[7 GC-12] Quenching of star formation in massive halos at $z \sim 2$

Raphael Gobat
KIAS

The gradual infall of small dark matter halos onto larger ones has become a relatively straightforward aspect of the standard hierarchical formation paradigm. What happens to the baryons they contain, however, is less well understood. Of special relevance are the processes that regulate and ultimately suppress star formation in galaxies in the early universe.

The $z=1.5-2.5$ epoch is then particularly interesting as a transition period when global star-formation in the universe starts peaking but also where the first ostensibly collapsed and virialized galaxy clusters appear, along with segregated galaxy populations. From a theoretical point of view, the mode of gas accretion in massive halos is also expected to change around this time, switching from a cold to a hot phase and affecting the build-up and evolution of the galaxies they host.

A lot of effort has thus been devoted to the search for high-redshift structures, in particular galaxy clusters, through a variety of methods. However, as the limited area for which deep datasets are available remains relatively limited, only few massive $z > 1.5$ structures have been found so far. Here I will instead discuss the regulation of star-formation in lower-mass, X-ray detected halos at $z \sim 2$ and its implication for galaxy quenching at high redshift. As these smaller, group-size halos are vastly more abundant and structurally simpler than massive clusters, they allow for true statistical studies and offer a novel way to probe environmental effects in this transitional epoch.

[7 GC-13] The Key role of the Bulge Compactness in Star-forming Activity in Late-type Galaxies

Woong-bae Jee in Suk-Jin Yoon

Department of Astronomy and Center for Galaxy Evolution Research, Yonsei University, Seoul 120-749, Republic of Korea

Which mechanism governs star-formation activity in galaxies is still one of the most important, open questions in galactic astronomy. To address this issue, we investigate the specific star formation rate (sSFR) of late-type galaxies as functions of various structural parameters including the morphology, mass, radius, and mass compactness (MC). We use a sample of $\sim 200,000$ late-type galaxies with $z = 0.02 \sim 0.2$ from SDSS DR7 and a catalog of bulge-disk decomposition (Simard et al. 2011; Mendel et al. 2013). We find a remarkably strong correlation between bulge's MC and galaxy's sSFR, in the sense that galaxies with more compact bulge tend to be of lower sSFR. This seems counter-intuitive given that galactic sSFR is driven predominantly by disks rather than bulges and suggests that the central mass density plays a key role in recent star-forming activity. We discuss the physical cause of the new findings in terms of the bulge growth history and AGN activities.

[7 GC-14] Revealing the complexity of ionized gas outflows in powerful Type 2 AGN in the local Universe

Marios Karouzos¹, Jong-Hak Woo¹, Hyun-Jin Bae^{1,2}
¹Department of Physics and Astronomy, Seoul National University, ²Department of Astronomy, Yonsei University

There exist scaling relations that link the mass of supermassive black holes with both the velocity dispersion and the mass of the central stellar cusp of their host galaxies. This implies that these two components grow in tandem. Feedback from actively accreting supermassive black holes (AGN), in the form of multi-phase gas outflows, has been argued to be the agent of this co-evolution. Here we employ the powerful GMOS integral field spectroscopy unit on the 8.2m Gemini-North telescope to investigate ionized gas outflows of luminous Type 2 AGN in the local Universe ($z < 0.1$). Our sample of 6 galaxies is drawn from the Sloan Digital Sky Survey (SDSS) and was selected based on their [OIII] dust-corrected luminosity (> 1042 erg/s) and signatures of outflows in the [OIII] line profile of their spatially integrated SDSS spectra. These are arguably the best candidates to explore AGN feedback in action since they are $< 1\%$ of a large local type 2 AGN SDSS sample selected based on their [OIII] kinematics. We combine a careful