

creates a central cavity and a compressed shell at the ionized boundary. We analytically solve for the temporal evolution of a thin shell, finding a good agreement with the numerical experiments. We estimate the minimum star formation efficiency required for a cloud of given mass and size to be destroyed by an HII region expansion. We find that typical giant molecular clouds in the Milky Way can be destroyed by the gas-pressure driven expansion of an H II region, requiring an efficiency of less than a few percent. On the other hand, more dense cluster-forming clouds in starburst environments can be destroyed by the radiation pressure driven expansion, with an efficiency of more than  $\sim 30$  percent that increases with the mean surface density, independent of the total (gas+stars) mass. The time scale of the expansion is always smaller than the dynamical time scale of the cloud, suggesting that H II regions are likely to be a dominant feedback process in protoclusters before supernova explosions occurs.

### [7 IM-03] Efficient simulation method for a gas inflow to the central molecular zone

Jihye Shin<sup>1</sup>, Sungsoo S. Kim<sup>2,3</sup>, Junichi Baba<sup>4</sup>, Takayuki R. Saitoh<sup>4</sup>, Kyungwon Chun<sup>2</sup>, and Shunsuke Hozumi<sup>5</sup>

<sup>1</sup>*Kavli Institute for Astronomy and Astrophysics at Peking University, P.R. China,*

<sup>2</sup>*School of Space Research, Kyung Hee University, Republic of Korea,*

<sup>3</sup>*Department of Astronomy & Space Science, Kyung Hee University, Republic of Korea,*

<sup>4</sup>*Earth-Life Science Institute, Tokyo Institute of Technology, Japan,*

<sup>5</sup>*Faculty of Education, Shiga University, Japan*

We present hydrodynamic simulations of gas clouds that inflowing from the disk to a few hundred parsec region of the Milky Way. Realistic Galactic structures are included in our simulations by thousands of multipole expansions that describe 6.4 million stellar particles of a self-consistent Galaxy simulation (Baba, Saitoh & Wada, in prep.). We find that a hybrid multipole expansion model with two different basis sets and a thick disk correction well reproduces the overall structures of the Milky Way. We find that the nuclear ring evolves into 240 pc at  $T \sim 1500$  Myr, regardless of the initial size. For most of simulation runs, gas inflow rate to the nuclear region is equilibrated as  $\sim 0.02$  Msun/yr, and thus accumulated gas mass and star formation activity is stabilized as  $6 \times 10^7$  Msun and  $\sim 0.02$  M/yr, respectively. These stabilized values are in a good agreement with

estimations for the CMZ. The nuclear ring is off-centered to the Galactic center by the lopsided central mass distribution of the Galaxy model, and thus an asymmetric mass distribution is arose accordingly. The lopsidedness also leads the nuclear ring to be tilted to the Galactic plane and to precess along the Galaxy rotation. In early evolutionary stage when gas clouds start to inflow and form the nuclear ring, the z-directional oscillations of the gas clouds results in the twisted, infinity-shaped nuclear ring. Since the infinity-shaped feature is transient only for first 100 Myr, the current infinity-shape observed in the CMZ may indicate that the CMZ forms quite recently.

### [7 IM-04] Simultaneous observations of the H<sub>2</sub>O and SiO masers toward the late-type stars using KVN

Youngjoo Yun, Se-Hyung Cho, Jaeheon Kim, Yoon Kyung Choi, and the members of evolved star sub-working group

*Korea Astronomy and Space Science Institute, 776 Daedeok-daero, Yuseong, Daejeon 305-348, Korea*

We present the results of simultaneous observations of the H<sub>2</sub>O and SiO masers emitted from the circumstellar envelopes of the late-type stars. These observations have been carried out at the K and Q bands using KVN since 2014 August and were scheduled to test the feasibility of multi-frequency phase referencing analysis on the maser lines. In order to increase the accuracy of group delay solution in the fringe search on the continuum source, the IF channels were randomly distributed within the available bandwidth of 500 MHz in each band. The positions of all maser spots are relatively described with respect to the position of the reference continuum source through the source frequency phase referencing technique, and this provides the astrometric position accuracy. Therefore, the relative locations of the H<sub>2</sub>O maser spots with respect to the SiO maser spots are determined from our observations, and the capability of the simultaneous multi-band observation of KVN is proved to be powerful to study the maser pumping mechanism around the late-type stars.

### [7 IM-05] Clustering properties and halo occupation of Lyman-break galaxies at $z \sim 4$

Jaehong Park<sup>1</sup>, Han-Seek Kim<sup>1</sup>, Stuart B. Wyithe<sup>1</sup>, Cedric G. Lacey<sup>2</sup>, and Carlton M. Baugh<sup>2</sup>

<sup>1</sup>*School of Physics, The University of Melbourne,*