Spin-Orbit Torque and Beyond

See-Hun Yang^{*} IBM Almaden Research Center, San Jose, CA 95120, USA ^{*}E-mail: seeyang@us.ibm.com

Efficient manipulation of magnetic elements are one of the main goals in spintronics. Discovery of spin transfer torque two decades ago has made a major impact on magnetic community and industry thus accelerating development of more powerful devices, i.e. STT-MRAM [1]. In particular, over the past few years a new type of torque, so-called spin-orbit torque has emerged and promised even more exciting advent of new type of devices in the future [2,3]. In this talk I will review the recent advancement of spin-orbit torque driven domain wall motion in magnetic nanowires and present other novel torques beyond spin-orbit torque [4,5,6].

References

- [1] Stuart Parkin and See-Hun Yang, "Memory on the Racetrack", Nature Nanotechnology 10, 195-198 (2015).
- [2] Kwang-Su Ryu, Luc Thomas, See-Hun Yang, and Stuart Parkin, "Chiral Spin Torque at Magnetic Domain Walls", Nature Nanotechnology 8, 527-533 (2013).
- [3] Satoru Emori, Uwe Bauer, Sung-Min Ahn, Eduardo Martinez, and Geoffrey S. D. Beach, "Current-driven dynamics of chiral ferromagnetic domain walls", Nature Materials **12**, 611-616 (2013).
- [4] See-Hun Yang, Kwang-Su Ryu, and Stuart Parkin, "Domain-wall velocities of up to 750 m s⁻¹ driven by exchange-coupling torque in synthetic antiferromagnets", Nature Nanotechnology **10**, 221-226 (2015).
- [5] Chirag Garg, See-Hun Yang, Timothy Phung, Aakash Pushp, and Stuart Parkin, submitted.
- [6] See-Hun Yang, Chirag Garg, and Stuart Parkin, submitted.