

to 0.6 and those of Shue et al. (1998) are slightly higher than those of the other models. Fourth, the predicted magnetopause radius based on Lin et al.(2010) well match the observed one within one earth radius, while that on Shue et al. (1998) overestimate the observed one by about 2 earth radii. Fifth, the PoD and CSI values of all the models are better for the solar maximum phase than those for the other phases, implying that the models are more optimized for the phase.

[구 SS-03] Collisionless Magnetic Reconnection and Dynamo Processes in a Spatially Rotating Magnetic Field

Junggi Lee , G. S. Choe, and Inhyeok Song
Kyung Hee University

Spatially rotating magnetic fields have been observed in the solar wind and in the Earth's magnetopause as well as in reversed field pinch (RFP) devices. Such field configurations have a similarity with extended current layers having a spatially varying plasma pressure instead of the spatially varying guide field. It is thus expected that magnetic reconnection may take place in a rotating magnetic field no less than in an extended current layer. We have investigated the spontaneous evolution of a collisionless plasma system embedding a rotating magnetic field with a two-and-a-half-dimensional electromagnetic particle-in-cell (PIC) simulation. In magnetohydrodynamics, magnetic flux can be decreased by diffusion in O-lines. In kinetic physics, however, an asymmetry of the velocity distribution function can generate new magnetic flux near O- and X-lines, hence a dynamo effect. We have found that a magnetic-flux-reducing diffusion phase and a magnetic-flux-increasing dynamo phase are alternating with a certain period. The temperature of the system also varies with the same period, showing a similarity to sawtooth oscillations in tokamaks. We have shown that a modified theory of sawtooth oscillations can explain the periodic behavior observed in the simulation. A strong guide field distorts the current layer as was observed in laboratory experiments. This distortion is smoothed out as magnetic islands fade away by the O-line diffusion, but is soon strengthened by the growth of magnetic islands. These processes are all repeating with a fixed period. Our results suggest that a rotating magnetic field configuration continuously undergoes deformation and relaxation in a short time-scale although it might look rather steady in a long-term view.

[구 SS-04] Dependence of spacecraft anomalies at different orbits on energetic electron and proton fluxes

Kangwoo Yi¹, Yong-Jae Moon^{1,2}, Ensang Lee² and Jae-Ok Lee¹
¹*School of Space Research, Kyung Hee University, Korea,* ²*Astronomy & Space Science, Kyung Hee University, Korea*

In this study we investigate 195 spacecraft anomalies from 1998 to 2010 from Satellite News Digest (SND). We classify these data according to types of anomaly : Control, Power, Telemetry etc. We examine the association between these anomaly data and daily peak particle (electron and proton) flux data from GOES as well as their occurrence rates. To determine the association, we use two criteria that electron criterion is >10,000 pfu and proton criterion is >100 pfu. Main results from this study are as follows. First, the number of days satisfying the criteria for electron flux has a peak near a week before the anomaly day and decreases from the peak day to the anomaly day, while that for proton flux has a peak near the anomaly day. Second, we found a similar pattern for the mean daily peak particle (electron and proton) flux as a function of day before the anomaly day. Third, an examination of multiple spacecraft anomaly events, which are likely to occur by severe space weather effects, shows that anomalies mostly occur either when electron fluxes are in the declining stage, or when daily proton peak fluxes are strongly enhanced. This result is very consistent with the above statistical studies. Our results will be discussed in view of the origins of spacecraft anomaly.

[구 SS-05] Prediction of Long-term Solar Activity based on Fractal Dimension Method

Rok-Soon Kim^{1,2}
¹*Korea Astronomy and Space Science Institute,* ²*University of Science and Technology*

Solar activity shows a self-similarity as it has many periods of activity cycle in the time series of long-term observation, such as 13.5, 51, 150, 300 days, and 11, 88 years and so on. Since fractal dimension is a quantitative parameter for this kind of an irregular time series, we applied this method to long-term observations including sunspot number, total solar irradiance, and 3.75 GHz solar radio flux to predict the start and maximum times as well as expected maximum sunspot number for