

the ambient medium, and are related to the gamma-ray flare which has been detected with VHE gamma-ray telescopes such as MAGIC and VERITAS.

[ㄱ KVN-02] LINEAR POLARIZATION OF CLASS I METHANOL MASERS IN MASSIVE STAR-FORMING REGIONS

Ji-hyun Kang¹, Do-Young Byun¹, Kee-Tae Kim¹, Jongsoo Kim¹, A-Ran Lyo¹, Mi-Kyung Kim¹, and W. H. T. Vlemmings²

¹*Korea Astronomy and Space Science Institute*

²*Onsala Space Observatory*

Class I methanol masers are found to be good tracers of the interaction between outflows from massive young stellar objects with their surrounding media. Although polarization observations of Class II methanol masers have been able to provide information about magnetic fields close to the central (proto)stars, polarization observations of Class I methanol masers are rare, especially at 44 and 95GHz. We present the results of linear polarization observations of 39 Class I methanol maser sources at 44 and 95GHz. These two lines are observed simultaneously with one of the 21m Korean VLBI Network telescopes in single-dish mode. Approximately 60% of the observed sources have fractional polarizations of a few percent in at least one transition. This is the first reported detection of linear polarization of the 44GHz methanol maser. We also observed 7 targets with the KVN in VLBI mode. We will present its preliminary results, too.

[ㄱ KVN-03] The Power of Simultaneous Multi-frequency Observations for mm-VLBI: Beyond Frequency Phase Transfer

Guang-Yao Zhao¹, Juan Carlos Algaba¹, Sang Sung Lee^{1,2}, Taehyun Jung^{1,2}, Richard Dodson³, Maria Rioja^{3,4,5}, Do-Young Byun^{1,2}, Jeffrey Hodgson¹, Sincheol Kang^{1,2}, Dae-Won Kim⁶, Jae-Young Kim⁷, Jeong-Sook Kim⁸, Soon-Wook Kim^{1,2}, Motoki Kino^{1,8}, Atsushi Miyazaki^{1,9}, Jong-Ho Park⁶, Sascha Trippe⁶, Kiyooki Wajima¹

¹*KASI*, ²*UST*, ³*ICRAR*, ⁴*CSIRO*, ⁵*OAN (IGN)*, ⁶*SNU*, ⁷*MPIFR*, ⁸*NAOJ*, ⁹*Hosei Univ*

Atmospheric propagation effects at millimeter wavelengths can significantly alter the phases of radio signals and reduce the coherence time, putting tight constraints on high frequency Very Long Baseline Interferometry (VLBI) observations.

In previous works it has been shown that non-dispersive (e.g. tropospheric) effects can be calibrated with the frequency phase transfer (FPT) technique. The coherence time can thus be significantly extended. Ionospheric effects, which can still be significant, remain however uncalibrated after FPT, as well as the instrumental effects. In this work, we implement a further phase transfer between two FPT residuals (i.e. so-called FPT2) to calibrate the ionospheric effects based on their frequency dependence. We show that after FPT2, the coherence time at 3 mm can be further extended beyond 8 hours, and the residual phase errors can be sufficiently canceled by applying the calibration of another source, which can have a large angular separation from the target ($> 20^\circ$). Calibrations for all-sky distributed sources with a few calibrators are also possible after FPT2. One of the strengths and uniqueness of this calibration strategy is the suitability for high frequency all-sky survey observations including very weak sources. We discuss the introduction of a pulse calibration system in the future to calibrate the remaining instrumental effects and allowing the possibility of imaging the source structure at high frequencies with FPT2, where all phases are fully calibrated without involving any sources other than the target itself.

[ㄱ KVN-04] Simultaneous VLBI observations of H₂O and SiO masers toward VX Sgr using KVN

Dong-Hwan Yoon^{1,2}, Se-Hyung Cho², Youngjoo Yun², Yoon Kyung Choi², Maria Rioja³, Richard Dodson³, Jaeheon Kim⁴, Dongjin Kim⁵, Hanul Yang¹, Hiroshi Imai⁶, Do-Young Byun²

¹*Astronomy program, Department of Physics and Astronomy, Seoul National University, Korea.*

²*Korea Astronomy and Space Science Institute, Korea.*

³*International Center for Radio Astronomy Research, M468, The University of Western*

Australia, Australia.

⁴*Shanghai Astronomical Observatory, Chinese Academy of Sciences, China.*

⁵*Department of Astronomy, Yonsei University, Korea.*

⁶*Department of Physics and Astronomy, Kagoshima University, Japan*

We performed simultaneous VLBI observations of H₂O 616–523 (22.2 GHz) and SiO $v=1, 2, J=1-0$ (43.1, 42.8 GHz) and $v=1, J=2-1, J=3-2$ (86.2, 129.3 GHz) masers toward VX Sagittarius using the Korean VLBI Network (KVN). The astrometrically registered maps of the 22.2 GHz H₂O and 43.1, 42.8, 86.2 SiO masers were successfully obtained at two epochs of 2016 February 27 and 2016 March

27 by adopting the Source Frequency Phase Referencing (SFPR) method. In addition we detected 129.3 GHz SiO maser at second epoch. These results make it possible to determine the accurate position of central star as a dynamical center of 22.2 GHz H₂O maser and relative locations of 43.1, 42.8, 86.2, 129.3 GHz SiO masers. In addition, it is possible to investigate the morphological and kinematic variations of clumpy structures from SiO maser to H₂O maser regions in future together with the development of asymmetric structure of H₂O maser region.

measured distance to R Crt enables us to estimate the actual 3D velocity of water masers around R Crt. Our research suggests the possibility of performing astrometric studies with the KVN. As a next step, we are going to enhance the astrometric accuracy by observing SiO masers.

[구 KVN-05] The recent activities for a precise astrometry using SFPR with KVN

Ilje Cho^{1,2}, Taehyun Jung^{1,2}, Bongwon Sohn^{1,2}, Motoki Kino^{1,3}, Guangyao Zhao¹, Maria Jose Rioja⁴, Richard Dodson⁴, Ivan Agudo⁵

¹*Korea Astronomy and Space Science Institute*

²*University of Science and Technology*

³*National Astronomical Observatory of Japan*

⁴*International Centre for Radio Astronomy Research*

⁵*Instituto de Astrofísica de Andalucía*

Thanks to the quasi-optics system of KVN, the relative astrometry at different frequencies can be effectively achieved. In this talk, as a part of the KVN special session, we are going to present an important method making it possible, so called the source frequency phase referencing (SFPR).

Based on the background, we will show our recent activities using the SFPR for the Galactic Center (Sgr A*) study, from its advantages on astrometric studies to the remained issues.

[구 KVN-06] Measurement of proper motion and annual parallax with maser emission

Dong-Jin Kim¹, Se-Hyung Cho², Young-Joo Yun², Yoon Kyung Choi², Dong-Hwan Yoon², Suk-Jin Yoon¹

¹*Department of Astronomy, Yonsei University,*

²*Korea Astronomy and Space Science Institute*

We present the results of astrometric observations using water masers around a semi-regular variable star R Crt. The observations were carried out for two years with the Korean VLBI Network (KVN). The absolute positions of the water masers from R Crt are successfully obtained at 10 epochs in total. By tracking the positions of the water maser emission, we directly measured the annual parallax and distance of R Crt. The