

[포 IM-06] Spitzer and Herschel observations of protostellar outflows in L1251B

Yunhee Choi¹, Jeong-Eun Lee¹, Joel D. Green², Sébastien Maret³, Yau-Lun Yang⁴, Edwin A. Bergin⁵, Geoffrey A. Blake⁶, Abraham. C. A. Boogert⁷, James Di Francesco^{8,9}, Neal J. Evans II⁴, Klaus M. Pontoppidan², and Annelia I. Sargent⁶
¹*Kyung Hee University, Korea*, ²*Space Telescope Science Institute, USA*, ³*Institut de Planétologie et d'Astrophysique de Grenoble, France*, ⁴*University of Texas at Austin, USA*, ⁵*University of Michigan, USA*, ⁶*California Institute of Technology, USA*, ⁷*NASA Ames Research Center, USA*, ⁸*National Research Council, Canada*, ⁹*University of Victoria, Canada*

L1251B is an excellent example of a small group of pre- and protostellar objects in low-mass star-forming region. Previous interferometer data with a high angular resolution resolved the outflows associated with L1251B into a few components. To understand the physical conditions of the protostellar outflows in L1251B, we mapped this region spectroscopically with Spitzer/IRS and obtained spectral line data from Herschel/PACS. Spitzer/IRS provides the S(0)-S(7) pure rotational lines of H₂ as well as fine-structure emissions produced in shocks such as S, [Si II], and [Fe II] and it is a powerful tool for studying shocked interstellar gas. In addition, [O I] lines observed with Herschel/PACS are described well by J-type shock models expected in the outflows from protostars. We will present an analysis of the L1251B protostellar outflow in the H₂ pure rotational lines and fine-structure emissions.

[포 IM-07] Polarized Infrared Emission from Polycyclic Aromatic Hydrocarbons and Implications

Thiem Hoang
Korea Astronomy and Space Science Institute & Korea University of Science and Technology

Polarized mid-infrared emission from polycyclic aromatic hydrocarbons (PAHs) can provide a crucial test of basic physics of alignment of nanoparticles and opens a potential new window into studying magnetic fields. In this talk, I will present a new model of polarized PAH emission that takes into account the effect of PAH alignment with the magnetic field due to resonance paramagnetic relaxation. I will then present our

predictions for the polarization level of the strong PAH emission features from the interstellar medium. I will present the first detection of polarization of PAH emission at 11.3micron which is consistent with our theoretical prediction. Finally, I will discuss important implications of this work for tracing magnetic fields via mid-IR PAH features and for constraining the polarization of anomalous microwave emission that is useful for the quest of CMB B-modes.

태양/태양계

[포 SS-01] Connection between a Small-Scale Emerging Island and Double Arc Loops Producing a M6.5 Flare in Active Region 12371

Jihye Kang¹, Satoshi Inoue², Yong-Jae Moon³, Tetsuya Magara³ and Kanya Kusano²
¹*Department of Astronomy and Space Science, Kyung Hee University*, ²*Institute for Space-Earth Environmental Research, Nagoya University*, ³*School of Space Research, Kyung Hee University*

In this paper, we report a small-scale emerging island and double arc loops, which are associated with a M6.5 flare, in Active Region 12371. We investigate the spatial and temporal changes of both photospheric magnetic fields using SDO/HMI data and coronal magnetic structures obtained from nonlinear force-free field (NLFFF) extrapolation. From the vector magnetograms, we find a small-scale emerging island near the main polarity inversion line about three hours before the flare. The island has a strong shear angle, which is determined by difference between transverse component of observed field and potential field, of around 90 degrees. Furthermore, the NLFFF well reproduces a sigmoidal structure seen in SDO/AIA 94, which is consistent with the double arc loops configuration suggested by Ishiguro and Kusano (2017) who introduced a magnetic configuration showing the double arc instability. The observed emerging island is located among the double arc loops, which is also supported by their model. Finally, there was an eruption (M6.5 flare) associated with the loops. We discuss a possible role of the double arc instability for the eruption.

[포 SS-02] Velocity Oscillations in the

Chromosphere and the Transition Region above Plage Regions

Hannah Kwak, Jongchul Chae
Astronomy Program, Department of Physics & Astronomy, Seoul National University

We investigate velocity oscillations in the active region plage by using the high-spatial, high-spectral and high-temporal resolution spectral data acquired by the Interface Region Imaging Spectrograph (IRIS). From the Mn I 2801.907 Å (lower chromosphere), C II (lower transition region) and Si IV (middle transition region) lines, we measure the line of sight Doppler velocity at different atmospheric layers, and present results of wavelet analysis of the plage region with a range of periods from 2 to 8 minutes. In addition, we present correlations of the oscillations from the lower chromosphere to the middle transition region. Finally, we will discuss the regional dependence of the oscillation properties on physical properties such as temperature and magnetic field inclination.

[ㄷ SS-03] Connection of Blobs along Post-CME Ray and EUV Flares

Yoojung Kim, Jongchul Chae
Seoul National University

After a coronal mass ejection occur, plasma blobs are often observed along the post-CME ray. Searching for features related to the plasma blobs would be important in understanding their origin. We investigated the morphology of solar flares at EUV wavelengths, around the estimated times when blobs were formed. We focused on three events - 2013 September 21 and 22, 2015 March 7 and 8, and 2017 July 13 and 14 - observed by Atmospheric Imaging Assembly (AIA) aboard Solar Dynamic Observatory (SDO). Around the blob ejection times on 2013 September 21 and 22 and 2017 July 13 and 14, we found regions with recurrent events of pronounced flux increase in EUV images. Around those of 2015 March 7 and 8, however, we could not observe such recurrent flux increase. This illustrates that even though blob ejections along different post-CME rays look similar in the high corona, the associated features in the low corona may differ. We conclude that magnetic morphology and CME triggering process should be carefully examined in order to classify plasma blobs by their nature.

[ㄷ SS-04] Current Status of

KMTNet/DEEP-South Collaboration Research for Comets and Asteroids Research between SNU and KASI

Yoonsoo P. BACH¹, Hongu YANG^{1,2}, Yuna G. KWON¹, Subin LEE¹, Myung-Jin KIM², Young-Jun CHO^{2,3}, Jintae Park², Masateru ISHIGURO¹, and Hong-Kyu Moon²

¹*Department of Physics and Astronomy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea,*

²*Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 34055, Korea*

³*Astronomy and Space Science, University of Science and Technology, 217 Gajeong-ro, Yuseong-gu, Daejeon 34113, Korea*

Korea Microlensing Telescope Network (KMTNet) is one of powerful tools for investigating primordial objects in the inner solar system in that it covers a large area of the sky (2×2 degree²) with a high observational cadence. The Deep Ecliptic Patrol of the Southern sky (DEEP-South) survey has been scanning the southern sky using KMTNet for non-bulge time (45 full nights per year) [1] since 2015 for examining color, albedo, rotation, and shape of the solar system bodies. Since 2017 January, we have launched a new collaborative group between Korea Astronomy and Space Science Institute (KASI) and Seoul National University (SNU) with support from KASI to reinforce mutual collaboration among these institutes and further to enhance human resources development by utilizing the KMTNet/DEEP-South data. In particular, we focus on the detection of comets and asteroids spontaneously scanned in the DEEP-South for (1) investigating the secular changes in comet's activities and (2) analyzing precovery and recovery images of objects in the NASA's NEOWISE survey region. In this presentation, we will describe our scientific objectives and current status on using KMTNet data, which includes updating the accuracy of the world coordinate system (WCS) information, finding algorithm of solar system bodies in the image, and doing non-sidereal photometry.

[ㄷ SS-05] Dependence of the peak fluxes of solar energetic particles on CME parameters and magnetic connectivity

Jinhye Park¹, Yong-Jae Moon^{1,2}, Harim Lee², S. W. Kahler³

¹*Department of Astronomy & Space Science, Kyung Hee University*