

symbiotic stars are known to exhibit broad wings around Balmer emission line. We show high resolution spectra of S-type symbiotic stars, Z Andromedae and AG Draconis, obtained with the ESPaDOnS and the 3.6 m Canada-France- Hawaii Telescope, in which we find prominent broad wings around Balmer lines. We adopt Monte-Carlo technique to consider two types of wing formation mechanisms, which are Thomson scattering by free electron in H II region and Raman scattering by atomic hydrogen in H I region. We find that Thomson wings of H α and H β have the same widths in the Doppler space due to the cross section independent of wavelength. In contrast, Raman H α wings are 3 times broader widths than H β counterparts, which is attributed to the different cross sections and branching ratios. Our CFHT data show that H α wings of Z Andromedae and AG Draconis are broader than H β wings, lending strong support to the Raman scattering origin of Balmer wings in these objects.

[포 SA-05] Chemical Abundance Analysis of Ultra Metal-Poor ([Fe/H] < -4.0) Stars

Mi Ji Jeong¹, Young Sun Lee²
¹*Department of Astronomy, Space Science, and Geology, Chungnam National University, Daejeon 34134, South Korea.* ²*Department of Astronomy and Space Science, Chungnam National University, Daejeon 34134, South Korea*

We present preliminary results of elemental abundances of six ultra-metal poor (UMP; [Fe/H] < -4.0) stars derived from high-resolution spectra obtained by Gemini/GRACES. The UMP candidates were selected for the high-resolution follow-up from the low-resolution spectra of Sloan Digital Sky Survey (SDSS). We investigate possible progenitors of the UMP objects by comparing the measured abundance patterns with yields that various supernova models predict. Our results can provide stringent constraints on the mass range of the first generation of stars, which are the progenitors of the UMP objects.

[포 SA-06] Magellan High Resolution Spectroscopy of Raman-Scattered He II, C II and O VI Lines in the Symbiotic Nova RR Telescopii

Jeong-Eun Heo¹, Hee-Won Lee¹, Francesco Di Mille², Tali Palma³ and Rodolfo Angeloni⁴
¹*Sejong University, Seoul, Korea,* ²*Las Campanas Observatory, La Serena, Chile,* ³*Universidad de*

Andrés Bello, Santiago, Chile, ⁴*Universidad de La Serena, La Serena, Chile*

RR~Telescopii is a symbiotic nova exhibiting accretion activities through gravitational capture of the slow stellar wind from a Mira variable. We present high resolution spectra of RR~Tel obtained with MIKE and the 6.5 m Magellan-Clay telescope, in which we find broad features with FWHM exceeding 10 Å at 6545, 6825, 7082, 7023 and 7053 Å. They are formed through Raman-scattering with atomic hydrogen of far-UV He II 1025, O~VI 1032, 1038 Å and C II 1036 and 1037 Å. We compute the Raman conversion efficiencies using the case B recombination theory for He II emissions, which are used in turn to infer the intrinsic line luminosities of O VI and C II. The Raman O~VI features are characterized by double-peaked profiles with a peak separation ~ 60km/s, pointing out the presence of an accretion disk with a physical size of ~ sub AU. In contrast, Raman C II features exhibit profiles with a simple peak and a narrower width ~40 km/s, indicating that C II is formed in a much more extended region. The weak C II multiplet at 1335, 1336 Å found in the IUE spectral archive and the absence of C II 1036, 1037 Å in the FUSE archive show that far-UV C II lines suffer heavy interstellar extinction consistent with the distance of ~ 2.5 kpc to RR Tel.

천문우주관측기술

[포 AT-01] Flight model development of the NISS structure for NEXTSat-1 payload

Bongkon Moon¹(문봉곤), Kyeongyeon Ko^{1,2}(고경연), Duk-Hang Lee¹(이덕행), Woong-seob Jeong^{1,2}(정웅섭), Sung-Joon Park¹(박성준), Dae-Hee Lee^{1,2}(이대희), Jeonghyun Pyo¹(표정현), Won-Kee Park¹(박윤희), Il-Joong Kim¹(김일중), Youngsik Park¹(박영식), Mingyu Kim³(김민규), Ukwon Nam¹(남옥원), Minjin Kim^{1,2}(김민진), Jongwan Ko¹(고종완), Myungshin Im⁴(임명신), Hyung Mok Lee⁴(이형목), Jeong-Eun Lee⁵(이정은), Goo-Hwan Shin⁶(신구환), Jangsoo Chae⁶(채장수), Toshio Matsumoto^{1,7}
¹*Korea Astronomy and Space Science Institute (한국천문연구원), Korea,* ²*University of Science and Technology (과학기술연합대학원대학교), Korea,* ³*Genesia co., Japan,* ⁴*Seoul National University (서울대학교), Korea,* ⁵*Kyung Hee University (경희대학교), Korea,* ⁶*Satellite Technology & Research Center (인공위성연구소), KAIST, Korea,* ⁷*ISAS/JAXA, Japan*