

우주론

[구 CO-01] Intrinsic alignments of emission line galaxies at $z \sim 1.4$ from the FastSound redshift survey

Motonari Tonegawa¹, Teppei Okumura², Tomonori Totani³, Gavin Dalton⁴, and Kiyoto Yabe⁵

¹*Korea Institute for Advanced Study, South Korea,*

²*Academia Sinica, Taiwan,*

³*The University of Tokyo, Japan,*

⁴*University of Oxford, United Kingdom,*

⁵*Kavli Institute for the Physics and Mathematics of the Universe, Japan*

Intrinsic alignments (IA), the coherent alignment of intrinsic galaxy orientations, can be a source of a systematic error of weak lensing surveys. The redshift evolution of IA also contains information about the physics of galaxy formation and evolution. We present the first measurement of IA at high redshift, $z \sim 1.4$, using the spectroscopic catalog of blue star-forming galaxies of the FastSound redshift survey, with the galaxy shape information from the Canada-Hawaii-France telescope lensing survey. The IA signal is consistent with zero with power-law amplitudes fitted to the projected correlation functions for density-shape and shape-shape correlation components, $\Delta_{\delta+} = -0.0040 \pm 0.0754$ and $\Delta_{++} = -0.0159 \pm 0.0271$, respectively. These results are consistent with those obtained from blue galaxies at lower redshifts (e.g., $\Delta_{\delta+} = 0.0035_{-0.0389}^{+0.0387}$ and $\Delta_{++} = 0.0045_{-0.0168}^{+0.0166}$ at $z=0.51$ from the WiggleZ survey), suggesting no strong redshift evolution of IA. The upper limit of the constrained IA amplitude corresponds to a few percent contamination to the weak-lensing shear power spectrum, resulting in systematic uncertainties on the cosmological parameter estimations by $-\Delta \sigma_8 < 0.026$ and $-\Delta \Omega_{\mathrm{m}} < 0.019$.

[구 CO-02] Barionic Acoustic Oscillations with 3-point Correlation Function of Quasars

Doohyun Choi¹, Graziano Rossi¹, Zachary Slepian², Daniel Eisenstein³

¹*Sejong University (optionally affiliation in Korea in parenthesis),* ²*Lawrence Berkeley National Laboratory,* ³*Harvard-Smithsonian Center for Astrophysics*

While quasars are sparse in number density, they reside at relatively high-redshift as compared to e.g. luminous red galaxies. Hence, they are likely to be less non-linearly evolved than the galaxy population, and thus have a distribution that more closely mirrors the primordial density field. Therefore, they offer an intriguing opportunity to search for Baryonic Acoustic Oscillations (BAO). To this end, the 3-point correlation function (3PCF) is an excellent statistical tool to detect BAO. In this work, we will make the first-ever measurement of the large-scale quasar 3PCF from the SDSS-IV DR14 quasar sample (spanning the largest volume to-date). This work will use the order N^2 -time 3PCF algorithm of Slepian & Eisenstein (2015), with N the number of objects.

[구 CO-03] Cosmological parameter constraints from galaxy-galaxy lensing with the Deep Lens Survey

Mijin Yoon and Myungkook James Jee
Yonsei University

The Deep Lens Survey (DLS), a precursor to the Large Synoptic Survey Telescope (LSST), is a 20 deg² survey carried out with NOAO's Blanco and Mayall telescopes. DLS is unique in its depth reaching down to ~ 27 th mags in BVRz bands. This enables a broad redshift baseline and is optimal for investigating cosmological evolution of the large scale structure.

Galaxy-galaxy lensing is a powerful tool to estimate averaged matter distribution around lens galaxies by measuring shape distortions of background galaxies. The signal from galaxy-galaxy lensing is sensitive not only to galaxy halo properties, but also to cosmological environment at large scales. In this study, we measure galaxy-galaxy lensing and galaxy clustering, which together put strong constraints on the cosmological parameters.

We obtain significant galaxy-galaxy lensing signals out to ~ 20 Mpc while tightly controlling systematics. The B-mode signals are consistent with zero. Our lens-source flip test indicates that minimal systematic errors are present in DLS photometric redshifts. Shear calibration is performed using high-fidelity galaxy image simulations. We demonstrate that the overall shape of the galaxy-galaxy lensing signal is well described by the halo model comprised of central and non-central halo contributions. Finally, we present our preliminary constraints on the matter

density and the normalization parameters.

[구 CO-04] HectoMAP and Horizon Run 4: Over- and Under-dense Large-scale Structures in the Real and Simulated Universe

Ho Seong Hwang¹, M. J. Geller², Changbom Park³, D. G. Fabricant², M. J. Kurtz², K. J. Rines⁴, Juhan Kim⁵, A. Diaferio⁶, H. J. Zahid², P. Berlind², M. Calkins², S. Tokarz², S. Moran²

¹Quantum Universe Center, Korea Institute for Advanced Study, ²Smithsonian Astrophysical Observatory, ³School of Physics, Korea Institute for Advanced Study, ⁴Western Washington University, ⁵Center for Advanced Computation, Korea Institute for Advanced Study, ⁶Università degli Studi di Torino/Istituto Nazionale di Fisica Nucleare

HectoMAP is a dense redshift survey of red galaxies covering a 53 square degree strip of the northern sky, and Horizon Run 4 is one of the densest and largest cosmological simulations based on the standard Lambda cold dark matter model. We use HectoMAP and Horizon Run 4 to compare the physical properties of observed large-scale structures with simulated ones in the redshift range $0.22 < z < 0.58$. We find that the properties of the largest over- and under-dense structures in HectoMAP are well within the distributions for the largest structures drawn from 300 Horizon Run 4 mock surveys. Overall the size, richness and volume distributions of observed large-scale structures when the universe is ~ 9.4 Gyrs old are remarkably consistent with predictions of the standard Lambda cold dark matter model.

[구 CO-05] Lyman alpha radiative transfer at the epoch of cosmic reionization

Hyo Jeong Kim¹, Hyunbae Park², Kyungjin Ahn¹
¹Department of Earth Science Education, Chosun University, ²Korea Astronomy and Space Science Institute

We present a numerical code for the random scattering histories of Lyman alpha photons in the intergalactic medium. The numerical code calculates the radiative transfer under generic three dimensional density, ionization fraction, and peculiar velocity fields based on N-body + radiation transfer simulations of the epoch of reionization. The code is tested with models having analytical solutions, which have idealized geometry and simplified velocity fields. The emergent line

profiles can give constraints to the ionization structure around Lyman alpha sources in the early universe.

[구 CO-06] Lagrangian Perturbation Theory for the Cosmological Structure Formation with 2-component Fluid

Kyungjin Ahn
Chosun University

We present the preliminary result of our Lagrangian perturbation theory for the large-scale structure formation, in the presence of the cold dark matter (CDM) and the baryonic fluid. In the linear order, two mutually independent pseudo-particles can describe the evolution of density fluctuations and the accuracy of the calculation is better than the 4-mode (growing, decaying, streaming, compensated) Eulerian linear perturbation theory. In the 2nd order, the separability of pseudo-particles is not as straightforward as in the linear order, and the related difficulty in developing the 2nd order theory will also be presented.

[구 CO-07] Marked correlation function as modified gravity probe

Yi Zheng
School of Physics, Korea Institute for Advanced Study, 85 Heogiro, Dongdaemun-gu, Seoul 130-722, Korea

For modified gravity models with screening mechanisms, the signal of modified gravity is larger at low density areas. We could add a density weighted mark to galaxy correlation function and increase the signal to noise of modified gravity detection. Based on mock galaxy catalogs from modified gravity simulations, we try different mark functions and parameters to find the best choices for discriminating modified gravity from GR. In this talk I will present our marked correlation function results and discuss its advantages and disadvantages.

[구 CO-08] Evidence for galaxy dynamics tracing background cosmology below the de Sitter scale of acceleration

Maurice H.P.M van Putten
Sejong university, Astronomy & Space science department