

[구 GC-12] Ly α Radiative Transfer and The Wouthuysen-Field effect

Kwang-Il Seon¹ (선광일), Chang-Goo Kim² (김창구)
¹Korea Astronomy and Space Science Institute
 (한국천문연구원), ²Princeton University, USA

A three-dimensional (3D) Ly α radiative transfer code is developed to study the Wouthuysen-Field effect, which couples the 21 cm spin temperature of neutral hydrogen and the Ly α radiation field, and the escape fraction of Ly α from galaxies. The Monte Carlo code is capable of treating arbitrary 3D distributions of Ly α source, neutral hydrogen and dust densities, gas temperature, and velocity field. It is demonstrated that the resonance-line profile at the center approaches to the Boltzmann distribution with the gas temperature. A plane-parallel ISM model, which is appropriate for the neutral ISM of our Galaxy, is used to calculate the Ly α radiation field strength as a function of height above the galactic plane. We also use a two-phase, clumpy medium model which is composed of the cold and warm neutral media (WNM). It is found that the Ly α radiation field is strong enough to thermalize the 21 cm spin temperature in the WNM to the gas kinetic temperature. The escape fraction of Ly α is found to be a few percent, which is consistent with the Ly α observations of our Galaxy and external galaxies.

[구 GC-13] Discovery of a significant population of CN-enhanced red clump stars in the Milky Way bulge: Implications for the formation of early-type galaxies

Young-Wook Lee, Seungsoo Hong, Dongwook Lim
 Yonsei University, Seoul 03722, Korea

We report our discovery of a significant population of CN-enhanced red clump stars in the classical bulge component of the Milky Way. Since CN-rich stars trace a population with enhanced Helium, Nitrogen, and Sodium originated in globular clusters (GCs), this is a direct evidence that proto-GCs were the major building blocks in the formation of the classical bulges and early-type galaxies in the hierarchical merging paradigm.

[구 GC-14] Three-Dimensional Structure of Star-Forming Regions in NGC 6822 Hubble V

Hye-In Lee¹, Heeyoung Oh², Huynh Anh N. Le³,
 Soojong Pak¹, Sungho Lee², Gregory Mace⁴, Daniel

T. Jaffe⁴, Quang Nguyen-Luong², Ken'ichi
 Tatematsu⁵

¹School of Space Research, Kyung Hee University,
²Korea Astronomy & Space Science institute,
³Department of Physics and Astronomy, Seoul
 National University, ⁴Department of Astronomy, the
 University of Texas, ⁵National Astronomical
 Observatory of Japan

NGC 6822 is a dwarf irregular galaxy in the Local Group and it is located in 500 kpc, further than the Large Magellanic Cloud and the Small Magellanic Cloud. Therefore, we can study star-forming processes by local condition in NGC 6822 instead of tidal force of the Galactic gravitational field. Hubble V is the brightest of several H II complexes in this galaxy. We observed Hubble V by using IGRINS attached on the 2.7 m telescope at the McDonald Observatory in Texas, US in May 2016. We performed a spectral mapping of 15" x 7" area on H and K bands, and detected emission lines of bright Br γ λ 2.1661 μ m and weak He I λ 2.0587 μ m. Molecular hydrogen lines of 1-0S(1) λ 2.1218 μ m, 2-1 S(1) λ 2.2477 μ m, and 1-0 S(0) λ 2.2227 μ m was also detected. These emission lines show the structure of an ionized core and excited surface of clouds by far-ultraviolet photons, photodissociation region (PDR). We present three-dimensional maps of emission line distributions through multi slit scanning data and compare these results with the previous study. This presentation shows the physical structure of the star-forming regions and we discuss a PDR model and an evolution of Hubble V complex.

[구 GC-15] Bar Fraction in Early-type and Late-type

Yun Hee Lee¹, Hong Bae Ann², and Myeong-Gu
 Park¹

¹Department of Astronomy and Atmospheric
 sciences, Kyungpook National University, ²Division
 of Science Education, Pusan National University

Bar fractions depend on the properties of host galaxies. However, the observational studies did not provide consistent tendency. We investigated the bar fractions and their dependence on properties of host galaxies using three bar classifications: visual inspection, ellipse fitting method and Fourier analysis from a volume-limited sample of 1,698 disk galaxies brighter than Mr=-15.2 within z = 0.01 from the Sloan Digital Sky Survey (SDSS) Data Release 7 (DR7). We found two causes to make the discrepancy in previous studies. One is caused by