

overview of the expected EM signals from GW sources and the current EM follow-up observations that have been undertaken in Korea and the world.

[구 GW-04] Gravitational Wave Astrophysics with the Superconducting Low-frequency Gravitational-wave Telescope

Sang-Hyeon Ahn¹, Yeong-Bok Bae¹, Gungwon Kang², Chunglee Kim¹, Whansun Kim³, John J. Oh³, Sang Hoon Oh³, Chan Park², Edwin J. Son³, Hyung Mok Lee⁴, Hyungwon Lee⁵, Hyunkyu Lee⁶, Chang-Hwan Lee⁷, Ho Jung Paik⁸

¹Korea Astronomy & Space Science Institute, Center for Theoretical Astronomy, ²Korea Institute of Science and Technology Information, Division of Supercomputing, ³National Institute for Mathematical Sciences, Division of Industrial Mathematics, ⁴Seoul National University, Dept. of Physics and Astronomy, ⁵Inje University, Dept. of Computer Simulation, ⁶Hanyang University, dept. of Physics, ⁷Pusan National University, dept. of Physics, ⁸U. of Maryland, USA, Dept. of Physics

Gravitational wave (GW) is a probe to observe compact objects (WD, NS, and BHs) in the Universe. Compact binary coalescences (CBCs) were expected to be primary sources of LIGO, VIRGO, and KAGRA. Indeed GW150914 from BH-BH binary coalescence at 430 Mpc was discovered by LIGO between 25-350 Hz. The total system mass of GW150914 is $\sim 70 M_{\odot}$, and about $3 M_{\odot}$ of energy is converted to GWs in 0.2s of the observation duration. In lower frequencies below 10 Hz, in addition to CBCs with $1-100 M_{\odot}$, more massive sources of $\sim 1,000-10,000 M_{\odot}$ are observable for seconds up to days in time scale. We introduce GW astrophysics and present highlights of target sources for the proposed super conducting low-frequency gravitational-wave telescope (SLGT).

[구 GW-05] Stochastic Gravitational Wave Background in 0.1-10 Hz

Chan Park¹, Sang-Hyeon Ahn², Yeong-Bok Bae², Gungwon Kang¹, Chunglee Kim², Whansun Kim³, John J. Oh³, Sang Hoon Oh³, Edwin J. Son³, Yong Ho Lee⁴

¹Korea Institute of Science and Technology Information, ²Korea Astronomy & Space Science Institute, ³National Institute for Mathematical Sciences, ⁴Korea Research Institute of Standards and Science

Stochastic gravitational wave background (SGWB)

is expected to be contributed by primordial sources (e.g. inflation signature) and astrophysical sources (e.g., incoherent superposition of a large numbers of compact binary inspirals throughout in the Universe). Theoretically, SGWB is predicted to span in a broad frequency range between less than nHz up to kHz. Many gravitational-wave (GW) detectors such as LIGO or LISA aim to detect or constrain SGWB in different frequency band that is most sensitive for each detector. In this talk, we focus on the perspectives of constraining the energy density of SGWB between 0.1-10 Hz. We introduce the characteristics of SGWB and representative models for primordial and astrophysical sources. Then, we propose a signal extraction scheme to detect SGWB using one or several omni-directional GW detectors such as SLGT(Superconducting Low-frequency Gravitational-wave Telescope). Considering SLGT sensitivity, we discuss how to observe SGWB in 0.1-10 Hz if we have SLGT network. Finally, we highlight interesting SGWB models that can be constrained in 0.1-10 Hz with SLGT.

[구 GW-06] Binary Black Hole Inspirals and GW detection in 0.1-10 Hz

Yeong-Bok Bae¹, Sang-Hyeon Ahn¹, Gungwon Kang², Chunglee Kim¹, Whansun Kim³, John J. Oh³, Sang Hoon Oh³, Chan Park², Edwin J. Son³, Yong Ho Lee⁴

¹Korea Astronomy & Space Science Institute, ²Korea Institute of Science and Technology Information, ³National Institute for Mathematical Sciences, ⁴Korea Research Institute of Standards and Science

The pilot study of SLGT (Superconducting Low-frequency Gravitational-wave Telescope) is being performed by KKN (KASI-KISTI-NIMS) collaboration. In this presentation, we discuss perspectives of detecting GWs in the low-frequency band (0.1-10 Hz), which is a target frequency band of SLGT, but can be hardly observed by advanced LIGO. IMBHs (Intermediate Mass Black Hole Binaries) and IMRIs (Intermediate Mass Ratio Inspirals) with total masses of $O(1000)$ up to $O(10,000)$ solar masses are most probable sources between 0.1-10 Hz. We estimate horizon distances and signal to noise ratios of IMBHs and IMRIs for different SLGT design sensitivities. Based on our calculations, detection rates for IMBHs and IMRIs with SLGT will be discussed.