

Department of Physics and Astronomy, Seoul National University, Republic of Korea

Oscillations in a sunspot are easily detected through the Doppler velocity observation. Although the sunspot oscillations look erratic, the wavelet analysis show that they consist of successive wave packets which have strong power near three or five minutes. Previous studies found that 3-min oscillation at the chromosphere is a visual pattern of upward propagating acoustic waves along the magnetic field lines. Recent multi-height observations help this like vertical study, however, we also focus on horizontal facet to extend three dimensional understand of sunspot waves. So, we investigate a fragmented sunspot expected to have complex wave profiles according to the positions in the sunspot observed by the Fast Imaging Solar Spectrograph. We choose 4 points at different umbral cores as sampling positions to determine coherence of oscillations. The sets of cross-correlation with three and five minutes bandpass filters during a single wave packet reveal interesting results. Na I line show weak correlations with some lags, but Fe I and Ni I have strong correlations with no phase difference over the sunspots. It is more remarkable at Ni I line with 3-min bandpass that all sets of cross-correlation look like the autocorrelation. We can interpret this as sunspot oscillations occur spontaneously over a sunspot at photosphere but not at chromosphere. It implies a larger or deeper origin of 3-min sunspot oscillation.

[7 SS-06] The Standard Processing of a Time Series of Imaging Spectral Data Taken by the Fast Imaging Solar Spectrograph on the Goode Solar Telescope

Jongchul Chae, Juhyeong Kang, Kyuhyoung Cho
Department of Physics and Astronomy, Seoul National University (서울대)

The Fast Imaging Solar Spectrograph (FISS) on the Goode Solar Telescope (GST) at Big Bear Solar Observatory is the imaging Echelle spectrograph developed by the Solar Astronomy Group of Seoul National University and the Solar and Space Weather Group of Korea Astronomy and Space Science Institute. The instrument takes spectral data from a region on the Sun in two spectral bands simultaneously. The imaging is done by the organization of intensity data obtained from the fast raster scan of the slit over the field of view. Since the scan repeats many times, the whole set

of data can be used to construct the movies of monochromatic intensity at arbitrary wavelengths within the spectral bands, and those of line-of-sight velocity inferred from different spectral lines. So far there are two standard observing configurations: one recording the H α line and the Ca II 8542 line simultaneously, and the other recording the Na I D2 line and Fe I 5435 line simultaneously. We have developed the procedures to produce the standard data for each observing configuration. The procedures include the spatial alignment, the correction of spectral shift of instrumental origin, and the wavelength measurement of the line wavelength. The standard data include the movie of continuum intensity, the movies of intensity and velocity inferred from a chromospheric spectral line, the movies of intensity and velocity inferred from a photospheric line. The processed standard data will be freely available online (fiss.snu.ac.kr) to be used for research and public outreach. Moreover, the IDL procedures will be provided on request as well so that each researcher can adapt the programs for their own research.

[7 SS-07] He I D3 and 10830 observations of the flare-productive active region AR 12673 on 2017 September 7

Yeon-Han Kim^{1,4}, Yan Xu³, Sujin Kim^{1,4}, Su-Chan Bong^{1,4}, Eun-Kyung Lim¹, Heesu Yang¹, Vasylyurchyshyn², Kwangsu Ahn², Young-Deuk Park¹, Phillip R. Goode²
¹KASI, ²BBSO/NJIT, ³CSTR/NJIT, ⁴UST

The active region NOAA AR 12673 is the most flare productive active region in the solar cycle 24. On 2017 September 07, it produced an X1.3 flare, three M-class, and several C-class flares. We successfully observed several C-class flares from 16:50 UT to 22:00 UT using the 1.6m Goode Solar Telescope (GST; formerly NST) at Big Bear Solar Observatory (BBSO). The GST provides us with unprecedented high-resolution data of the Sun since 2009. Interestingly, we observed the active region in He I D3 and 10830 lines simultaneously. The data shows several interesting features: (1) D3 emission seems to be much weaker than 10830 emission around 21:29 UT; (2) a small loop seen in 10830 is moving upward and is brightened around 21:16 UT, but it is not clear in D3; (3) there are waves in the penumbra seen in 10830 line center; (4) there is a jet with twisting motion. In this presentation, we will give the results of our analysis and interpretations.