

Although mass accretion from the disk to the central protostar is a key process of low mass star formation, the accretion mechanism is still poorly understood. To investigate “episodic accretion”, which has been suggested as an accretion mechanism in low mass star formation, we have carried out near-infrared spectroscopic observations of three very low-luminosity objects (VeLLOs) and one background source, using InfraRed Camera onboard the AKARI space telescope. The ice absorption features of H<sub>2</sub>O, CO<sub>2</sub>, and CO were detected around the wavelengths of 3.0, 4.26, and 4.67 μm, respectively. In addition, we revealed the XCN ice feature, which is attributed to high energy UV photons produced by the episodic burst accretion. The comparisons of the ice abundances of our targets with those of other YSOs observed previously with AKARI IRC imply that the three VeLLOs had experienced burst accretions although they are now in a very quiescent phase.

#### [석 IM-09] New insights on the origin of multiple stellar populations in globular clusters

Jaeyeon Kim (김재연) and Young-Wook Lee (이영욱)  
*Center for Galaxy Evolution Research & Department of Astronomy, Yonsei University*

In order to investigate the origin of multiple stellar populations in the halo and bulge of the Milky Way, we have constructed chemical evolution models for the low-mass proto-Galactic subsystems such as globular clusters. Unlike previous studies, we assume that supernova blast waves undergo blowout without expelling the pre-enriched gas, while relatively slow winds of massive stars, together with the winds and ejecta from low and intermediate mass asymptotic-giant-branch stars, are all locally retained in these less massive systems. We find that the observed Na-O anti-correlations in metal-poor GCs can be reproduced when multiple episodes of starbursts are allowed to continue in these subsystems. A specific form of star formation history with decreasing time intervals between the stellar generations, however, is required to obtain this result, which is in good agreement with the parameters obtained from our stellar evolution models for the horizontal-branch. The “mass budget problem” is also much alleviated by our models without ad-hoc assumptions on star formation efficiency and initial mass function. We also applied these models to investigate the origin of super helium-rich red clump stars in the metal-rich bulge as recently suggested by Lee et

al. (2015). We find that chemical enrichments by the winds of massive stars can naturally reproduce the required helium enhancement ( $dY/dZ = 6$ ) for the second-generation stars. Disruption of proto-globular clusters in a hierarchical merging paradigm would have provided helium enhanced stars to the bulge field.

#### [구 IM-10] Radial distribution of blue straggler stars in Magellanic Cloud clusters

Jongsuk Hong  
*IKavli Institute for Astronomy and Astrophysics, Peking University, China*

Using the high-resolution observational data obtained by the Hubble Space Telescope, we found that there is the diversity of the radial trends of blue straggler stars (BSSs) in young massive clusters (YMCs) in the Large Magellanic Cloud unlike BSSs in old globular clusters usually showing the segregated radial distributions. To understand the dynamical processes that lead to the none-segregated or even inversely-segregated radial distribution of BSSs, we performed direct N-body simulations for YMCs. Our numerical simulations show that the presence of black hole subsystems inside the cluster centre can significantly affect the dynamical evolution of BSSs and eventually lead to none- or inversely-segregated radial distribution of BSSs.

#### [구 IM-11] An exosolar planetary system N-body simulator II

ChaeLin<sup>1</sup> Hong, Maurice van Putten<sup>1,2</sup>  
<sup>1</sup>*Center for High Energy Astrophysics, UNIST,*  
<sup>2</sup>*Department of Physics and Astronomy, Sejong University*

We present a general N-body exosolar system simulator in anticipation of upcoming searches for exoplanets and even exomoons by next generation telescopes such as James Webb Space Telescope. For habitable zones, traditionally defined by temperature, we here address the essential problem of dynamical stability of planetary orbits. Illustrative examples are presented on P-type orbits in stellar binary systems, that should be fairly common as in Kepler 16b. Specific attention is paid to reduced orbital lifetimes of exoplanets in the habitable zone by the stellar binary, that is proposed by Maurice van Putten (2017). Especially, we focused on a classic work of complex three-body problem that is well known by Dvorak(1986). We change his elliptic restricted three-body problem to extend unrestricted three-body problem to look into dynamical motions