

Fine mapping of rice bacterial leaf blight resistance loci to major Korean races of Xoo (*Xanthomonas oryzae*)

Myung-Chul Lee*, Yu-Mi Choi, Sukyeung Lee,
Hyemyeong Yoon, Sejong Oh

National Agrobiodiversity Center, National Institute of Agricultural Sciences, RDA,
Jeonju 54874, Republic of Korea

Bacterial leaf blight (BLB), caused by *X. oryzae* pv. *oryzae* (Xoo), is one of the most destructive diseases of rice due to its high epidemic potential. Understanding BLB resistance at a genetic level is important to further improve the rice breeding that provides one of the best approaches to control BLB disease. In the present investigation, a collection of 192 accessions was used in the genome-wide association study (GWAS) for BLB resistance loci against four Korean races of Xoo that were represented by the prevailing BLB isolates under Xoo differential system.

A total of 192 accessions of rice germplasm were selected on the basis of the bioassay using four isolated races of Xoo such as K1, K2, K3 and K3a. The selected accessions were used to prepare 384-plex genotyping by sequencing (GBS) libraries and Illumina HiSeq 2000 paired-end read was used for GBS sequencing. GWAS was conducted using TASSEL 5.0. The TASSEL program uses a mixed linear model (MLM). The results of the bioassay using a selected set of 192 accessions showed that a large number of accessions (93.75%) were resistant to K1 race, while the least number of accessions (34.37%) resisted K3a race. For races K2 and K3, the resistant germplasm proportion remained between 66.67 to 70.83%. The genotypic data produced SNP matrix for a total of 293,379 SNPs. After imputation the missing data was removed, which exhibited 34,724 SNPs for association analysis. GWAS results showed strong signals of association at a threshold of $[-\log_{10}(P\text{-value})]$ more than 5 (K1 and K2) and more than 4 (K3 and K3a) for nine of the 39 SNPs, which are plausible candidate loci of resistance genes. These SNP loci were positioned on rice chromosome 2, 9, and 11 for K1 and K2 races, whereas on chromosome 4, 6, 11, and 12 for K3 and K3a races. The significant loci detected have also been illustrated, NBS-LRR type disease resistance protein, SNARE domain containing protein, Histone deacetylase 19, NADP-dependent oxidoreductase, and other expressed and unknown proteins. Our results provide a better understanding of the distribution of genetic variation of BLB resistance to Korean pathogen races and breeding of resistant rice

[Acknowledgements]

This work was supported by a grant No. PJ0135392018) from the National Institute of Agricultural Sciences, Rural Development Administration, Republic of Korea.

Keywords: Rice. Bacterial leaf blight, GWAS, Population structure