

although the moment of the outburst was not caught. The embedded protostar IRAS 16316-1540 observed with the Immersion Grating Infrared Spectrograph (IGRINS, $R = \Delta\lambda/\lambda \sim 45000$) shows the broad absorption features in atomic and CO transitions, as seen in FU Orionis objects (FUors), indicative of an outburst event. We examine whether the spectra of IRAS 16316-1540 arise from the rotating inner hot gaseous disk. Using the IGRINS spectral library, we show that the line profiles of IRAS 16316-1540 are more consistent with an M1.5 V template spectrum convolved with a disk rotation profile than the protostellar photosphere absorption features with a high stellar rotation velocity. We also note that the absorption features deviated from the expected line profile of the accretion disk model can be explained by a turbulence motion generated in the disk atmosphere. From previous observations that show the complex environment and the misaligned outflow axes in IRAS 16316-1540, we suggest that an impact of infalling clumpy envelope material against the disk induces the disk precession, causing the accretion burst from the inner disk to the protostar.

[구 IM-06] The JCMT Transient Survey: Examination of Periodic Variability in nearby Star-forming Regions

Yong-Hee Lee¹, Jeong-Eun Lee¹, Doug Johnstone^{2,3}, Gregory J. Herczeg³, Steve Mairs^{2,3,5}

¹*School of Space Research, Kyung Hee University, Republic of Korea* ²*NRC Herzberg Astronomy and Astrophysics Canada* ³*Department of Physics and Astronomy, University of Victoria, Canada* ⁴*Kavli Institute for Astronomy and Astrophysics, Peking University, People's Republic of China* ⁵*East Asian Observatory, USA*

We perform the Lomb-Scargle Periodogram analysis to protostars identified by the JCMT Transient Survey, which monitors 8 nearby star forming regions. The observations have been done monthly for over 3 years using SCUBA-2 (the Submillimetre Common User Bolometer Array 2) in two wavelengths, 450 and 850 μm . Under the threshold of 1% False Alarm Probability, we found 16 variable sources including EC53, which is the first variable protostar detected by the JCMT Transient Survey. Most of the variable sources are cataloged as protostars (classified via the Spitzer data, Megeath et al. 2012; Dunham et al. 2015), but SerpS-MM19, which has a clear 1-year period, is a candidate of a first hydrostatic core (Maury et al. 2011; Young et al. 2018).

[구 IM-07] Removing Large-scale Variations

in Regularly and Irregularly Spaced Data

Jungyeon Cho (조정연)
Chungnam National University

In many astrophysical systems, smooth large-scale variations coexist with small-scale fluctuations. For example, a large-scale velocity or density gradient can exist in molecular clouds that have small-scale fluctuations by turbulence. In redshifted 21cm observations, we also have two types of signals - the Galactic foreground emissions that change smoothly and the redshifted 21cm signals that fluctuate fast in frequency space. In many cases, the large-scale variations make it difficult to extract information on small-scale fluctuations. We propose a simple technique to remove smooth large-scale variations. Our technique relies on multi-point structure functions and can obtain the magnitudes of small-scale fluctuations. It can also be used to design filters that can remove large-scale variations and retrieve small-scale data. We discuss how to apply our technique to irregularly spaced data, such as rotation measure observations toward extragalactic radio point sources.

[구 IM-08] The distribution of magnetic field strength in Orion A region

Jihye Hwang^{1,2}, Jongsoo Kim^{1,2}
¹*Korea Astronomy and Space Science Institute,*
²*University of Science and Technology, Korea*

Magnetic fields play an important role in supporting molecular clouds against gravitational collapse. The measured magnetic field strengths in molecular clouds enable us to see the effect of magnetic fields in star-forming regions. People have used the Chandrasekhar and Fermi (CF) method to estimate magnetic field strength from observational quantities of molecular cloud density, turbulent velocity and polarization angle dispersion. However, previous studies obtained just one magnetic field strength over the quite large region of a molecular cloud by using the CF method. We here suggest a way to estimate magnetic field strength distribution in Orion A region. We used 450 and 850-micron polarization data of James Clerk Maxwell Telescope (JCMT). Magnetic field strengths were estimated in two wavelengths with 4 pixel resolutions of 16, 20, 24 and 28". Through statistical analysis, we proved the difference of magnetic field strengths between two wavelengths were caused by the difference of their beam sizes. Additionally, we calculated the radii of curvature of polarization segments to