

Estimation of Na Abundance from Low-resolution Stellar Spectra.

Hyejin Park, Young Sun Lee
Chungnam National University

It is inferred that many stars in the Galactic halo or bulge were once members of globular clusters (GCs), which are now dissolved. To distinguish the GC-originated stars, which can provide valuable information on the origin of the bulge and halo, from the in situ field stars, the Na abundance plays an important role. However, the interstellar Na in certain directions can unnecessarily enhance the estimate of the Na abundance from stellar spectra due to blended Na D lines unless the spectral resolution is very high, which allows to resolve the lines from the interstellar Na. In this study, we present a means of correcting the Na abundance affected by the interstellar Na in the low-resolution of the Sloan Digital Sky Survey stellar spectra.

[포 IM-11] ALMA Observations of a Massive-star-forming Infrared Dark Cloud Core MSXDC G053.11+00.05 MM1

Hyun-Jeong Kim¹, Bon-Chul Koo¹, Kee-Tae Kim², Chang-Hee Kim²
¹Seoul National University, ²Korea Astronomy and Space Science Institute

We present the ALMA observations of the infrared dark cloud (IRDC) core MSXDC G053.11+00.05 MM1 at the distance of 1.7 kpc. While the core was first identified at 1.2 mm with a mass of 124 Msun, recent near- and mid-infrared observations have revealed a parsec-scale molecular hydrogen (H₂ 1-0 S(1) at 2.12 micron) outflow and two early class young stellar objects (YSOs) at the center of the core, one of which is likely massive (M > 8 Msun). From the ALMA Band 7 observations with a resolution of 0.5", we have found a dust filament of < 0.1 pc in which five dense cores are embedded in the 870 micron continuum. The brightest core is consistent with one of the two previously-detected YSOs, but the other four are newly discovered implying their very deeply embedded status. We have also detected several molecular line emission including H₁₃CO⁺ and C₁₇O as well as ¹³CO outflow with complicated morphology. At the brightest core, the methanol line (CH₃OH) shows velocity gradients, which may support the existence of a circumstellar disk around a high-mass protostar. Based on the derived properties of the dense cores, we discuss their association with the two YSOs and H₂ outflow detected in infrared and high-mass star-formation

process occurring in IRDC cores.

[포 IM-12] TRAO Survey of Nearby Filamentary Molecular Clouds, the Universal Nursery of Stars (TRAO FUNS). III. Dynamics of filaments in different star forming environments

Eun Jung Chung¹, Shinyoung Kim^{1,2}, Hyunju Yoo¹, and Chang Won Lee^{1,2}
¹Korea Astronomy and Space Science Institute, ²University of Science and Technology

Recent high resolution IR observations reveal that molecular clouds are filamentary and such a structure is ubiquitous over various star-forming environments, and it is clear that filaments play a crucial role in the formation of cores and stars. However, the formation process of dense cores in the filaments are still unknown. To investigate this issue in detail, we have carried out TRAO FUNS (TRAO survey of nearby Filamentary molecular clouds, the Universal Nursery of Stars) toward various star forming filamentary molecular clouds.

In this presentation, we will report the first look results of filaments and dense cores in MCLD 123.5+24.9 and IC 5146, which are known as a quiescent, non-star-forming region and an active, high-mass star forming region, respectively. By comparing the kinematic properties of filaments and dense cores in different star forming environments, we verified the formation scenario of filaments and dense core, i.e., gravoturbulent fragmentation via supersonic motions.

[포 IM-13] JCMT-CHIMPS2 Survey

Kee-Tae Kim¹, Toby Moore², Tetsuhiro Minamidani³, Oscar Morata⁴, Erik Rosolowski⁴, Yang Su⁵, David Eden², and the CHIMPS2 Team
¹KASI, ²Liverpool John Moores University, UK, ³NAOJ, Japan, ⁴ASIAA, Taiwan, ⁵University of Alberta, Canada, ⁶Purple Mountain Observatory, China

The CHIMPS2 survey is to extend the JCMT HARP ¹³CO/¹⁸O J=3-2 Inner Milky-Way Plane Survey (CHIMPS) and the ¹²CO J=3-2 survey (COHRS) into the inner Galactic Plane, the Central Molecular Zone (CMZ), and a section of the Outer Plane. When combined with the complementary ¹²CO/¹³CO/¹⁸O J=1-0 survey at the Nobeyama 45m (FUGIN) at matching 15" resolution and sensitivity, and other current CO surveys, the results will provide a complete set of transition data with which to calculate accurate column densities, gas temperatures and turbulent Mach numbers. These will be used to: analyze molecular cloud properties