

(한국천문연구원), ²University of Science & Technology, Korea (과학기술연합대학원대학교),
³Hanyang University (한양대학교)

Studying the amount and kinematics of circum- and intergalactic medium (CGM and IGM) is key to understanding the role of feedback and environment (cold streams and galactic winds) in the evolution of galaxies. In particular, Ly α emission line has been utilized to investigate the density structure and kinematics of the (most abundant) H I gas in the CGM and IGM around galaxies. Therefore, modeling Ly α radiative transfer through multiphase interstellar medium (ISM), CGM and IGM is crucial in understanding the galaxy evolution. As discussed in Kakiichi & Dijkstra (2018), most Ly α RT effects would occur on interstellar scales. This is because the main source of Ly α photons would be H II regions, which are in most cases, if not all, surrounded by “cold” photo-dissociation regions. However, most Ly α RT studies have been performed in the CGM and IGM environments with $T \sim 10,000\text{K}$. In this talk, we present how the Ly α RT effect in the cold ISM with $T \sim 100\text{K}$ regulates the Ly α spectral properties.

[포 GC-05] Hubble Space Telescope Survey of Host Galaxies of Hard X-ray-Selected AGNs

Hyunmo Hwang¹, Minjin Kim¹, Aaron J. Barth², Luis C. Ho^{3,4}

¹Department of Astronomy and Atmospheric Sciences, Kyungbook National University

²Department of Physics and Astronomy, University of California at Irvine

³Kavli Institute for Astronomy and Astrophysics

⁴Department of Astronomy, Peking University

We present an ongoing imaging survey of the host galaxies of hard X-ray-selected active galactic nuclei (AGNs) observed with the Hubble Space Telescope (HST). The snapshot images are taken with the Advanced Camera for Surveys through an HST gap-filler program. The sample, selected from the 70-month Swift-BAT X-ray source catalog, represents an unbiased and uniform AGN population, which will enable us to test the AGN unification model and explore the physical connection between host galaxies and central supermassive black holes. We also plan to investigate the AGN triggering mechanism by examining merger signatures and searching for dual nuclei. We present the pipeline for imaging analysis and the current status of the survey.

[포 GC-06] Spectroscopic observation of the massive high- z ($z=1.48$) galaxy cluster SPT-CL J2040-4451 using Gemini Multi-Object Spectrographs

Jinhyub Kim^{1,2}, Myungkook J. Jee^{1,2}, Seojin F. Kim¹, and Jongwan Ko³.

¹Department of Astronomy, Yonsei University,

²Department of Physics, University of California, Davis,

³Korea Astronomy and Space Science Institute

Mass measurement of high-redshift galaxy clusters with high accuracy is important in constraining cosmological parameters. Extremely massive clusters at high redshift may impose a serious tension with the current Λ CDM paradigm. SPT-CL J2040-4451 at $z=1.48$ is considered one such case given its redshift and mass estimate inferred from the SZ data. The system has also been confirmed to be indeed massive from a recent weak-lensing (WL) analysis. Comparison of the WL mass with the spectroscopic result may provide invaluable information on the dynamical stage of the system. However, the existing spectroscopic coverage of the cluster is extremely poor; only 6 blue star-forming galaxies have been found within the virial radius, which results in highly inflated and biased velocity dispersion. In this work, we present a spectroscopic analysis of the member candidates using Gemini Multi-Object Spectrographs (GMOS) observation in Gemini South. The observation was designed to find early-type member galaxies within the virial radius and to obtain reliable velocity dispersion. We explain our selection scheme and preliminary results of the spectra. We also compare the dynamical mass estimate inferred from the velocity dispersion with the WL mass.

[포 GC-07] Star-gas misalignment in Horizon-AGN simulation

Donghyeon J. Khim¹, Sukyoung K. Yi¹

¹Department of Astronomy and Yonsei University Observatory, Yonsei University, Seoul 03722, Republic of Korea

Recent Integral Field Spectroscopy (IFS) studies revealed that not only late type galaxies (LTGs) but also early type galaxies (ETGs) have various kinds of kinematic rotation. (e.g. not clearly detectable rotation, disk-like rotation, kinematically distinct core (Cappellari 06)) Among the various studies about galactic kinematics, one of the most notable anomalies is the star-gas misalignment. The gas