

[포 CD-03] Graph Database Solution for Higher Order Spatial Statistics in the Era of Big Data

Cristiano G. Sabiu¹, Juhan Kim²

¹*Yonsei University, Department of Astronomy*

²*Korea Institute for Advanced Study, Center for Advanced Computation*

We present an algorithm for the fast computation of the general N-point spatial correlation functions of any discrete point set embedded within an Euclidean space of \mathbb{R}^n . Utilizing the concepts of kd-trees and graph databases, we describe how to count all possible N-tuples in binned configurations within a given length scale, e.g. all pairs of points or all triplets of points with side lengths $< r_{max}$. Through benchmarking we show the computational advantage of our new graph-based algorithm over more traditional methods. We show that all 3-point configurations up to and beyond the Baryon Acoustic Oscillation scale (~ 200 Mpc in physical units) can be performed on current Sloan Digital Sky Survey (SDSS) data in reasonable time. Finally we present the first measurements of the 4-point correlation function of ~ 0.5 million SDSS galaxies over the redshift range $0.43 < z < 0.7$.

We present the publicly available code GRAMSCI (GRAph Made Statistics for Cosmological Information; bitbucket.org/csabiu/gramsci), under a GNU General Public License.

[포 CD-04] The Joint analysis of galaxy clustering and weak lensing from the Deep Lens Survey to constrain cosmology and baryonic feedback

Mijin Yoon¹ (윤미진), M. James Jee^{1,2} (지명국), and J. Tony Tyson²

¹*Yonsei University (연세대학교)* ²*University of California, Davis*

Based on three types of 2-point statistics (galaxy clustering, galaxy-galaxy lensing, and cosmic shear power spectra) from the Deep Lens Survey (DLS), we constrain cosmology and baryonic feedback. The DLS is a deep survey, so-called a precursor to LSST, reaching down to ~ 27 th magnitude in BVRz' over 20 deg^2 . To measure the three power spectra, we choose two lens galaxy populations centered at $z \sim 0.27$ and 0.54 and two source galaxy populations centered at $z \sim 0.64$ and 1.1 , with more than 1 million galaxies.

We perform a number of consistency tests to confirm the reliability of the measurements. We calibrated photo-z estimation of the lens galaxies

and validated the result with galaxy cross-correlation measurement. The B-mode signals, indicative of potential systematics, are found to be consistent with zero. The two cosmological results independently obtained from the cosmic shear and the galaxy clustering + galaxy-galaxy lensing measurements agree well with each other. Also, we verify that cosmological results between bright and faint sources are consistent.

While there exist some weak lensing surveys showing a tension with Planck, the DLS constraint on S8 agrees nicely with the Planck result. Using the HMcode approach derived from the OWLS simulation, we constrain the strength of baryonic feedback. The DLS results hint at the possibility that the actual AGN feedback may be stronger than the one implemented in the current state-of-the-art simulations.

[포 CD-05] Detection of Intrinsic Spin Alignments in Isolated Spiral Pairs

Hanwool Koo^{1,2}, Jounghun Lee³

¹*Korea Astronomy and Space Science Institute,*

²*University of Science and Technology (Korea),*

³*Seoul National University*

Observational evidence for intrinsic galaxy alignments in isolated spiral pairs is presented. From the catalog of the galaxy groups identified by Tempel et al. in the flux-limited galaxy sample of the Sloan Digital Sky Survey Data Release 10, we select those groups consisting only of two spiral galaxies as isolated spiral pairs and investigate if and how strongly the spin axes of their two spiral members are aligned with each other. We detect a clear signal of intrinsic spin alignment in isolated spiral pairs, which leads to the rejection of the null hypothesis at the 99.9999% confidence level via the Rayleigh test. It is also found that those isolated pairs comprising two early-type spiral galaxies exhibit the strongest signal of intrinsic spin alignment and that the strength of the alignment signal depends on the angular separation distance as well as on the luminosity ratio of the member galaxies. Using the dark matter halos consisting of only two subhalos resolved in the EAGLE hydrodynamic simulations, we repeat the same analysis but fail to find any alignment tendency between the spin angular momentum vectors of the stellar components of the subhalos, which is in tension with the observational result. Several possible sources of this apparent inconsistency between the observational and the numerical results are discussed.

[포 CD-06] Cosmology in UOS: Case with