

## Lectotypification of *Anaphalis morii* Nakai (Asteraceae) with Brief Discussions of its Taxonomic Entity

**Dong Hyuk Lee<sup>1</sup>, Jun Gi Byeon<sup>1</sup>, Tae Im Heo<sup>1</sup>, Byeong Joo Park<sup>1</sup>,  
Ji Dong Kim<sup>1</sup>, Jun Woo Lee<sup>1</sup>, Byeong Kwon Lee<sup>1</sup> and Byeong-Hee Choi<sup>2\*</sup>**

<sup>1</sup>Baekdudaegan National Arboretum

<sup>2</sup>Inha University

Systematic studies for genus *Anaphalis* in Korea are relatively scarce. As a fundamental step to further taxonomic studies of this genus, we here propose lectotype for *A. morii*, endemic taxa in Korea with a brief re-evaluation of its taxonomic entity. During the examination of herbarium specimen for *A. morii*, we found nine sheets of five collections in three herbaria. Among the original materials, we selected a specimen in TI which was first cited by the author and include an additional annotation, matching with his own description. Also, we were able to determine the taxonomic relationships between *A. morii* and its relatives, *A. yakusimensis*. Based on our observation, we identified that several morphological characters are clearly differ from *A. yakusimensis* occurring only Isl. Yakushima in Japan.

[This research was supported by a grant (18172MFDS201) from Ministry of Food and Drug Safety in 2020.]

\*(Corresponding author) E-mail: bhchoi@inha.ac.kr, Tel: +82-32-860-7695