

Fungal Flora in Bamboo Forest of Korea (I)

Ji Yul Lee and Duck Hyun Cho*

Department of Food Science, Seoul Woman's College, Seoul, Korea

*Kyungduck Commercial High School, Andong, Kyungsangbuk-Do, Korea

韓國 竹林產 高等 菌類

李 址 烈 · 趙 德 炫

서울여자대학 식품과학과
 경상북도 안동시 경덕상업고등학교

Abstract: 40 specimens of higher fungi were collected at the bamboo forest of Damyang, Jyunranam-Do, and Kyungju, Kyungsangman-Do during May 6, August 17, 29 and October 23, 1976. These fungi were identified. They were composed of 3 orders, 10 families, 19 genera and 23 species. Among them, *Clitocybe cataneifloccosa* S. Ito et Imal, *Collybia matris* S. Ito et Imal, *Marasmiellus lactea* (Pers. ex Fr.), *Lepiota hetieri* Boud. and *Paxillus atrotomentosus* (Fr.) Fr. var. *bambusinus* Baker et Dale were found to be new in Korea.

Yoshiko shidei reported fungi which were found at the bamboo forest of Japan(1974 and 1976). Attempts to investigate Korean bamboo forest fungi have been made by the authors. 20 specimens were collected at Damyang on August 17. And 20 specimens were collected at Kyungju on May 6, August 29 and October 3.

Bamboos of Damyang were *phyllostachys edulis* Riv. and those found in Kyungju were *phyllostachys reticulata* Koch. Fungi of Damyang were of various kinds and the dominant species of Kyungju was *Dictyophora indusiata* (Pers.) Fishch.

Comparative study of the fungi which were separately found at Damyang and Kyungju will appear in near future. Their located places will be referred to D as to Damyang and K as to Kyungju.

Basidiomycetes 담자균강

Homobasidiae	동담자균강
Hymenomycetes	균실류
Agaricales	송이목
Tricholomataceae	송이과

Clitocybe castaneifloccosa S. Ito et Imai(D)
 꼬마깔대기버섯(신칭)
 S. Ito & Imai, Trans. Sapporo Nat. Hist. Soc. 16: 12. 1939.

Pileus 0.5-1cm broad, funnel-shaped, plane or depressed in center, dark, yellowish brown or grayish yellow brown. Lamellae 1mm wide, sparse, brown. Stipe 0.5-1cm long 0.6-1mm thick brown, hollow, decurrent, thick and roundish downward. Spores subglobose, 7.2-8.6×5.7-7.2 μ , white, enclosed one or some oil drops, nonamyloid, basidia four spores, 31.5-38.6×4.3-7.2 μ .
 Hab. Caespitose on cut bamboo.

C. clavipes (Fr.) Quél. (D) 배불뚝이깔대기버섯
Tricholomopsis platyphylla (Fr.) Sing. (D) 넓
 은주름버섯

Collybia confluens (Fr.) Kummer (D) 밀버섯

C. matris S. Ito et Imai 황갈색밀버섯 (신칭)

S. Ito & Imai, Trans. Sapporo Nat. Hist. Soc.
 16 : 16, 1939.

Pileus 1.6-2.7cm broad, convex, light brown,
 brown at center, dark at margin when old adult.
 Lamellae dense, brown or dark brown. Stipe
 2.5-4.2cm long, 1-3mm thick, equal, yellowish
 gray, adnate. Spores subcylindrical or elliptical.
 7.2-8.6×4.3-5.0μ, enclosed two or three oil
 drops, seldom germ pore present, light brown,
 amyloid. Cystidia 48.6-52.9×17.2-22.9μ, basidia
 four spores, 7.2-20.2×4.3-5.7μ.

Hab. Solitary or clustered on rotten bamboo.

Armillariella mellea (Fr.) Karst. (K) 썩나무버

섯 *Oudemansiella radicata* (Fr.) Sing. (K) 긴
 뿌리버섯

Marasmiellus lactea (Pers. ex Fr.) (D) 꼬마
 흰갓주름 버섯 (신칭)

Agaricus anuns Bull. Herb. Fr. pl. 563, f. N-o.
 1971.

Pileus 2mm broad, white, pale orange when
 dry, with spherical or conic furrow. Lamellae
 sparse. Stipe 1cm long, 1mm thick, adnate, white
 hair of star-shaped distributed upward. Spores
 seed-shaped, white, 6.5-8.6×2.5-4μ, nonamyl-
 oid, formed projection on the surface, basidia
 four spores, 17.2-21.5×5.7-7.2μ Cystidia
 clavate or flask shaped, 35.3-77.6×12.9-13.3μ,
 hyphae from gill trama 4.3-5.7×7.5μ, clamp
 present seldom.

Hab. Caespitose on cut bamboo.

Marasmius siccus (Schw.) Fr. (K) 분홍갓버섯

Agaricaceae 들버섯과

Lepiota hetieri Boud. (D) 대나무우산버섯 (신칭)

Boudier, Bull. Soc. Myc. Fr. 18 : 137. pl. 6
 f.1. 1902

Pileus 1-5cm broad, spherical at first then
 convex, plane when old adult, margin upward,
 farinaceous compactly. pale yellow or pale yello-
 wish brown, flesh tough, white, thick. Lamellae

0.5-2cm wide, dense, pale yellow or pale yello-
 wish brown. Stipe 1.5-5.5cm long, 3-10mm
 thick, scattered white scales and upward of
 annulus. Annulus upward, free, stuffed, scattered
 white powders on the surface. Spores subcylin-
 drical or elliptical, 4.3-5.7×2.6-2.9μ, white,
 nonamyloid, basidia four spores, subcylindrical,
 17.2-18.6×4.3μ, Cystidia clavate, 20.6×13.3μ,
 hyphae from gill trama 35.8-57.2×7.2 9.2μ,
 clamp absent.

Hab. Solitary or clustered on fallen leaves of
 bamboo or on the soils

Coprinaceae

Coprinus friesii Quél. (D,K) 꼬마떡풀버섯

Naematoloma fasciculare (Fr.) Karst. (K) 느
 란타발 (독)

Cortinariaceae 끈적버섯과

Inocyber imosa (Fr.) Quél. (K,D) 팜버섯 (독)

Paxillaceae 우단버섯과

Paxillus atrotomentosus (Fr.) Fr. var. *bambu-*
sinus Baker et Dale (K) 대나무우단버섯 (신칭)

Pileus 3-9.5cm broad, plane or depressed at
 center, glabrous, fan-shaped, mixed pale yellow
 with light brown, dark brown or black when
 old adult, wrinkled, flesh grayish, tough.
 filamentous, margin incurved. Lamellae 4mm
 wide, curly-shaped, sparse, dull orange, then
 changed bluish gray or pale yellowish brown,
 farinaceous. yellow when young black when old
 adult. Stipe 4-6cm long, 0.8-2cm thick, some
 attached laterally, non-cylindrical, brownish
 velvet hair covered irregularly, stuffed, tough,
 flesh white, decurrent. Spores elliptical, 4.3-5.8
 ×3.6-4.3μ, light yellowish, pseudo-amyloid,
 basidia four spores, 27.2-32.9×5.7-7.2μ, hyp-
 hae from stem trama 57.2-72.5×5.7-10μ. Clamp
 present very seldom.

Hab. Solitary or clustered on rotten bamboo.

Boletaceae 그물버섯과

Boletus rubellus Krombh. 붉은두메그물버섯

Russlaccase 무당버섯과

Russla cutifracta Cocke (D) 청버섯

R. fragilis (Pers.) Fr. (D) 애기무당버섯

Polyporaceae 구멍장이버섯과

Elfvigia applanata (Pers.) Karst. (D) 진나
비결상
Ganoderma lucidum (Fr.) Karst. 불노초
Gasteromycetes 북균류
Sclerodermataceae 어리알버섯목
Calostomataceae 연지버섯과
Astracus hygrometricus(Pers.) Morgan (K)
연지버섯
Phallales 말뚝버섯목
Phallaceae 말뚝버섯과
Dictyophora indusiata (Pers.) Fisch. (K) 망
태버섯
Mutinus caninus (Pers.) Fr. (K) 뱀버섯
Pseudocolus schellenbergiae (Sumst.) Johnson
(D) 새발버섯

적 요

1976년 5월 6일, 8월 17, 29일, 10월 3일에 담양
과 경주의 죽림에서 40여개의 고등균류물 채집하여
동정한 결과 3목10과19속23종을 분류하였으며 그 중
에서 한국미 기록종은 다음과 같다.

Clitocybe castaneifloccosa S. Ito et Imai,
Colybia matris S. Ito et Imai, *Marasmiellus*
lactea (Pers. ex Fr.), *Lepiota hetieri* Boud.
Paxillus atrotomentosus (Fr.) Fr. var. *bambusi-*
nus Baker et Dale.

References

Atkinson, Geo. F. (1930): Mushrooms, Hafner, Co.
Hongo, T. (1973): Enumeration of Hygrophora-
ceae, Boletaceae and Strobilomycetaceae. Bu-
lletin of the National Science Museum 16(3),
Tokyo, Japan.
Hong, S.W. (1974): Scientific research of Mt.
Naejang areas, Korea Nature Preservation
Society.
Imazeki, R. and T. Hongo (1957) Colored illu-
strations of fungi of Japan, Hoikusha Publishing
Co. Ltd.
_____. (1965): Colored illustrations of
fungi of Japan vol. II. Hoikusha Publishing
Co. Ltd.

_____, _____ and Tubaki, K. (1970): Common
fungi of Japan in color, Hoikusha Publishing
Co. Ltd.
Ito, S. (1955): Myc. Fl. Japan 2(4), Yokendo.
_____. (1959): Myc. Fl. Japan 2(5), Yokendo.
Kauffman, C.H. (1918): The Agaricus of Mich-
igan vol. I: 1-442, vol. II: 443-924 Dover
Publication Inc.
Kim et al., (1975): Taxonomic Study on Korean
Basidiomycetes. Kor. J. Mycol. 3(2): 31-33
Lee, E.R. and H.S. Jeong (1972): Floral Studies
on the Basidiomycetes in Korea. R-72-82.
Ministry of Science and Technology.
Lee, J.Y. (1957): The list of the fungi of Korea,
Seoul High School.
_____. (1973): The list of the fungi of Korea,
Kor. J. Mycol. I(1): 35-43.
_____. (1973): The higher fungiof Kwangneung
Forest Institute (I).
_____. and D.M. Cho (1975): Notes on Korean
Higher Fungi, Kor. J. Mycol. 3(2): 13-18
Lee, J.Y. et al., (1959): Colored illustrations
of fungi of Korea, Baemungak, Seoul.
Lee, T.S. and Y.W. Lee, (1957): A List of the
Korean Fungi part I, Central Forest Experi-
ment Station, Seoul.
_____. (1958): A List of the Korean Higher
Fungi, part II, Central Forest Experiments
Station, Seoul
Lee, Y.W. (1959): A List of the Korean Higher
Fungi Part III, Central Forest Experiment
Station, Seoul.
Lim, J.H. and B.K. Kim (1972): Taxonomic
investriation on Korean higher fungi (I). Ko.
J. Pharmacog. 3(I): 11-20.
Shidei, Y. (1974): Report of fungal florain ba-
mboo forest. Trans. Mycol. Soc. Japia. 15(3):
251-257.
_____. (1976): Fungal flora and environmental
factors of bamboo forest in Kyoto. Trans.
Mycol. Soc. Japn. 17: 91-94.
Singer, R. (1962): The Agaricales in modern
taxonomy (2nd Ed.) 915pp. 73 pls.

The Explanation of plate I

- A. *Clitocybe castaneifloccossa* S. Ito et Imai
a. carpophores×2 b. spores×1000
c. basidia×1000
- B. *Collybia matris* S. Ito et Imei
a. carpophores×2 b. spores×1000
c. basidia×1000 d. cystidia×1000
- C. *Marasmiellus lactea* (Pers. ex Fr.)
a. carpophores×10 b. spores×1000
c. cystidia×1000
d. hyphae from gill trama×1000
e. basidia×1000

Plate II

- A. *Lepiota hetieri* Boud.
a. carpophores×1 b. spores×1000
c. cystidia×1000 d. basidia×1000
e. hyphae from gill trama×1000
- B. *Paxillus atrotomentosus* (Fr.) Fr. var. *bamb-
usinus* Baker et Dale
a. carpophores×2/3 spores×1000
c. basidia×1000
d. hyphae from stem trama×1000

Plata I






Plate II

