

The Use of Radioimmunoassay to Monitor Reproductive Status of Cheju Native Cattle and the Effect of Supplementary Feeding on Reproduction.

1. Body Weight Changes, Breeding Performances and Progesterone Levels from Weaning until First Calving

Chung, D. C. J. K. Kim, D. C. Kim*, Y. K. Beak* and S. H. Moon*

Radioisotope Research Institute Cheju National University.

Summary

Studies were conducted to investigate the relationship between hormone levels and nutritional levels for improving performance of Cheju native cattle.

In June 1984 a trial was initiated using 8 Cheju native calves after weaning, fed at two supplementary feeding levels (NRC 100% and 70%). The body weight, breeding performance, change in progesterone level during pregnancy and estrus cycle were evaluated.

Mean body weight at 6 months of age was 155kg when fed 100% NRC ration but it was only 137kg when heifers received the 70% NRC ration. At 10, 15 and 20 months of age the body weight was 66, 160 and 115kg, respectively, showing that heifers fed the standard ration gained weight rapidly ($P < 0.01$).

Average size of the left ovary in the standard group was 2.1 x 1.6cm and right ovary was 2.6 x 1.8cm. However in the restricted feeding group the ovaries were found to be smaller. Diameter of graffian follicles showed a similar tendency to ovarian size in the two groups.

The first oestrus in the standard feeding group appeared at 14.6 months when body weight was 265kg. Age at first calving was on average 28.9 months at a body weight of 436kg. On the other hand when heifers were fed the restricted ration the first oestrus appeared at 23.0 months at a body weight of 250kg. Average age at first calving was 38.9 months which was 10 months later than the average in the standard feeding group ($P < 0.01$).

* Cheju Experiment Station.

** This work was supported by International Atomic Energy Agency (Contract No. 3783/RB).

In standard feeding group the progesterone level was 2.0ng/ml at two weeks after pregnancy and gradually increased up to 14 weeks and peaked at 18 weeks. This peak (6.4-6.5ng/ml) was maintained up to 24 weeks when progesterone level decreased until it reached 2.1ng/ml at the end of pregnancy. In the restricted group progesterone level up to 16 weeks followed a similar pattern to the standard group but there was a tendency in the restricted group to have lower progesterone levels ($P < 0.01$).

The standard and restricted groups showed similar patterns of progesterone concentration during the oestrus cycle. There were no statistically significant differences in progesterone levels between standard and restricted groups but there was variation between individual animals.

Introduction

Cheju Island is situated in the Korean strait, South of the Korean mainland. The island, dominated by Mount Halla which rises to 1950 meters, and has a subtropical climate. It possesses 110,000 hectares of undeveloped grassland which comprises 51% of the total land mass. This area of grassland provides tremendous potential for the grazing of cattle and horses.

During the 1970s there were in excess of 70,000 head of cattle kept on this island but by 1985 this number had been reduced to 45,000. The main reason for this reduction was the poor breeding performance of the native cattle and a dramatic increase in meat consumption.

Kim et al (1980) reported that Cheju native cattle were on average 27.8 months old at first oestrus and the average age at first calving was 38.0 months. When compared with mainland cattle and exotic beef breeds Cheju native cattle were on average 10 months later in reaching maturity. On average 174 days elapsed from calving to first oestrus when bi-annual calving cows were included. Kim et al (1983) stated that the average percentage rate of bi-annual calvers was 15.4%. This poor breeding performance is undoubtedly the main limiting factor to increased cattle numbers, reflected in a loss of farmers income. Poor breeding performance is always thought to be due to under-nutrition and lack of

essential mineral supplementation. Although this conjecture is probably correct it is nevertheless necessary to clarify the true reasons for poor fertility. The development of a satisfactory livestock industry depends upon solving these problems.

Much work has been done on the measurement of progesterone, estradiol, estrone and estrogen in cow blood plasma using radioimmunoassay techniques (Graverick et al., 1970; Chew et al., 1977; Humphery et al., 1983). Henricks et al., (1971) and Gonsales et al., (1975) measured the progesterone and estradiol levels in heifers prior to reaching maturity and Plotka et al. (1967), Sprague et al. (1971) and Swanson et al. (1972) measured progesterone levels during cows oestrus cycle. However in most cases (Wattmann et al., 1972; Christensen et al., 1974; Ireland et al. 1984; Walterus et al., 1984) measured estradiol together with progesterone. Chung et al. (1984) measured hormone levels from the non-pregnant period to early stage of pregnancy.

Eley et al. (1981) measured changes in estradiol and progesterone levels during pre- and postpartum. Considerable work has been carried out on postpartum hormone measurements but little work has been done on prepartum hormone evaluation. Verne La Voie et al. (1981) evaluated the effect of different

calf suckling methods on progesterone levels. Other research works have been done on progesterone and estradiol (Humphery et al., 1983), estradiol – 17 β (Kesler et al., 1979; Goodal et al., 1978) and steroid hormone of postpartum cow. However the aforementioned works only concentrated on physiological aspects and did not include nutritional levels.

To improve the breeding performance of Cheju native cattle the relationship between hormone levels and nutritional levels should be investigated. This investigation was necessary to clarify the reasons for poor fertility, delay in oestrus after parturition and silent heat.

Materials and Methods.

The trial was initiated in June 1984 and 8 Cheju native calves after weaning were randomly assigned in equal numbers on two supplementary feeding levels, these were designed to provide either 100% (standard group) or 70% (restricted group) NRC. All animals in standard group were fed silage, hay and balanced ration over winter period according to NRC feeding standard. On the other hand animals in the restricted group were fed good quality hay but not balanced ration. Heifers in both groups were grazed on improved pasture from May to October. All animals were weighed twice per month at the same time during the experimental period. Rectal palpation was performed on the corpus luteum of ovaries, ovarian size and graffian follicle size at the onset of oestrus in heifers. After parturition ovarian activity, such as the first postpartum ovulation day, corpus luteum diameter, active follicles and ovarian size were determined by rectal examination at 7 days intervals. Blood samples were collected 1400 and 1600 hrs from November 1984 until early 1986.

Blood samples for progesterone and estra-

diol assay were taken at the onset of oestrus cycle and thereafter every 48 hrs until the 19th day and then every 24 hrs until the end of cycle. During the heat period, blood sample were taken 3 times daily.

The collected samples were kept at room temperature for 2-4 hrs and centrifuged at 5,000 r.p.m. after which serum was decanted and stored at -20°C, until radioimmunoassay. Progesterone and estradiol were assayed using Coat-A-Count progesterone and Coat-A-Count Estadiol, products, produced by the Diagnostic Products Corporation, U.S.A.

Results and Discussion

1. Body weights

Table 1 shows body weight changes according to different feeding levels. Mean body weight at 6 months of age was 155kg when fed 100% NRC ration but it was only 137 kg when heifers received the 70% NRC ration. At 10, 15 and 20 months of age the body weight was 66, 160 and 115 kg, respectively, showing that heifers fed the standard ration gained weight rapidly. There was a difference in body weight between the standard and restricted feeding groups. The body weight gain of the restricted group is similar to Sul's (1981a) data on the mainland Korean native cattle (177 kg at 12 months and 229 kg at 24 months). The standard group gave similar body weight gains to Holstein and Korean cross breeds (255 kg at 12 months and 406 kg at 24 months). The body weight gain at weaning of the standard group is much less than the Hereford, Aberdeen Angus, Charolais, and Charolais cross Korean (Sul, 1981b). The small size of the Cheju native cattle is mainly due to the poor feeding management of the local farmer.

Table 1. Body weight change on supplementary feeding levels (kg)

Nutritional level	Cow No.	Body weight at				
		6	10	15	20	25 (months)
NRC 100%	1	127	173	230	317	360
	2	154	219	281	305	370
	3	172	252	230	441	552
	4	167	218	282	352	392
	Mean	155.0	215.5	255.8	353.8	411.0
NRC 70%	1	106	132	179	200	295
	2	144	144	186	193	276
	3	150	167	212	246	323
	4	148	153	165	264	287
	Mean	137.0	149.0	185.5	225.8	295.3

2. Ovarian activity

The ovarian activity judged by palpation at the first oestrus is presented in Table 2. The rectal palpation was performed when female young stock was reached at puberty.

Average size of left ovary in the standard feeding group was 2.1 cm long, and 1.6 cm wide and the right ovary was 2.6 cm long and 1.8 cm wide. However in the restricted feeding group the ovaries were found to be smaller (left 1.5 x

Table 2. Ovarian activities judged by palpation at first oestrus

Nutritional levels	Cow No.	Corpus luteum		Ovarian size (cm)		Follicle size Dia. (mm)		Oestrus condition
		L	R	L	R	5-10	11-15	
NRC 100%	1		1	1.5×1.0	1.8×1.2	1	1	+++
	2	1		2.2×1.8	3.0×2.2		1	+++
	3		1	2.3×2.1	2.8×2.0	1	1	+++
	4		1	2.6×1.5	2.8×1.8	1	1	+++
	Total or Mean		1	3	2.2×1.6	2.6×1.8	3	3
NRC 70%	1		1	2.0×1.5	2.6×1.7	1	1	+++
	2	1		2.0×1.3	2.5×1.5		1	++
	3		1	1.2×1.6	2.0×1.6	1	1	+++
	4		1	1.0×0.8	1.2×0.8	1	1	+++
	Total or Mean		1	3	1.6×1.2	2.6×1.4	3	1

L; Left, R; right, Dia; diameter rl

1.2 cm, right 2.0 x 1.4 cm).

Diameter of Graffian follicles followed a similar tendency to ovarian size in the two groups. Graffian follicles in three heifers were

over 10mm in diameter in the standard group, but only one heifer in the restricted group had follicles in excess of 10mm in diameter. When all heifers fed the standard ration showed normal

oestrus and all conceived.

Two corpus luteum were found in the left ovary and 6 in the right, bearing a close relationship to the findings of Rajakaski (1960). The size of the corpus luteum was larger in the standard group than in the restricted group. This result was consistent with the work of Apgar et al. (1975) who reported higher ovary weights in heifers receiving a high level of nutrition. Ovary sizes in the present trial were found to be similar to the results obtained by Chang and Kim (1982) but were much smaller than those found in Holstein (Rajakoshi, 1960; Salisbury and Van Demark et al., 1961).

The smaller ovaries found in the experimental animals were probably due to the small body

size of Cheju native heifers. Swanson et al. (1972) reported 0.9 cm diameter of ovaries at first oestrus but after recurrence of three consecutive oestrus cycles, the ovaries increased their diameter to 1.2-1.4 cm. Suzuki et al. (1982) studied on the relationship between oestrus and nutritional levels and proved conclusively that low nutrition causes reduction in oestrus length and increases silent heat, leading to infertility.

3. Reproductive performance

Table 3 shows reproductive performance of heifers according to level of supplementary feeding. The first oestrus in the standard feeding group appeared at 14.6 months when the body weight was 265 kg. Age at first pregnancy was

Table 3. Reproductive status and body weight on supplementary feeding levels

Nutritional levels	Cow No.	First oestrus		First pregnancy		First calving	
		Age (month)	Body weight (kg)	Age (month)	Body weight (kg)	Age (month)	Body weight (kg)
NRC 100%	1	13.3	200	13.9	207	23.3	360
	2	13.0	234	15.4	234	25.0	370
	3	12.3	291	26.0	303	36.1	555
	4	19.8	335	21.3	353	30.5	459
	Mean	14.6	265	19.1	274.3	28.9	436.0
NRC 70%	1	20.6	197	25.3	285	34.9	379
	2	26.0	257	36.0	336	45.1	360
	3	25.0	297	27.1	318	44.6	425
	4	20.2	250	21.2	256	30.8	340
	Mean	23.0	250.1	27.4	298.8	38.9	376.0

on average 19.1 months when the average body weight was 274.3 kg and the age at first calving was on average 28.9 months at a body weight of 436 kg. On the other hand when heifers were fed the restricted ration the first oestrus appeared at 23.0 months at a body weight of 250 kg. Average age at first calving was 38.9 months,

which was 10 months later than the average in the standard feeding group.

With exception of one case of distocia in the standard group and a case of retained placenta in the restricted group no other calving disorders were experienced. All calves were born healthy and in good condition.

4. Hormone levels

The progesterone concentration during pregnancy and oestrus cycle at two different feeding levels is shown in Figure 1 and 2. In the standard feeding group the progesterone level was 2ng/ml after two weeks of pregnancy and gradually

increased to 4.9ng/ml at 10 weeks. The level of progesterone peaked at 6.4ng/ml at 8 weeks and this peak was maintained for 24 weeks. After 26 weeks it gradually started to fall back and fell to 2.1 ng/ml at 40 weeks. The progesterone level in the restricted group up to 12 weeks was 5.0




Fig. 1. Change in the serum progesterone concentration in cows during pregnancy on different feeding levels (ng/ml)

ng/ml displaying little difference from the standard group but from 14 weeks the progesterone level in the restricted group dropped, and this reduction (4-4.5ng/ml) was maintained to 36 weeks when the progesterone level drops rapidly over the remaining 4 weeks until it reached 1.5 ng/ml.

When the pregnancy period was divided into three equal parts there were significant differences in hormone levels in the middle section (12-28 weeks) but no hormone difference in part 1 and 3 between the two groups. Few studies so far have been being conducted on the relationship between progesterone level and nutrition. Corah et al. (1974) reported that plasma progesterone immediately prepartum and

postpartum was not affected by energy restriction 100 days prior to calving. Heifers fed high energy rations tend to have slightly higher progesterone levels 3-5 days before calving.

In the present trial progesterone levels were measured throughout the entire pregnancy. Although Randal and Erb (1971) measured progesterone levels without regard to nutritional status their findings were nevertheless similar. Chung et al. (1978, 1984) reported that the progesterone levels at 20-60 dya after conception were 5.9-9.1 ng/ml in Korean native cows. Henricks et al. (1971) stated that the progesterone level in heifers which returned to oestrus 18-20 days after mating was significantly lower than in-calf heifers.

Considerable work has been done on progesterone levels during pregnancy but the major emphasis has been on pre and post-partum. Chew et al. (1977) found that progesterone levels were sigmoid during pregnancy. Shin et al. (1979) found higher progesterone levels (5.1-11.7 ng/ml) from 30 days to 3 days before calving and these levels declined to 1.6 ng/ml at 24 hours prior to calving. Progesterone levels during present trial were slightly lower than those

reported by Shin et al. (1979), Randal and Erb (1971), Henricks et al. (1971), and Ariji et al. (1974) but results obtained by Shemesh et al. (1973) and Corsh et al. (1974) were similar.

Variations in progesterone levels may be due to different hormone assay techniques or use of Coat-A-Count for human diagnosis.

Standard and restricted groups followed similar patterns of progesterone concentration during the oestrus cycle (Fig. 2); progesterone




Fig. 2. Plasma progesterone concentration during oestrus cycle (ng/ml)

concentration peaked 10-12 days in both groups, 2.4 ng/ml in standard and 2.25 ng/ml in restricted groups. These levels were maintained up to 16 days and thereafter the concentration gradually dropped towards the end of the cycle. There were no statistically significant differences in progesterone levels between standard and restricted groups but there was variation between individual animals.

Estradiol measurements were not possible due to the strong interference between bovine and human matrix in the RIA kit supplied. However as E2 levels in cows are higher than

those in humans the samples should be diluted in zero matrix prior to assay.

REFERENCES

1. Apgar, J., D. Aspros, J.E. Hixon, R.R. Saatchman and W. Hansel. 1975. Effect of restricted feed intake on the sensitivity of the bovine corpus luteum to LH in vitro. *J. Anim. Sci.*, 41: 1120-1123.
2. Arije, G.R., J.N. Wiltbank and M.L. Hopwood. 1974. Hormone levels in pre- and

- post-parturient beef cows. *J. Anim. Sci.*, 39: 338-347.
3. Chang, G.C. and J.K. Kim. 1982. Investigation of ovarian size and morphological change in the imported Holstein. *Korean J. Anim. Sci.*, 24: 427-431.
 4. Chew, B.P., H.F. Keller, R.E. Erd and P.V. Malven. 1977. Periparturient concentrations of prolactin, progesterone and the estrogens in blood plasma of cows retaining and not retaining fetal membranes. *J. Anim. Sci.*, 44: 1055-1061.
 5. Christensen, D.S., M.L. Hopwood and J.N. Wiltbank. 1974. Levels of hormones in the serum of cycling beef cows. *J. Anim. Sci.*, 38: 577-583.
 6. Chung, Y.C., C.K. Kim. 1978. Studies on the multiple births in cattle. 1. Study on the early diagnosis of pregnancy in cow. *Korean J. Anim. Sci.*, 20: 342-354.
 7. Chung, Y.C., C.K. Kim and K.S. Lee. 1984. A study on changes in serum hormone level and body weight during estrus and early pregnancy in Korean Native Heifers and Charolais Crossbred Heifers. *Korean J. Anim. Sci.*, 26: 497-508.
 8. Corah, L.R., A.P. Quealy, T.G. Dunn and C.C. Kaltenbach. 1974. Prepartum and postpartum levels of progesterone and estradiol in beef heifers fed two levels of energy. *J. Anim. Sci.*, 39: 380-385.
 9. Eley, D.S., W.W. Thatcher, H.H. Head, Collier, G.J. Wilcox and E.P. Call. 1981. Periparturient and postpartum endocrine changes of conceptus and maternal units in Jersey cows bred for milk yield. *J. Dairy Sci.*, 64: 312-320.
 10. Garverick, H.A., R.E. Erd, G.D. Niswender and C.J. Callahan. 1970. Reproductive steroids in the bovine, III. Changes during the estrous cycle. *J. Anim. Sci.*, 32: 946-955.
 11. Goodale, W.S., H.A. Garverick, D.J. Kesler, C.J. Bierschwal, R.G. Elmore and R.S. Youngquist. 1978. Transitory changes of hormones in plasma of postpartum dairy cows. *J. Dairy Sci.*, 61: 740-746.
 12. Henricks, D.M., D.R. Lamond, J.R. Hill and J.F. Dickey. 1971. Plasma progesterone concentrations before mating and in early pregnancy in the beef heifer. *J. Anim. Sci.*, 33: 450-455.
 13. Humphrey, W.D., C.C. Kaltenbach, T.G. Dunn, D.R. Koritnik and G.D. Niswender. 1983. Characterization of hormonal patterns in the beef cow during postpartum anestrus. *J. Anim. Sci.*, 56: 445-453.
 14. Ireland, J.J., R.L. Fogwell, W.D. Oxender, K. Ames and J.L. Cowley. 1984. Production of estradiol by each ovary during the estrous cycle of cows. *J. Anim. Sci.*, 59: 764-771.
 15. Kesler, D.J., H.A. Garverick, C.J. Bierschwal, R.G. Elmore and R.S. Youngquist. 1979. Reproductive hormones associated with normal and abnormal changes in ovarian follicles in postpartum dairy cows. *J. Dairy Sci.*, 62: 1290-1296.
 16. Kim, J.K. and S.C. Kim. 1980. Studies on the cause of occurrence and the treatments for the reproductive disorder in Jeju Native Cattle. *Korean J. Anim. Sci.*, 22: 161-166.
 17. Kim, J.K., M.C. Kim and S.C. Kim. 1983. Study of feeding and reproductive performance in Jeju Native Cattle. *J. Anim. Sci.*, 77: 87.
 18. Plotka, E.D., R.E. Erb, C.J. Callahan and W.R. Gomes. 1967. Levels of progesterone in peripheral blood plasma during the estrous cycle of the bovine. *J. Dairy Sci.*,

- 50: 1158-1161.
19. Rajakaski, E. 1960. The ovarian follicular system in sexually mature heifers with special reference to seasonal cyclical and right-left variations. *Acta Endocrinol.* 52: 1-68.
 20. Randel, R.D. and R.E. Erb. 1971. Reproductive steroids in the bovine VI. Changes and interrelationships from 0 to 260 days of pregnancy. *J. Anim. Sci.*, 33: 115-123.
 21. Salisbury, G.W. and N.L. VanDemark. 1961. *Physiology of reproduction and artificial insemination of cattle.* Acad. Press. New York and London, pp.23-25.
 22. Shemesh, M., N. Ayalon and H.R. Lindner. 1973. Early pregnancy diagnosis based upon plasma progesterone levels in the cow and ewe. *J. Anim. Sci.*, 36: 726-729.
 23. Shin, W.J., S.H. Chee, Y.B. Lee and J.K. Kwun. 1979. Studies on the changes in serum steroid hormone concentrations and serum metabolite contents during the late pregnancy in Korean Native cow. *Korean J. Anim. Reprod.* 3: 32-41.
 24. Sprague, E.A., M.L. Hopwood, G.D. Niswender and J.N. Wiltbank. 1971. Progesterone and luteinizing hormone levels in peripheral blood of cycling beef cows. *J. Anim. Sci.*, 33: 99-103.
 25. Sul, D.S. 1981 a. Study on change of body weight and measurements of Korean Cattle by crossbreeding of Charolais. *Korean J. Anim. Sci.*, 23: 162-169.
 26. Sul, D.S. 1981 b. Proposal of Korean Native Cattle improvement procedure for production. *Korean J. Anim. Sci.*, 23: 359-398.
 27. Suzuki, O., M. Sato and Y. Kubota. 1982. Effects of underfeeding on estrous behavior and ovarian function in postpubertal beef heifers. *J. Anim. Reprod.*, 28: 204-211.
 28. Swanson, L.V., H.D. Hafs and D.A. Morrow. 1972. Ovarian characteristics and serum LH, prolactin, progesterone and glucocorticoid from first estrus to breeding size in Holstein heifers. *J. Anim. Sci.*, 34: 284-293.
 29. LaVoie, V., D.K. Han, D.B. Foster and E.L. Moody. 1981. Suckling effect on estrus and blood plasma progesterone in postpartum beef cows. *J. Anim. Sci.*, 52: 802-811.
 30. Walters, D.L., D. Schams and E. Schallenberger. 1984. Pulsatile secretion of gonadotrophins, ovarian steroids and ovarian oxytocin during the luteal phase of the oestrous cycle in the cow. *J. Reprod. Fert.*, 71: 479-491.
 31. Wettemann, R.P., H.D. Hafs, L.A. Edgerton and L.V. Swanson. 1972. Estradiol and progesterone in blood serum during the bovine estrous cycle. *J. Anim. Sci.*, 34: 1020-1025.