


A Facile Synthesis of Bis-alkylmercaptomethanephosphonates

Dae Young Kim and Dong Young Oh*

Department of Chemistry, Korea Advanced Institute of Science and Technology, Seoul 131.


Received August 13, 1986

Bis-alkylmercaptomethanephosphonates which are the derivatives of the practically unknown formylphosphonate are the intermediates of the conversion under Wittig-Horner reaction condition into the corresponding ketene thioacetals¹. Ketene thioacetals are key intermediates in a wide variety of organic syntheses².



Until now, bis-alkylmercaptomethanephosphonate have been obtained by the following methods; Arbuzov reaction of trialkyl phosphites with bis-(alkylmercapto) methyl chlorides³, addition of dialkyl sulfide to alkylthiomethanephosphonate carbanion⁴, and reaction of diethoxymethanephosphonate with mercaptans⁵. These known methods are successful to some extent but not sufficient both in yields and in reaction conditions.

In the course of our studies on α -heterosubstituted phosphonates we have recently reported a synthesis of 1-alkoxy-1-arylmethanephosphonates⁶ which involves the Arbuzov reaction of diethyl trimethylsilyl phosphite (DTSP) with arylaldehyde dialkylacetals. In an extension of this work we now wish to report a convenient synthetic method for the preparation of bis-alkylmercaptomethanephosphonates by the reaction of DTSP with orthothioformates.



DTSP reacted with orthothioformates in the presence of Lewis acid highly to yield the desired phosphonates (Table 1). In order to find out optimum condition, we have studied the reaction of DTSP and tributyl orthothioformate in the presence

of four different Lewis acids. The best yield was obtained in stannic chloride. Treatment of orthothioformates with the Lewis acid presumably leads to formation of intermediate thionium ions⁶. In a subsequent nucleophilic reaction DTSP reacts with this ions to yield product.

The advantages of this approach are to give the high yield, to shorten the reaction time and to be easy preparation of starting materials⁷.

The following is a typical experimental procedure. To a solution of orthothioformate (2 mmol) in methylene chloride under nitrogen at -10°C is added dropwise stannic chloride (2.1 mmol). After 10 min stirring at -10°C , DTSP (2 mmol) is added into the reaction mixture. The resulting solution is left to slowly return to room temperature for 3-4h, diluted with methylene chloride (5 ml) and water (5 ml), and stirred for 10 min. The organic layer is separated, dried with magnesium sulfate, and evaporated to leave a yellow or colourless oil. The product was purified by Kugelrohr distillation.

References

- (a) G.C. Kruse, N.L.J. Broekhof, a, Wijsman and A. van der Gen, *Tetrahedron Lett.*, 885 (1977); (b) M. Mikolajczyk, S. Grzejszczak, A. Zatorski and B. Mlotkowska, *Tetrahedron Lett.*, 2731 (1976); (c) M. Mikolajczyk, S. Grzejszczak, A. Zatorski, B. Mlotkowska, H. Gross and B. Costisella, *Tetrahedron*, **34**, 3810 (1978); (d) E. Juaristi, B. Gordillo and L. Valle, *Tetrahedron*, **42**, 1963 (1986).
- (a) D. Seebach, M. Kolb and B.T. Grobel, *Chem. Ber.*, **106**, 2277 (1973); (b) M.S. Chauhan, H. Junjappa, *Synthesis*, 880 (1974); (c) E.J. Corey and A.P. Kozikowski, *Tetrahedron Lett.*, 925 (1975).
- (a) B. Mlotkowska, H. Gross, B. Costisella, M. Mikolajczyk, S. Grzejszczak and A. Zatorski, *J. Prakt. Chem.*, **319**, 17 (1977); (b) H. Gross, I. Keitel, B. Costisella, M. Mikolajczyk and W. Midura, *Phosphorus and Sulfur*, **16**,

Table 1. Preparation of Bis-alkylmercaptomethanephosphonates

R	Lewis Acid	Yield(%) ^a	b.p.(°C/mmHg)	³¹ P-NMR ^b (CDCl ₃ /H ₃ PO ₄) (ppm)
CH ₂ CH ₂ CH ₂ CH ₃	SnCl ₄	92	140-143/0.4	+ 19.14
	TiCl ₄	80		
	BF ₃ Et ₂ O	83		
	ZnI ₂	65		
CH ₂ CH(CH ₃) ₂	SnCl ₄	94	136-38/0.4	+ 19.10
CH ₂ CH ₂ CH ₃	SnCl ₄	88	133-135/0.4	+ 19.50
CH(CH ₃) ₂	SnCl ₄	83	131-134/0.4	+ 19.70
CH ₃ CH ₃	SnCl ₄	85	134-136/0.6 ^c	+ 19.85

^a Isolated yield by Kugelrohr distillation. ^b The conversion of positive ³¹P-NMR to down field from H₃PO₄ is used. ^c Lit⁸; 95-100°C/0.05.

- 257 (1983).
 4. (a) M. Mikolajczak, P. Balczewski and S. Grzejszczak, *Synthesis*, 127 (1980).
 5. D.Y. Kim and D.Y. Oh, *Synthetic Commun.*, 859 (1986).
 6. R.T. Reetz and H. Muller-Strake, *Tetrahedron Lett.*,

3301 (1984).

7. (a) M. Sekine, K. Okimoto, K. Yamada and T. Hata, *J. Org. Chem.*, **46**, 2097 (1981); (b) D. Seebach, K.H. Geiss, A.K. Beck, B. Graf and H. Daum, *Chem. Ber.*, **105**, 3280 (1972).

Formation of *p*-Cymene from the Catalytic Reactions of Unsaturated Aldehydes and Alcohols with a Rhodium(I) Complex

Hyung Mok Jeong, Jeonghan Park, and Chong Shik Chin*


Department of Chemistry, Sogang University, CPO Box 1142, Seoul 121

Received, August 13, 1986

Catalytic cyclization of unsaturated aldehydes to give cycloketones^{1,5} and unsaturated cycloalcohols⁶ with various transition metal complexes has been observed. Aromatization of olefin substituted cyclohexenones by a transition metal complex has been also reported.⁷ In the presence of transition metal catalyst, alcohols and ketones undergo the hydrogenation reaction to give hydrocarbons under H₂,^{8,11} in the presence of hydrogen donor,¹² or simply under N₂ (bimolecular disproportionation reaction, 2R₂CHOH → R₂CH₂ + R₂CO + H₂O).¹³ Dehydrogenation of cyclohexanes with transition metal complexes has been known.^{14,15}

NMR spectra. The reactions also yield some other products (besides *p*-cymene) which have not been fully characterized. It should, however, be mentioned that *p*-isopropenyl toluene (ca. 12%) has been found in the reactions of unsaturated aldehydes, and H₂O in all reactions (eqs. 1-3). It is noticed in eqs. 1-3 that the yield is always close to 50%. Accordingly, it may be suggested that the catalytic reactions (eqs. 1-3) involve a bimolecular disproportionation reaction such as 2R₂CHOH → R₂CH₂ + R₂CO + H₂O as previously reported.¹³
Acknowledgement. We wish to thank the Korea Science and Engineering Foundation and the Ministry of Education for their financial supports.

References



During our investigation on the catalytic activities of M(ClO₄)(CO)(PPh₃)₂ (M = Rh, Ir) for the reactions of unsaturated nitriles,^{16,17} aldehydes¹⁸ and alcohols,¹⁹ we have found the formation of *p*-cymene from the catalytic reactions of the unsaturated aldehydes (neral and geranial) and alcohols (nerol, geraniol and linalool) with Rh(ClO₄)(CO)(PPh₃)₂ (**1**) (eqs. 1-3). In all experiments, 0.7 mmole of **1** and 14 mmoles of the corresponding aldehyde or alcohol were used in 5 ml of CHCl₃ to produce ca. 7.0 mmoles of *p*-cymene. The product, *p*-cymene was identified by GC, mass (m/e = 134) and ¹H-

1. B.R. James and C.G. Young, *J. Organomet. Chem.*, **285**, 321 (1985).
2. R.E. Campbell, Jr., C.F. Lochow, K.P. Vora, and R.G. Miller, *J. Am. Chem. Soc.*, **102**, 5824 (1980).
3. R.C. Larock, K. Oertle, and G.F. Potter, *J. Am. Chem. Soc.*, **102**, 190 (1980).
4. C.F. Lochow and R.G. Miller, *J. Am. Chem. Soc.*, **98**, 1281 (1976).
5. K. Sakai, J. Ide, O. Oda, and N. Nakamura, *Tetrahedron Lett.*, 1287 (1972).
6. K. Sakai and O. Oda, *Tetrahedron Lett.*, 4375 (1976).
7. P.A. Grieco and N. Marinovic, *Tetrahedron Lett.*, 2454 (1978).
8. R. Durand, P. Geneste, C. Moreau, and J.L. Pirat, *J. Catal.*, **90**, 147 (1984).
9. R.K.M.R. Kallury, T.T. Tidwell, D.G.B. Boocock, and D.H.L. Chow, *Can. J. Chem.*, **62**, 2540 (1984).
10. D.J. Drury, M.J. Green, D.J.M. Ray, and A.J. Stevenson, *J. Organomet. Chem.*, **236**, C23 (1982).
11. W.F. Marier, K. Bergman, W. Bleicher, and P.V.R. Schleyer, *Tetrahedron Lett.*, 4227 (1981).
12. G.A. Olah and G.K.S. Prakash, *Synthesis*, 397 (1978).
13. J. Blum, *J. Mol. Catal.*, **3**, 33 (1978).
14. R.H. Crabtree, *J. C. S. Chem. Commun.*, 1829 (1985).
15. D. Brady, M. Ephritikhine, H. Felkin, and R. Homes-Smith, *J. C. S. Chem. Commun.*, 789 (1983).