

- Jpn.*, **54**, 2712 (1981); (c) Y.H. Kim, and B.C. Chung, *J. Org. Chem.*, **48**, 1562 (1983).
 8. E. Lee-Ruff *Chem. Rev.*, **6**, 195 (1977).
 9. M.J. Givian, D.T. Sawyer, R. Tanpoonpholivat, T. Ungerman, and M.M. Morrison, *J. Am. Chem. Soc.*, **101**, 640 (1979).
 10. Actually, sodium sulfite was oxidized with O₂ to sodium

sulfate in quantitative yield.

11. 2-Bromo-4-methylthioacetanilide reacted with O₂ in DMSO at 20°C for 24 h to give the corresponding amide in 5% yield together with the 90% recovery of the starting material, and in 25% yield of amide in the presence of 18-crown-6-ether under the same reaction condition.

An Efficient Route for the Synthesis of Glorin

Yoon-Sik Lee, Soong-Wha Lee, and Woon-Jong Park


Department of Chemical Technology, Seoul National University, Seoul 151. Received October 20, 1986

Chemotaxis is referred to the directed movement of cells toward an attractant along a concentration gradient. Many di- or tripeptides with N-formyl-methionine, which are related to inflammation mechanisms of body are well known examples of chemoattractant for neutrophils and macrophages¹.

Recently, Shimomura *et al.*^{2,3} has isolated a chemotactic peptide, N-propionyl- γ -L-glutamyl-L-ornithine- δ -lactam α -ethyl ester (**1**) (glorin) for social amoeba *Polysphondylium violaceum*, and conformed the structure by comparing it with synthetic glorin for its chemotactic activity. To study the relationship between the structures of various derivatives of glorin and their chemotactic activities on amoeba or leukocytes, we had to synthesize them in large quantity. However the reported route for the synthesis of glorin⁴ appeared somewhat crude without mentioning of reaction condition. Furthermore, no physical data were available except MS and IR spectra².

We now wish to report here a simple and efficient route for the synthesis of glorin and its derivatives. L-Glutamic acid (**2**) was first transformed into N-benzyloxycarbonyl-L-glutamic acid α -ethyl ester (**3**) in the usual manner.^{5,6} L-Ornithine- δ -lactam (**5**) was prepared from L-ornithine methyl ester dihydrochloride (**4**)⁷ by the known procedures⁸. The two fragments, **3** and **5** were coupled by mixed anhydride method⁹ affording N-benzyloxycarbonyl- γ -L-glutamyl-L-ornithine- δ -lactam α -ethyl ester (**6**) in 70% yield. Since **5** was known to be hygroscopic and unstable, 3 eq of **5** was used in the coupling step, and excess of **5** could be removed by washing it with water. The coupled product, **6** was very stable as nice crystalline form¹⁰ and gave satisfactory H-nmr spectral data; NMR (CDCl₃): δ 7.35 (5H, 2, phenyl), 6.64 (1H, d, amide), 5.93 (1H, br, amide), 5.74 (1H, d, amide), 5.11 (2H, s, -CH₂-of benzyl), 4.4-4.0 (2H, m, two methines), 4.20 (2H, q, -OCH₂-), 3.32 (2H, d of t, -CH₂ NH- of lactam), 2.57-1.64 (8H, m, four -CH₂-), 1.27 (3H, t, -CH₃). Treatment of **6** with H₂/Pd in methanol for 7 h, and evaporation of the solvent gave oily product. It was reacted without further purification with 10 eq of propionic anhydride in CH₂Cl₂ for 10 h at room temperature. Removal of the solvent in high vacuum and crystallization of the resulting solid in EtOH-EtOAc afforded **1** in quantitative yield; MP 139-140°C. TLC, Rf 0.69, silica gel (2-butanone-H₂O-HOAc = 7:1.5:1.5), [α]_D²⁰ = +37.77 (c = 0.4, CHCl₃); NMR(CDCl₃), δ 6.93 (1H, br, amide), 6.70 (1H, br, amide), 6.15 (1H, br, amide), 4.7-4.0 (2H, br, two methines), 4.20 (2H,

q, -OCH₂-), 3.37 (2H, m, -CH₂NH- of lactam), 2.6-1.6 (10H, m, five -CH₂-), 1.28 (3H, t, -CH₃), 1.16(3H, t, -CH₃); Anal. calcd. for C₁₅ H₂₅ N₃ O₅: C 55.03, H 7.70, N 12.84; found: C 55.26, H 8.13, N 12.52. Unlike the result of Shimomura *et al.*², **1** was recovered from the reaction mixture in a highly purified



⁷ZnCl₂, H₂O-Et₂O, 0°C, 1.5h and rt, 24h, 91%; ⁸Ac₂O, rt, 17h, 80%; ⁹EtOH, dicyclohexyl amine, rt, 15h, 70%; ¹⁰Dowex 50 WX4, H₂O-MeOH(1:1), rt, 30 min; ¹¹NaOMe, MeOH, rt, 3h, 91%; ¹²for activation, N-methylmorpholine, ClCO₂-isobutyl, -15°C, 5 min; for coupling, -15°C, 30 min and rt, 10h, 70%; ¹³H₂, Pd-C, rt, 7h; ¹⁴propionyl anhydride, CH₂Cl₂, rt, 10h.

form, and no chromatographic purification was necessary. Furthermore, this reaction scheme shows a general route for the synthesis of various kinds of N-substituted Glorin.

Acknowledgement. We wish to thank Ministry of Education for a grant of financial support for this work.

References and Notes

1. E. Schiffmann, B.A. Corcoran, and S.M. Wahl, *Proc. Natl. Acad. Sci. USA*, **72**, 1059 (1975).
2. O. Shimomura, H.L.B. Suthers, and J.T. Bonner, *ibid.*, **79**, 7376 (1982).
3. O. Shimomura, H.L.B. Suthers, and J.T. Bonner, *FEBS Letters*, **155**, 155 (1983).
4. N-Propionyl-L-glutamic acid- α -ethyl- γ -benzyl ester was prepared from L-glutamic acid. Treatment of the diester with HF give N-propionyl-L-glutamic acid α -ethyl ester, which was coupled with **5** yielding **1**. The final

product was purified by column chromatography on Sephadex LH-20 using acetonitrile/water, 9:1 (vol/vol) as the eluent.

5. M. Bergmann and L. Zervas, *Ber.*, **65**, 1192 (1932).
6. E. Klieger, E. Schroder, H. Gibian, *Liebigs Ann. Chem.*, **640**, 157 (1961).
7. K. Golankiewicz, M. Wiewioroski, *Acta. Biochim. Pol.*, **10**, 433 (1964).
8. K. Golankiewicz, M. Dezor, M. Wiewiorowski, *ibid.*, **13**, 1 (1966).
9. G.W. Anderson, J.E. Zimmerman, F.W. Collahan, *J. Am. Chem. Soc.*, **89**, 5012 (1967).
10. The compound, **6** was purified by crystallization in EtOAc. MP 124–126°C; TLC, Rf 0.21, silica gel (CHCl₃, 95% EtOH = 15:1); $[\alpha]_D^{25} = +52.6$ (c = 0.5, CHCl₃); Anal. calcd. for C₂₀H₂₆N₃O₆: C 59.40, H 6.48, N 10.39; found: C 59.38, H 6.63, N 10.03.

A Biogenetic-Type Synthesis of Rose Furan

Eun Lee*, Yong Pyo Hong, Junghun Suh, and Sae-Hee Chang

Department of Chemistry, College of Natural Sciences, Seoul National University, Seoul 151. Received October 20, 1986

Rose furan(**1**) was isolated from Bulgarian rose oil in 1968 and the structure was assigned and confirmed by synthesis by G. Büchi and coworkers¹. Since then it has been isolated from other natural sources².

Following the first synthesis by Büchi¹, many successful schemes were published. In a number of syntheses³⁻⁸, pre-formed furan precursors were used and different strategies for introduction of the prenyl group were investigated. Other synthetic routes involve primarily oxidative furan ring formation of acyclic intermediates⁹⁻¹¹.

In this report, we wish to describe a new biogenetic-type synthesis of rose furan(**1**) from nerol(**2**), one of its probable biogenetic progenitors.

2,3-Epoxynerol(**3**) was prepared by epoxidation of nerol(**2**) with t-butyl hydroperoxide in benzene in the presence of vanadyl acetylacetonate as reported by Sharpless and coworkers¹². The reaction was complete in 3 hours at room temperature and the product **3** was isolated in almost quantitative yield.

The reaction of 2,3-epoxynerol(**3**) with titanium isopropoxide in dichloromethane at room temperature for 30 hours yielded the enediol **4** in 63% overall yield from nerol(**2**).¹³ In the nmr spectrum, a new vinylic proton (H-4) signal appeared at δ 5.36 as a broad triplet and C-5 bisallylic methylene protons gave rise to another broad triplet at δ 2.67.

Partial esterification of **4** with p-toluenesulfonyl chloride in pyridine produced the monotosylate **5** in 89% yield which was cleanly cyclized to yield the epoxide **6** upon treatment with excess sodium hydride in THF for 2 hours. Reaction with methanolic sodium bicarbonate for 36 hours resulted in the formation of several byproducts. In the nmr spectrum of **6**, the characteristic triplet at δ 3.35 was assigned to H-2.

