

## Glutamate-and NMDA-induced calcium influx at synaptosomes and the difference of their actions

Chung-Soo Lee<sup>1</sup>, Jae-Keon Sim, Yong-Kyoo Shin and Kwang-Soo Lee

Department of Pharmacology, College of Medicine and Korea Medical research Institute, Chung-Ang University, Seoul 156-756, Korea

### ABSTRACT

Glutamate and aspartate may evoke an increase in membrane permeability to monovalent cations and  $\text{Ca}^{++}$ . However, it is uncertain whether  $\text{Ca}^{++}$  influx is mediated by voltage dependent  $\text{Ca}^{++}$  channels or by excitatory amino acid activated channels. In addition, the influences of excitatory amino acids on  $\text{Ca}^{++}$  uptake by neuronal tissues as well as the responses of their actions to extracellular  $\text{Mg}^{++}$  concentration are different.

$\text{K}^+$  induced  $\text{Ca}^{++}$  uptake by synaptosomes was dependent on extracellular  $\text{Mg}^{++}$  up to 5 mM and at concentration of 10 mM,  $\text{Ca}^{++}$  influx was rather reduced. In  $\text{Na}^+$  rich media, glutamate-and aspartate-induced  $\text{Ca}^{++}$  uptake was increased by  $\text{Mg}^{++}$  in a dose independent manner. However, the response for NMDA was inhibited by  $\text{Mg}^{++}$  at concentrations above 2 mM.  $\text{K}^+$ -and glutamate-induced  $\text{Ca}^{++}$  influxes were inhibited by 2, 4-dinitrophenol, chlorpromazine and verapamil but not by tetraethylammonium chloride. Tetrodotoxin effectively inhibited the action of glutamate but did not affect that of  $\text{K}^+$ . The response for NMDA was inhibited by 2, 4-dinitrophenol and tetrodotoxin, slightly inhibited by verapamil, and not affected by tetraethylammonium chloride. In  $\text{Na}^+$  rich medium, depolarizing action of glutamate, aspartate and NMDA on synaptosomes was not demonstrated, whereas these agents stimulated  $\text{Ca}^{++}$  uptake and caused  $\text{Ca}^{++}$  influx induced depolarization at mitochondria. On the other hand, the activities of synaptosomal ATPases were not affected by excitatory amino acids at 5 mM. The results suggest that glutamate or NMDA induced  $\text{Ca}^{++}$  influx at synaptosomes exhibits different responses for extracellular  $\text{Mg}^{++}$ . Excitatory amino acids induced  $\text{Ca}^{++}$  influx at synaptosomes may be associated with increased permeability of membrane for  $\text{Na}^+$  and  $\text{Ca}^{++}$  except  $\text{K}^+$  and membrane depolarization due to increased ionic permeability.

**Key Words:** Glutamate, NMDA, Aspartate,  $\text{Ca}^{++}$  influx at Synaptosomes

### INTRODUCTION

Glutamate may be the principal neurotransmitter that mediate fast excitatory synaptic transmission in the vertebrate central nervous system (Watkins and Evans, 1981). It is suggested that glutamate not only has direct excitatory effects on neurones intrinsic to the striatum,

but also stimulates the release of dopamine presumably at presynaptic receptor sites (Giorguieff *et al.*, 1977; Nieoullon *et al.*, 1978; Roberts and Anderson, 1979).

Glutamate seems to activate two or three distinct types of receptors, namely, N-methyl-D-aspartate (NMDA), quisqualate and kainate (Watkins and Evans, 1981; Ishida and Neyton, 1985). All these agonists increase membrane permeability to monovalent cations (Hablitz and Langmoen, 1982; Crunelli *et al.*, 1984), but NMDA also activates a conductance that per-

<sup>1</sup>To whom all correspondences should be addressed.

mits significant calcium influx (MacDermott *et al.*, 1986) which is blocked in a voltage dependent manner by extracellular magnesium (Mayer *et al.*, 1984).

The excitatory action of glutamate upon neurones is shown to be accompanied by a considerable increase of  $aK^+e$  in the spinal cord of the frog (Sonnhof *et al.*, 1975) and in the cerebral cortex of the cat (Heinemann and Pumain, 1980). This increase of  $aK^+e$  is sufficient to depolarize cells in other neurone, causing them to release transmitters.

It is established that depolarization of the plasma membranes of synaptosomes which retain considerably functional integrity with either veratridine or high external  $K^+$  leads to an enhanced uptake of  $Ca^{++}$  and consequent transmitter release (Blaustein, 1975). Since  $Ca^{++}$  uptake induced by  $K^+$  depolarization is unaffected by tetrodotoxin, it appears that this  $Ca^{++}$  uptake does not occur through the  $Na^+$  channel (Nachshen and Blasustein, 1980).

Many of the neurotransmitters, hormones and drugs may modulate the efficiency of neurotransmitter release by modifying the permeability for  $Ca^{++}$  channels through their effects on cAMP-dependent protein kinases and subsequent phosphorylation in specific proteins at the plasma membrane (Reuter, 1983).

On the other hand, recently it is reported that exposure of mammalian brain cells to glutamate or its analogues causes enhanced hydrolysis of inositol phospholipids (Nicoletti *et al.*, 1986) and its metabolite, inositol triphosphate, triggers intracellular  $Ca^{++}$  mobilization (Michell, 1975).

Although it is well known that glutamate stimulates  $Ca^{++}$  uptake in brain slices (Ichida *et al.*, 1982) or in synaptosomes (Retz *et al.*, 1982), the mechanism of action involved in this effect is uncertain. The possibility that the stimulatory effect of glutamate on  $Ca^{++}$  uptake could be due to the effect of cell swelling is suggested (Termer *et al.*, 1950). Since electrophoretic administration of EDTA sometimes induces generalized excitation, chelating effect of glutamate is also proposed (Krnjević, 1970) as the mechanism. On the other hand, the synaptosomal accumulation of  $Ca^{++}$  induced by glutamate may result from decreased  $Ca^{++}$  efflux rather than from increased  $Ca^{++}$  influx. Furthermore, the different responses of gluta-

mate of  $K^+$  induced  $Ca^{++}$  uptake for tetrodotoxin is suggested (Retz and Coyle, 1984). Thus, the action of glutamate on  $Ca^{++}$  transport system at the plasma membrane is obscure.

In the present study, effects of glutamate, aspartate and NMDA on action sites which regulate  $Ca^{++}$  influx at synaptosomes were investigated. For this purpose, effects of inhibitors of membrane phosphorylation, calmodulin inhibitor and specific ionic channel blockers on the actions of excitatory amino acids were studied. Influences of these agents on the membrane potential and activities of synaptosomal ATPases such as  $Na^+-K^+$  ATPase and  $Ca^{++}$ ,  $Mg^{++}$  ATPase were also examined.

## MATERIALS AND METHODS

### Chemicals

Glutamic acid, N-methyl-D-aspartic acid (NMDA), arsenic acid, carbonyl cyanide m-chlorophenylhydrazone (CCCP), chlorpromazine, tetrodotoxin, tetraethylammonium chloride, lanthanum chloride, tris-ATP, choline chloride and arsenazo III were purchased from Sigma Chemical Co.. Verapamil was obtained from Knoll AG; 2, 4-dinitrophenol, aspartic acid,  $MgCl_2$  and  $KCl$  from E. Merck;  $CaCl_2$  and  $NaCl$  from Kanto Chemical Co.; safranin o from Junsei Chemical Co.. Other chemicals were of analytical reagent grade.

### Preparation of rat cerebral synaptosomes

Synaptosomal particles were prepared from rat cerebrum by a modification of the method of Gray and Whittaker (1962). Male Sprague-Dawley rats weighing about 150 g were used. The animals were killed by decapitation. After removing blood clot, the brains were placed in 9 volume of cold 0.32 M sucrose and homogenized with a teflon glass homogenizer. A crude nuclear fraction was removed by centrifugation at 1,000 g for 15 min and the supernatant was transferred and centrifuged at 12,000 g for 60 min. The resulting residue was resuspended in cold 0.32 M sucrose and homogenized with a teflon glass homogenizer. Seven ml of suspension in 0.32 M sucrose (contained about 2 mg protein) was applied into a discontinuous den-

sity gradient consisting of 13 ml layer of 0.8 M sucrose and 13 ml layer of 1.2 M sucrose. After centrifugation in BECKMAN L5-50B-ultracentrifuge using fixed 42.1 Ti rotor at 50,000 g for 2 hr, a synaptosomal fraction was obtained at the interface of the 0.8-1.2 M sucrose layer. The synaptosomal band was broken and centrifuged at 15,000 g for 15 min. The synaptosomal pellet was resuspended in small volume of 0.32 M sucrose and protein concentration was determined by the method of Lowry et al. (1951).

In order to measure  $\text{Ca}^{++}$  uptake and release by synaptosomes, the synaptosomal band was diluted with 1.2 volume of distilled water and centrifuged at 15,000 g for 15 min. The synaptosomal pellet was resuspended in preincubation medium for  $\text{Ca}^{++}$  transport study (124 mM NaCl, 5 mM KCl, 1.3 mM  $\text{MgCl}_2$  and 20 mM Tris HCl, pH 7.4).

#### Measurement of $\text{Ca}^{++}$ uptake by synaptosomes

The rate and extent of  $\text{Ca}^{++}$  uptake by and release from synaptosomes were measured through the absorbance changes of calcium sensitive dye, arsenazo III, at 675-685 nm with a dual wavelength-split beam spectrophotometer (Aminco Chance dual wavelength-split beam recording spectrophotometer, American Instrument Co., U.S.A.) (Ahmed and Connor, 1979; Akerman and Heinonen, 1983). Measurement of  $\text{Ca}^{++}$  uptake by synaptosomes was done in the reaction mixture consisting of 0.75 mg protein/ml synaptosomal particles, 130 mM KCl (or 124 mM NaCl), 1.3 mM  $\text{MgCl}_2$ , 20 mM Tris-maleate, pH 7.4, 100  $\mu\text{M}$  arsenazo III and other compounds. After 10 min preincubation at 30°C, 20  $\mu\text{l}$  of 4.5 mM  $\text{CaCl}_2$  (final 30  $\mu\text{M}$ ) was added to above reaction medium and then absorbance change was measured for 1 min. Finally,  $\text{Ca}^{++}$  uptake by synaptosomes was initiated by 20  $\mu\text{l}$  of 150 mM tris-ATP (final 1 mM) in a 3 ml of total reaction volume and the concentration of  $\text{Ca}^{++}$  was determined.

#### Measurement of membrane potential

The membrane potential was measured with the safranin method (Akerman and Wikaström 1976; Akerman, 1978). The spectrum of safranin changes in a way typical of

stacking upon induction of potentials across the membranes of liposomes or mitochondria. The spectral shift was determined with Aminco Chance dual wavelength-split beam spectrophotometer at 484-524 nm in the same reaction medium as in  $\text{Ca}^{++}$  uptake measuring medium except that it contained 50  $\mu\text{M}$  safranin o.

#### Measurement of ATPase activity

Synaptosomal particles (0.1 mg/ml) were incubated in the reaction medium containing 50 mM imidazole-HCl, pH 7.4; 150 mM NaCl, 5 mM  $\text{MgCl}_2$  and 10 mM KCl for  $\text{Na}^+$ - $\text{K}^+$ ,  $\text{Mg}^{++}$  ATPase; 5 mM  $\text{MgCl}_2$  and 1 mM ouabain for  $\text{Mg}^{++}$  ATPase; 2 mM  $\text{CaCl}_2$ , 5 mM  $\text{MgCl}_2$  and 1 mM ouabain for  $\text{Ca}^{++}$ - $\text{Mg}^{++}$  ATPase. Isoosmolarity of the reaction medium was adjusted with choline chloride. After 5 min preincubation at 37°C, the reaction was initiated by the addition of 0.1 ml of 20 mM tris-ATP (final 2 mM) and the total volume was 1.0 ml. After 10 min incubation at 37°C, the reaction was stopped with 1.0 ml of 15% cold trichloroacetic acid and the mixture was centrifuged at 1,000 g for 10 min. One ml of the supernatant was used for determination of Pi according to the method of Horwitz (1952). ATPase activity was expressed in  $\mu\text{moles Pi}$  liberated per mg protein per hr.

## RESULTS

#### Effects of $\text{Mg}^{++}$ on the excitatory amino acid induced $\text{Ca}^{++}$ uptake

$\text{Mg}^{++}$  have been shown to antagonize the depolarizing action of NMDA on frog motoneurons (Evans *et al.*, 1977; Ault *et al.*, 1980). NMDA induced  $\text{Ca}^{++}$  influx is inhibited by extracellular  $\text{Mg}^{++}$  (MacDermott *et al.*, 1986). The response of glutamate for  $\text{Mg}^{++}$  is more complex because this agonist activates both NMDA receptors and other amino acid receptors linked to a voltage insensitive mechanism (Mayer and Westbrook, 1984). Furthermore, effects of excitatory amino acids on  $\text{Ca}^{++}$  uptake by striatal slices from brain are variable (Retz *et al.*, 1984). Thus, effects of  $\text{Mg}^{++}$  on  $\text{K}^+$ -and excitatory amino acid-induced  $\text{Ca}^{++}$  uptake by synaptosomes were investigated.

Fig. 1 showed that  $\text{K}^+$  induced  $\text{Ca}^{++}$  uptake




Fig. 1.  $K^+$  induced  $Ca^{++}$  uptake in the presence of  $Mg^{++}$ ,  $Ca^{++}$  transport at synaptosomes was spectrophotometrically measured through the absorbance changes of  $Ca^{++}$  sensitive dye, arsenazo III, at 675–685 nm. The reaction mixture contained 0.75 mg/ml of synaptosomes, varying concentration of  $MgCl_2$ , 130 mM KCl, 5 mM NaCl, 30  $\mu$ M  $CaCl_2$ , 100  $\mu$ M arsenazo III and 20 mM Tris-maleate, pH 7.4. Each point represents an average of 5 experiments.




Fig. 2. Effect of  $Mg^{++}$  on excitatory amino acids induced  $Ca^{++}$  uptake in 96 mM NaCl and 44 mM KCl containing medium. Each point represents an average of 4–6 experiments. Concentration of excitatory amino acid is 5 mM. ●, no addition; ○, glutamate; Δ, aspartate; □, NMDA.

by synaptosomes was stimulated by  $Mg^{++}$  at the concentration range of 0.5 mM–5.0 mM and at 10 mM of  $Mg^{++}$ ,  $Ca^{++}$  influx was reduced rather than increased. On the other hand, at  $Mg^{++}$  concentrations below 0.5 mM the release of  $Ca^{++}$  from synaptosomes was observed.

To explore the difference between response of  $K^+$  induced  $Ca^{++}$  uptake for  $Mg^{++}$  and that of excitatory amino acid induced  $Ca^{++}$  uptake for  $Mg^{++}$ , effects of  $Mg^{++}$  on glutamate, aspartate and NMDA induced  $Ca^{++}$  uptakes were observed in  $Na^+$ -rich medium. In reaction medium containing 44 mM KCl, 96 mM NaCl and  $MgCl_2$  less than 1.0 mM,  $Ca^{++}$  release rather than uptake was observed. Accordingly, the presence of  $Mg^{++}$  may be required for glutamate or aspartate induced  $Ca^{++}$  uptake, however, these stimulatory effects were reduced by high concentration (10 mM) of  $Mg^{++}$ . Effect of NMDA was inhibited by  $Mg^{++}$  at concentrations above 2 mM (Fig. 2).

Similar findings were also investigated in  $Na^+$ -rich (124 mM) medium. Fig. 3 illustrated that NMDA induced response was significantly inhibited by  $Mg^{++}$ . Thus, it is suggested that  $Ca^{++}$  concentration.




Fig. 3. Effect of  $Mg^{++}$  on excitatory amino acids induced  $Ca^{++}$  uptake in 132 mM NaCl and 5 mM KCl containing medium. Each point represents an average of 4–6 experiments. Concentration of excitatory amino acid is 5 mM. ●, no addition; ○, glutamate; Δ, aspartate; □, NMDA.

### Effects of inhibitors of phosphorylation and ion channel blockers on K<sup>+</sup> induced Ca<sup>++</sup> uptake

Changes of ion permeability of synaptosomes retaining functional integrity during K<sup>+</sup> induced depolarization are relatively not clarified. Thus, character of Ca<sup>++</sup> transport in K<sup>+</sup> induced depolarization was examined.

As can be seen in Table 1, K<sup>+</sup> induced Ca<sup>++</sup>

**Table 1.** Effects of inhibitors of membrane phosphorylation and ion channel blockers on K<sup>+</sup> induced Ca<sup>++</sup> uptake

| Additions         |        | Ca <sup>++</sup> uptake during 2 min<br>nmol/mg protein |
|-------------------|--------|---|
| None              |        | 4.14 ± 0.67   |
| Dinitrophenol     | 10 μM  | 1.70 ± 0.23   |
| Arsenate          | 10 μM  | 1.78 ± 0.17   |
| CCCP              | 1 μM   | 1.58 ± 0.28   |
| CPZ               | 10 μM  | 3.47 ± 0.16   |
| TTX               | 10 μM  | 4.23 ± 0.27   |
| TEA               | 100 μM | 4.11 ± 0.11   |
| La <sup>+++</sup> | 10 μM  | 3.50 ± 0.15   |
| Verapamil         | 0.5 mM | 2.96 ± 0.24   |

The reaction mixtures were the same as described in Fig. 1. Each value represents mean ± S.E. of 6 experiments. The agents were present during preincubation period of 10 min.

**Table 2.** Effects of inhibitors of membrane phosphorylation and ion channel blockers on excitatory amino acids induced Ca<sup>++</sup> uptake

| Additions     | Ca <sup>++</sup> uptake during 2 min (nmol/mg protein) |             |             |
|---------------|--|-------------|-------------|
|               | Without  | + Glutamate | + NMDA      |
| None          | 2.63 ± 0.16  | 6.08 ± 0.12 | 2.77 ± 0.18 |
| Dinitrophenol | 10 μM  | 2.09 ± 0.23 | 3.58 ± 0.48 |
| CPZ           | 100 μM   | 2.67 ± 0.19 | 5.03 ± 0.10 |
| TTX           | 10 μM  | 2.37 ± 0.24 | 3.54 ± 0.07 |
| TEA           | 1 mM   | 2.34 ± 0.19 | 4.91 ± 0.29 |
| Verapamil     | 0.5 mM   | 1.82 ± 0.13 | 3.40 ± 0.21 |

The reaction mixture contained 0.75 mg/ml of synaptosomes, 96 mM NaCl, 44 mM KCl, MgCl<sub>2</sub> (2 mM for glutamate, 1.3 mM for NMDA), 30 μM CaCl<sub>2</sub>, 100 μM arsenazo III and 20 mM Tris-maleate, pH 7.4 with 5 mM of amino acid or not. Each value represents mean ± S.E. of 6 experiments. The agents were present during preincubation period of 10 min.

uptake was remarkably inhibited by inhibitors of protein phosphorylation, such as 2,4-dinitrophenol, arsenate and CCCP. Therefore, these results suggest that phosphorylation of protein at the plasma membrane opens K<sup>+</sup> activated Ca<sup>++</sup> channels. K<sup>+</sup> induced Ca<sup>++</sup> uptake was inhibited by a calmodulin inhibitor, chlorpromazine and Ca<sup>++</sup> channel blocker, La<sup>+++</sup> and verapamil but not by Na<sup>+</sup> channel blocker, La<sup>+++</sup> and verapamil but not by Na<sup>+</sup> channel blocker tetrodotoxin and K<sup>+</sup> channel blocker, tetraethylammonium chloride.

Thus, it is suggested that K<sup>+</sup> may activate action sites which are directly linked to Ca<sup>++</sup> channels.

### Effects of inhibitors of phosphorylation and ion channel blockers on excitatory amino acid induced Ca<sup>++</sup> uptake

Although opening of Ca<sup>++</sup> channels during membrane depolarization and the depolarizing action of excitatory amino acids are well known, the conductance mechanism underlying their depolarizing action has not been clearly defined. In the present study, Ca<sup>++</sup> transport process stimulated by excitatory amino acids was compared with K<sup>+</sup> induced Ca<sup>++</sup> uptake.

The results represented in Table 2 showed that glutamate-induced Ca<sup>++</sup> uptake was inhibited by 2, 4-dinitrophenol, tetrodotoxin and verapamil but not by chlorpromazine and tetraethylammonium chloride. NMDA induced Ca<sup>++</sup> influx was inhibited by 2, 4-dinitrophenol and

tetrodotoxin, and slightly reduced by verapamil.




Fig. 4. Effects of excitatory amino acids on synaptosomal membrane potential. The spectral shift of positive charged safranin o was measured with dual wavelength-split beam spectrophotometer at 484–524 nm. The reaction mixtures were same as described in Table 2. a, no addition ; b, glutamate ; c, NMDA ; d, aspartate.




Fig. 5. Effects of excitatory amino acids on energized mitochondrial  $Ca^{++}$  uptake.  $Ca^{++}$  uptake was spectrophotometrically measured at 675–685 nm. The reaction mixture contained 0.5 mg/ml of mitochondria, 5 mM succinate, 250 mM sucrose, 100  $\mu$ M arsenazo III and 20 mM HEPES-tris, pH 7.4. a, no addition ; b, 1 mM glutamate ; c, 5 mM NMDA ; d, 5 mM aspartate.

### Effects of excitatory amino acids on membrane potential

Since isolated synaptosomes have a potassium diffusion potentials similar to those found in intact nerve cells and it is suggested that they contain specific mechanisms for the transport of  $Ca^{++}$ ,  $Na^{+}$  and  $K^{+}$  across the plasma membrane (Blaustein *et al.*, 1972; Swanson *et al.*, 1974; Blaustein, 1975), effects of excitatory 1974; Blaustein, 1975), effects of excitatory amino acids on ionic permeability of synaptosomes were investigated in relation to their depolarizing actions. Depolarizing action of glutamate, aspartate and NMDA was observed  $Na^{+}$  and  $K^{+}$ , but their action was not detected in the medium containing 44 mM KCl and 90




Fig. 6. Effects of excitatory amino acids on mitochondrial membrane potential. The spectral shift of positive charged safranin o was measured at 484–524 nm. The reaction mixtures contained safranin o instead of arsenazo III were the same as described in Fig. 4. S indicates succinate. a, no addition ; b, 5 mM glutamate ; c, 5 mM glutamate plus 10  $\mu$ M 2, 4-dinitrophenol; d, 10  $\mu$ M 2,4-dinitrophenol; e, 5 mM glutamate plus 5  $\mu$ M ruthenium red ; f, 5  $\mu$ M ruthenium red.

**Table 3.** Effects of excitatory amino acids on synaptosomal  $\text{Na}^+-\text{K}^+$ ,  $\text{Mg}^{++}$  ATPase,  $\text{Mg}^{++}$  ATPase and  $\text{Ca}^{++}-\text{Mg}^{++}$  ATPase activities

| Additions | $\text{Na}^+-\text{K}^+$ , $\text{Mg}^{++}$ ATPase | $\text{Mg}^{++}$ ATPase | $\text{Ca}^{++}$ , $\text{Mg}^{++}$ ATPase |
|-----------|--|-------------------------|--|
| None      | $20.08 \pm 0.26$                                   | $10.46 \pm 0.16$        | $10.90 \pm 0.21$                           |
| Glutamate | $20.72 \pm 0.29$                                   | $9.90 \pm 0.32$         | $11.03 \pm 0.22$                           |
| Aspartate | $22.24 \pm 0.28$                                   | $10.59 \pm 0.38$        | $11.98 \pm 0.45$                           |
| NMDA      | $20.69 \pm 0.13$                                   | $10.78 \pm 0.11$        | $11.11 \pm 0.32$                           |

Concentration of excitatory amino acid is 5 mM. Experimental conditions were the same as described in Materials and Methods. All values are expressed in  $\mu\text{mol Pi/mg protein/hr}$  and represent mean  $\pm$  S.E. of 6 experiments.

mM NaCl (Fig. 4). Thus, excitatory amino acids induced depolarization was studied with mitochondria, in which the spectral shift to safranin, the cationic membrane potential probe was well established.

The driving force of mitochondrial  $\text{Ca}^{++}$  transport seems to be a membrane potential with negative intramitochondrial polarity (Akerman, 1978). As can be seen in Fig. 5 and 6, excitatory amino acids stimulated  $\text{Ca}^{++}$  uptake in energized conditions and facilitated a  $\text{Ca}^{++}$  uptake in energized conditions and facilitated a  $\text{Ca}^{++}$  influx mediated depolarization which was effectively prevented by ruthenium red and 2, 4-dinitrophenol. Thus, the result indicated that glutamate, aspartate and NMDA increase  $\text{Ca}^{++}$  permeability of the mitochondrial membrane.

#### Effects of excitatory amino acids on $\text{Na}^+-\text{K}^+$ , $\text{Mg}^{++}$ ATPase activities

Action of excitatory amino acids on regulatory apparatus of ion transport across the plasma membrane was examined. Glutamate, aspartate and NMDA at concentration of 5 mM did not affect the activities of  $\text{Na}^+-\text{K}^+$ ,  $\text{Mg}^{++}$  ATPase,  $\text{Mg}^{++}$  ATPase and  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  ATPase (Table 3).

## DISCUSSION

Excitatory amino acids such as L-glutamate and L-aspartate are thought to act as fast synaptic transmitters, the ionic mechanism of which is not well understood. It has been suggested that neurones of the mammalian central nervous system possess several different receptors including NMDA receptor for a-

cidic amino acids (Watkins and Evans, 1981), which in turn are coupled to separate conductance mechanisms (MacDonald and Poriatis, 1982). Excitatory amino acids may cause neuronal depolarization by increasing the membrane permeability to  $\text{Na}^+$  (Bührle and Sonnhof, 1983). Glutamate evokes influx of  $\text{Na}^+$  and  $\text{Ca}^{++}$  into motoneurons (Bührle and Sonnhof, 1983). NMDA receptors gate channels that are permeable to  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{++}$  (Mayer and Westbrook, 1985). The voltage dependence of the NMDA receptor-linked conductance is reported (Dingledine, 1983). Various experiments suggest that  $\text{Ca}^{++}$  influx is also associated with the activation of excitatory amino acid receptors on vertebrate neurones (Pumain and Heinemann, 1985). Thus, whether  $\text{Ca}^{++}$  enters through voltage dependent  $\text{Ca}^{++}$  channels or through excitatory amino acid activated channels of one or more subtype is unclear.

$\text{Mg}^{++}$  depresses depolarizing responses of motoneurons of the immature rat evoked by carbachol, norepinephrine and substance p (Ault *et al.*, 1980). It has also been reported that  $\text{Mg}^{++}$  depresses acetylcholine-as well as amino acid-induced excitation in the cat central nervous system (Davis and Watkins, 1977). The excitatory action of acidic amino acids acting at NMDA receptors is remarkably sensitive to the membrane potential and it has been suggested that the NMDA receptor is coupled to a voltage sensitive conductance (Mayer *et al.*, 1984). The blockade of inward current flow through ion channels linked to NMDA receptors by  $\text{Mg}^{++}$  is observed (Ault *et al.*, 1980). Glutamate and aspartate also induce a voltage dependent conductance increase (Nowak *et al.*, 1984; Pumain and heinemann, 1985). However, the responses of glutamate, aspartate and

NMDA induced  $\text{Ca}^{++}$  uptake by neuronal tissues for extracellular  $\text{Mg}^{++}$  concentration are different. Thus, in synaptosomes of rat brain the changes of  $\text{K}^{+}$ -and excitatory amino acids-induced  $\text{Ca}^{++}$  uptake by  $\text{Mg}^{++}$  were examined.

The result represented in Fig. 1 indicates that in  $\text{K}^{+}$  rich medium  $\text{Mg}^{++}$  have not inhibitory effect on  $\text{Ca}^{++}$  flux into synaptosomes because  $\text{K}^{+}$  induced  $\text{Ca}^{++}$  uptake by synaptosomes was stimulated by extracellular  $\text{Mg}^{++}$  at concentrations of 0.5 mM through inhibitory action of  $\text{Mg}^{++}$  was observed at 10 mM. This finding does not correspond with the report of Nachshen (1984). Moreover, in  $\text{Mg}^{++}$  free medium release of  $\text{Ca}^{++}$  from synaptosomes occurred. Thus, it is suggested that  $\text{K}^{+}$  induced  $\text{Ca}^{++}$  uptake is dependent upon  $\text{Mg}^{++}$  which activates ATPases such as  $\text{Na}^{+}\text{-K}^{+}$  ATPase and  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  ATPase. Fig. 3 shows that although stimulatory effect of glutamate is rather reduced by high concentration of  $\text{Mg}^{++}$  compared to low concentration, glutamate induced  $\text{Ca}^{++}$  uptake may be dependent upon extracellular  $\text{Mg}^{++}$ . However, actions of NMDA on  $\text{Ca}^{++}$  influx in  $\text{Na}^{+}$  rich media were antagonized by extracellular  $\text{Mg}^{++}$ . The results suggest that  $\text{Ca}^{++}$  influx associated with NMDA actions may be inhibited by  $\text{Mg}^{++}$  induced voltage dependent channel block, whereas glutamate affects other ion transport systems than the voltage dependent channel.

Since action of glutamate on  $\text{Ca}^{++}$  transport system at synaptosomes was uncertain, effects of uncouplers of membrane phosphorylation and specific ion channel blockers on glutamate induced  $\text{Ca}^{++}$  uptake were investigated, along with their effects on  $\text{K}^{+}$  induced  $\text{Ca}^{++}$  influx. Glutamate-and  $\text{K}^{+}$ -induced  $\text{Ca}^{++}$  influx was inhibited by 2, 4-dinitrophenol, inhibitor of membrane phosphorylation and verapamil, a  $\text{Ca}^{++}$  channel blocker but not by tetraethylammonium chloride, a  $\text{K}^{+}$  channel blocker. On the other hand, tetrodotoxin, a  $\text{Na}^{+}$  channel blocker effectively inhibited glutamate's action, but tetrodotoxin did not affect  $\text{K}^{+}$  induced  $\text{Ca}^{++}$  influx. The insensitivity of glutamate to tetrodotoxin in the present experiment is in agreement with earlier studies in striatal synaptosomes from the rat (Retz *et al.*, 1982). Therefore, Table 1 and 2 suggest that the increased permeability for  $\text{Ca}^{++}$  by glutamate and  $\text{K}^{+}$  may be mediated through their actions on membrane phosphorylation, and these

agents activate divalent cation selective pathways for  $\text{Ca}^{++}$  entry. Stimulated  $\text{Ca}^{++}$  uptake by glutamate may be partially attributable to the increased  $\text{Na}^{+}$  permeability and to the membrane depolarization because tetrodotoxin inhibits  $\text{Ca}^{++}$  influx by glutamate. NMDA induced  $\text{Ca}^{++}$  influx change was also reduced by 2, 4-dinitrophenol and tetrodotoxin but slightly inhibited by verapamil. Some experiments reported that  $\text{Ca}^{++}$  influx is associated with the activation of excitatory amino acid receptors and excitatory amino acids increase the intracellular  $\text{Ca}^{++}$  activity by activation of NMDA receptor channels rather than actions on voltage dependent  $\text{Ca}^{++}$  channels (MacDermott *et al.*, 1986). Accordingly, although opening of voltage dependent  $\text{Ca}^{++}$  channel chiefly regulates NMDA induced  $\text{Ca}^{++}$  influx, activation of NMDA receptor channels in part also may have a minor role.

The preceding article shows that synaptosomes may have membrane potentials which behave like  $\text{K}^{+}$  diffusion potentials. It is established that veratridine and high  $\text{K}^{+}$ , which increase  $\text{Ca}^{++}$  transport across membranes, through a mechanism involving membrane depolarization (Blaustein, 1975). The experiment of Cho *et al.* (1988) showed that in the reaction mixture containing  $\text{Ca}^{++}$  without  $\text{K}^{+}$  the addition of ATP induced the synaptosomal  $\text{Ca}^{++}$  uptake only slightly. The further addition of  $\text{K}^{+}$  is followed by a marked increase in calcium uptake with concomitant development of membrane depolarization. The depolarizing action of glutamate and NMDA can be easily demonstrated in electrophysiological studies using either cultured neurones or brain slices. However, in the present reaction medium containing  $\text{Na}^{+}$  and  $\text{K}^{+}$  depolarizing actions of glutamate, aspartate and NMDA would not be detected by the safranin method. It is reported that glutamate causes depolarization of synaptosomes in the presence of  $\text{Ba}^{++}$  or 4-aminopyridine, whereas in  $\text{Na}^{+}$  rich media it causes no or insignificant changes of membrane potential (Akerman *et al.*, 1987). thus, this finding suggests that outflow of  $\text{K}^{+}$  limits the membrane potential changes induced by glutamate and the changes of safranin spectrum shift in synaptosomes by excitatory amino acids may be relatively smaller than  $\text{K}^{+}$  and measurement of excitatory amino acids induced depolarization using safranin is limited. In addition, examination for sensitivity differences between



safranine and cyanine dye in synaptosomal depolarization may be necessary. The changes in membrane potential during  $\text{Ca}^{++}$  influx across the mitochondrial membrane are well demonstrated using safranine. The extent of safranine spectral change correlates linearly to the magnitude of membrane potential at least up to 200 mV and thus accurate estimates of the mitochondrial membrane potential in different conditions can be obtained (Akerman and Wikström, 1976). As can be seen in Fig. 5 and 6, glutamate, aspartate and NMDA stimulated mitochondrial  $\text{Ca}^{++}$  uptake and induced  $\text{Ca}^{++}$  diffusion potential. Accordingly, depolarizing action of excitatory amino acids is investigated as the changes of mitochondrial membrane potential.

Finally, increases of intracellular  $\text{Ca}^{++}$  ion concentration evoked by excitatory amino acids are not explained by their action on synaptosomal  $\text{Na}^{+}\text{-K}^{+}$  ATPase and  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  ATPase.

The results suggest that excitatory amino acids induced  $\text{Ca}^{++}$  influx may be associated with increased permeability of membrane to ions other than  $\text{K}^{+}$ .

## REFERENCES

- Ahmed Z and Connor JA: *Measurement of calcium influx under voltage clamp in molluscan neurones using the metallochromic dye arsenazo III.* *J Physiol* 286:61-82, 1979
- Akerman KEO: *changes in membrane potential during calcium ion influx and efflux across the mitochondrial membrane.* *Biochim Biophys Acta* 502: 359-366, 1978
- Akerman KEO and Heinonen E: *Qualitative measurements of cytosolic calcium ion concentration within isolated guinea pig nerve endings using entrapped arsenazo III.* *Biochim Biophys Acta* 732:117-121, 1983
- Akerman KEO, Scott IG, Heikkilä JE and Heinonen E: *Ionic dependence of membrane potential and glutamate receptor-linked responses in synaptoneuroosomes as measured with a cyanine dye, Dis-C<sub>2</sub>-(5).* *J Neurochem* 48: 522-559, 1987
- Akerman KEO Wikström MK: *Safranine as a probe of the mitochondrial membrane potential.* *FEBS Lett* 68(2):191-197, 1976
- Ault B, Evans RH, Francis AA, Oakes DJ and Watkins JC: *Selective depression of excitatory amino acid induced depolarizations by magnesium ions in isolated spinal cord preparations.* *J Physiol* 307:413-428, 1980
- Blaustein MP: *Effects of potassium, veratridine and scorpion venom on calcium accumulation and transmitter release by nerve terminals in vitro.* *J Physiol* 247:617-655, 1975
- Blaustein MP, Johnson EMJr and Needleman P: *Calcium-dependent norepinephrine release from presynaptic nerve endings in vitro.* *Proc Natl Acad Sci USA* 69:2237-2240, 1972
- Bührle ChPh and Sonnhof U: *The ionic mechanism of the excitatory action of glutamate upon the membranes of motoneurons of the frog.* *Pflügers Arch* 396:154-162, 1983
- Cho SJ, Lee CS, Shin YK and Lee KS: *Modulation of synaptosomal calcium transport by catecholamines through adrenergic receptors.* *Chung-Ang J Med in press*
- Crunelli V, Forda S and Kelly JS: *The reversal potential of excitatory amino acid action on granule cells of the rat dentate gyrus.* *J Physiol* 351:327-342, 1984
- Davis J and Watkins JC: *Effects of magnesium ions on the responses of spinal neurones to excitatory amino acids and acetylcholine.* *J Physiol* 124: 370-384, 1977
- Dingledine R: *N-methyl aspartate activates voltage dependent calcium conductance in rat hippocampal pyramidal cells.* *J Physiol* 343:385-405, 1983
- Evans RH, Francis AA and Watkins JC: *Selective antagonism by  $\text{Mg}^{2+}$  of amino acid-induced depolarization of spinal neurones.* *Experientia* 33:489-491, 1977
- Giorguieff MR, Kenel ML and Glowinski J: *Presynaptic effects of L-glutamic acid on the release of dopamine in rat striatal slices.* *Neurosci Lett* 6:73-77, 1977
- Gray EG and Whittaker VP: *The isolation of nerve endings from brain: an electron-microscopic study of cell fragments derived by homogenization and centrifugation.* *J Anat* 96:79-87, 1962
- Hablitz JJ and Langmoen IA: *Excitation of hippocampal pyramidal cells by glutamate in the guinea pig and rat.* *J Physiol* 325:317-331, 1982
- Heinemann U and Pumain R: *Extracellular calcium activity changes in cat sensorimotor cortex induced by iontophoretic application of amino acids.* *Exp Brain Res* 40:247-250, 1980
- Horwitt BN: *Determination of inorganic serum phosphate by means of stannous chloride.* *J Biol Chem* 199:537-541, 1952

- Ichida S, Tokunaga H, Moriyama M, Oda Y, Tanaka S and Kita T: *Effects of neurotransmitter candidates on <sup>45</sup>Ca uptake by cortical slices of rat brain: stimulatory effect of L-glutamic acid.* *Brain Res* 248:305-311, 1982
- Ishida AT and Neyton J: *Quisqualate and L-glutamate inhibit retinal horizontal-cell responses to kainate.* *Proc Natl Acad Sci USA* 82:1837-1841, 1985
- Krnjevic K: *Glutamate and  $\gamma$ -aminobutyric acid in brain.* *Nature* 228:119-124, 1970
- Lowry OH, Rosebrough NJ, Farr AL and Randal RJ: *Protein measurement with the folin phenol reagent.* *J Biol Chem* 193:266-275, 1951
- MacDermott AB, Mayer ML, Westbrook GL, Smith SJ and Barker JL: *NMDA-receptor activation increases cytoplasmic calcium concentration in cultured spinal cord neurones.* *Nature* 321:519-522, 1986
- MacDonald JF and Poriectis AV: *DL-Quisqualic and L-aspartic acids activate separate excitatory conductances in cultured spinal cord neurones.* *Brain Res* 245:175-178, 1982
- Mayer ML and Westbrook GL: *Mixed-agonist action of excitatory amino acids on mouse spinal cord neurones under voltage clamp.* *J Physiol* 354:29-53, 1984
- Mayer ML and Westbrook GL: *The action of N-methyl-D-aspartic acid on mouse spinal neurones in culture.* *J Physiol* 361:65-90, 1985
- Mayer ML, Westbrook GL and Guthrie PB: *Voltage-dependent block by Mg<sup>2+</sup> of NMDA responses in spinal cord neurones.* *Nature* 309:261-263, 1984
- Michell RH: *Inositol phospholipids and cell surface receptor function.* *Biochim Biophys Acta* 415:81-147, 1975
- Nachshen DA: *Selectivity of the Ca binding site in synaptosome Ca channels: Inhibition of Ca influx by multivalent metal cations.* *J Gen Physiol* 83:941-967, 1984
- Nachshen DA and Blaustein MP: *Some properties of potassium-stimulated calcium influx in pre-synaptic nerve endings.* *J Gen Physiol* 76:709-728, 1980
- Nicoletti F, Meek JL, Iadarola MJ, Chuang DM, Roth BL and Costa E: *Coupling of inositol phospholipid metabolism with excitatory amino acid recognition sites in rat hippocampus.* *J Neurochem* 46:40-46, 1986
- Nieoullon A, Cherany A and Glowinski J: *Release of dopamine evoked by electrical stimulation of the motor and visual areas of the cerebral cortex on both caudate nuclei and in the substantia nigra in the cat.* *Brain Res* 145:69-83, 1978
- Nowak L, Bregestovski P, Ascher P, Herbel A and Prochiantz A: *Magnesium gates glutamate-activated channels in mouse central neurones.* *Nature* 307:462-465, 1984
- Pumain R and Heinemann U: *Stimulus and amino acid induced calcium and potassium changes in rat neocortex.* *J Neurophysiol* 53:1-16, 1985
- Retz KCD and Coyle JT: *The differential effects of excitatory amino acids on uptake of <sup>45</sup>CaCl<sub>2</sub> by slices from mouse striatum.* *Neuropharmacol* 23:89-94, 1984
- Retz KC, Young AC and Coyle JT: *Glutamate stimulation of <sup>45</sup>Ca uptake by rat striatal synaptosomes.* *Eur J Pharmacol* 79:319-322, 1982
- Reuter H: *Calcium channel modulation by neurotransmitters, enzymes and drugs.* *Nature* 301:569-574, 1983
- Roberts PJ and Anderson SD: *Stimulatory effect of L-glutamate and related amino acids on [<sup>3</sup>H] dopamine release from rat striatum. An in vitro model for glutamate actions.* *J Neurochem* 32:1539-1545, 1979
- Sonnhof U, Grafe P, Krumnikl J, Linder M and Schindler L: *Inhibitory postsynaptic actions of taurine, GABA and other amino acids of motoneurons of the isolated frog spinal cord.* *Brain Res* 100:327-341, 1975
- Swanson PD, Anderson L and Stahl WL: *Uptake of calcium ions by synaptosomes from rat brain.* *Biochim Biophys Acta* 356:174-183, 1974
- Terner C, Eggleston LV and Krebs HA: *The role of glutamic acid in the transport of potassium in brain and retina.* *Biochem J* 47:139-140, 1950
- Watkins JC and Evans RH: *Excitatory amino acid transmitters.* *Ann Rev Pharmacol Toxicol* 21:165-204, 1981

## == 국문초록 ==

### Glutamate와 NMDA에 의한 Synaptosome에서의 칼슘 유입과 이들의 작용의 차이

중앙대학교 의과대학 약리학교실 및 한국의학연구소

이정수 · 심재건 · 신용규 · 이광수

Glutamate와 aspartate는 단가 양이온과 칼슘에 대한 세포막의 투과성을 증가시키는 것으로 시사되고 있다. 그러나 칼슘 유입이 voltage에 의존적인 칼슘 통로에 의하여 또는 흥분성 아미노산에 활성적인 통로에 의하여 이루어지는가는 분명하지 않다. 더욱이, 신경세포의 칼슘 유입에 미치는 흥분성 아미노산의 영향과 세포의 마그네슘에 대한 이들의 반응이 다른 것으로 추정하고 있다.

Synaptosome에서 포타슘에 의한 칼슘 흡수는 세포의 마그네슘에 의존적이었으나 10 mM 농도에서는 그 이하의 농도에서보다 오히려 감소하였다. 소듐이 주된 반응액에서 glutamate와 aspartate에 의한 칼슘 흡수는 마그네슘에 의하여 용량에 비존적인 양상으로 증가하였다. 그러나 NMDA의 작용은 2 mM 이상의 마그네슘에 의하여 억제되었다. 포타슘과 glutamate에 의한 칼슘 흡수는 2,4-dinitrophenol, chlorpromazine과 verapamil에 의하여 억제되었으나 tetraethylammonium chloride의 영향은 받지 아니하였다. Tetrodotoxin은 효과적으로 glutamate의 작용을 억제하였으나 K<sup>+</sup>의 작용에는 영향을 주지 않았다. NMDA의 작용은 2,4-dinitrophenol과 tetrodotoxin에 의하여 억제되었고 verapamil에 의하여 약간 억제되었으며 tetraethylammonium chloride의 영향은 받지 아니하였다. 소듐이 주된 반응액에서 glutamate, aspartate와 NMDA에 의한 synaptosome의 탈분극은 관찰되지 않았으나 이들은 mitochondria에서 칼슘 유입에 따른 탈분극을 초래하였다. 한편, 흥분성 아미노산은 synaptosome의 ATPase 활성도에 영향을 나타내지 않았다.

이상의 결과로부터 glutamate 또는 NMDA에 의한 synaptosome의 칼슘 흡수는 세포의 마그네슘에 각기 다른 양상을 나타내며 이들에 의한 칼슘 흡수는 포타슘을 제외한 소듐과 칼슘에 대한 세포막 투과성의 증가 그리고 이에 따른 탈분극에 연관이 있을 것으로 시사되었다.