

## Optical Resolution of Dabsyl Amino Acids in Reversed-Phase Liquid Chromatography

Sun Haing Lee\*, Tae Sub Oh, and Young Cheal Lee

Department of Chemistry, Kyungpook National University, Taegu 702-701. Received May 11, 1990

The dabsylation of amino acids has been applied to resolve their optical isomers with the use of chiral mobile phase in high performance liquid chromatography. The dabsyl amino acids were successfully separated on reversed phase column(C<sub>18</sub>) by adding a chiral L-benzylproline-Cu(II) chelate to the mobile phase. The separation selectivity of the dabsyl amino acid enantiomers was not less than that of dansyl amino acids. The retention order of the dabsyl amino acid enantiomers was as those of the dansyl amino acid enantiomers except dabsyl threonine.

The optical selectivity of the dabsyl amino acids increase with pH of the mobile phase and concentration of the chelate, but slightly decreases with concentration of buffer and organic solvent composition. However serine, methionine, valine, and leucine showed a slight decrease in the optical selectivity with increase in pH. The retention times of the dabsyl amino acids decreases with increasing pH and acetonitrile concentration but increases with the concentration of the chiral chelate added. The mechanism of the optical resolution is based on a stereospecific interaction including a intramolecular hydrophobic effect and SN-2 reactivity of the ligand exchange chromatography.

It is advantageous to detect absorption at 436 nm, which is less interferent than the other detection systems. The derivatized dabsyl amino acids are stable for a month.

### Introduction

Separation of optical isomers of amino acids is very important for synthesis of peptides and for the determination of the chemical structure<sup>1</sup>. Optical isomers of amino acids can be resolved by conversion into diastereomers. In order to carry out such a separation, an optically active agent must be present as part of either the mobile phase or stationary phase. This first attempt was carried out successfully in gas chromatography by Gil-Av and co-workers<sup>2</sup>. But recently, the separation of optical isomers of amino acids by using reversed phase high performance liquid chromatography has been of great interest<sup>3-7</sup>.

There are two different methods for the optical resolution of amino acids. The one involves the use of chiral stationary phase (CSP)<sup>8-10</sup>. The other involves the use of chiral mobile phase additives in conjugation with achiral stationary phase<sup>11,18</sup>. The previous method(CSP) is particularly applicable to the separation of enantiomers when recovery of the purified isomer is desired. Davankov and his groups<sup>9-10</sup> used the stationary phase of L-proline attached to the resin for the separation of enantiomers of amino acids. The latter method(CMPA) has advantage capable of using less expensive conventional packings. It is easier to apply this separation system to the high performance liquid chromatography. So it has been used for the optical separation of free amino acids or amino acid derivatives<sup>11-13</sup>. Karger *et al.*<sup>16,17</sup> reported the use of L-2-alkyl-4-octyldiethylenetriamine metal complexes in aqueous mobile phases for the separation of optically active dansylated amino acids(DNS-AA). Hara and Gil-Av(24) successfully separated free amino acids by use of a proline-Cu(II) eluent on an ion exchange column.

In this paper, we tried the optical resolution of dabsyl amino acid (DABS-AA) on a C<sub>18</sub> column by adding the chiral chelate of Cu(II)-N-benzyl-L-proline to the mobile phase. The resolution of optical isomers of dabsyl amino acids is compared with that of dansylated amino acids. The different elution behaviors between DNS-AA and DABS-AA are illu-


strated based on the ligand exchange reaction and hydrophobic interaction which show the stereoselectivity during the chromatographic elution.

### Experiments

**Instrument.** The liquid chromatograph from Waters Associates consisting of Series 440 absorbance detector, Model 6000 A solvent delivery system, Model U6K LC injector and Model 730 Data Module was used for this work. The UV detector was set at 436 nm. The mobile phase flow rate was 1.0 ml/min at a pressure of about 1500 psi. The column was  $\mu$ -Versapak C<sub>18</sub>(30 cm x 4.1 mm i.d.). The pH of the mobile phase was adjusted by a pH meter(Model 292) of the Fisher Scientific Company.

**Reagents.** Dabsylation was done through the precolumn derivatization of free amino acids to enhance the detectability and to see the optical separation. Free amino acids such as D, L-valine(Val), D,L-asparagine(Asn), and D,L-tyrosine(Tyr) were obtained from Sigma Chemical Co., (St. Louis, Mo, U.S.A.). D, L-leucine(Leu), D,L-phenylalanine(Phe), D,L-threonine(Thr), and D,L-methionine(Met) was obtained from Yoneyama(Osaka, Japan). Dabsyl chloride(DEBS-Cl) was obtained from Dojido Laboratories (Kumamoto, Japan). The dry amino acids were dabsylated according to the procedure described below<sup>25,26</sup>. A 1 ml dabsyl chloride solution (0.65 mg/ml in acetonitrile) was mixed with 1 ml of 0.1 M sodium bicarbonate solution containing free amino acids. The reaction mixture was shaken in water bath at 70 °C for 30 minutes when the red color of DABS-Cl was changed to orange color. The resulting solutions were directly introduced to the injector of the liquid chromatograph for the separation.

The ligand, N-benzyl-L-proline(BzPro), was synthesized by treating L-proline with benzyl chloride under a basic condition<sup>27-29</sup>. The product was identified by UV, IR, <sup>1</sup>H and <sup>13</sup>C NMR spectra. The yield was 58%, N-benzyl-L-proline was used as the ligand for the copper(II) chelate. The mobile



**Figure 1.** Typical chromatogram of DABS-amino acids with chiral mobile phase addition method. The mobile phase containing 35% acetonitrile and 65%  $5 \times 10^{-3}$  M chelate solution containing  $2 \times 10^{-2}$  M  $\text{NH}_4\text{Ac}$  buffer at pH 6.5. The chelate is  $\text{Cu(II)-(BzPro)}_2$ .

phase containing *N*-benzyl-*L*-proline and  $\text{Cu(II)}$  with 2:1 molar ratio was used for the resolutions of optical isomers of DABS-amino acids, and adjusting the pH to the desired value with acetic acid or sodium hydroxide solution. Mobile phase were degassed and purified by a solvent clarification kit just before use.

## Results and Discussion

The separation of the optical isomers of derivatized amino acids for all these compounds is of interest because the elution orders between *D* and *L* derivatized amino acids are different from those obtained in the use of different ligands such as *L*-proline, *L*-hydroxyproline, *N*-benzyl-*L*-proline, *N*-benzyl-*L*-hydroxyproline, and *N*-xylenylproline<sup>20,21,29-31</sup>. However, the elution orders of the dabsylated amino acids were the same as three of the dansylated amino acids except DABS-threonine as shown in Figure 1. On the other hand, the elution order for DABS-threonine enantiomers were reversed compared with DNS-threonine. This behavior in-

dicates that the relatively long chain of dabsyl group affects differently for the resolution of DABS-amino acids relative to the short chain of naphthyl group in DNS-amino acids. The retention behaviors for separation of the DABS-amino acids can be illustrated on the stereospecific interaction of the ligand exchange reactions. The capacity factors of DABS-amino acids and the separation selectivity of their enantiomers varied with the pH of the mobile phase as shown in Table 1. The capacity factors of DABS-amino acid enantiomers decreased with pH of the mobile phase. This behavior was the same as that of DNS-amino acids. The decrease of the capacity factor on pH indicates that the binary complexes are more stable than the ternary with increase in pH due to the greater deprotonation of the ligand and thus the ligand exchange reaction with DABS-amino acids under these conditions seems much more difficult. We believe that the optical selectivity increases with increasing pH because of difference in stereospecificity coming from the ligand exchange reaction. The optical resolution of the DABS-amino acids is also dependent on the concentration of the copper(II)-*N*-benzyl-*L*-proline chelate added as shown in Table 2. The capacity factor and selectivity of the optical isomers of DABS-amino acids increased as the concentration of the chiral chelate additive increased. The increase of capacity factors with increasing the chelate concentration indicates that the ternary complexes retain longer than the corresponding binary complex. The increase of the optical selectivity with increase in the concentration of the ternary chelate indicates that the hydrophobic interaction of the ternary complexes with the stationary phase shows enantioselectivity.

The effect of concentration of acetonitrile on the retention behaviors is shown in Table 3. It shows that the lower the concentration of acetonitrile, the greater is the retention of the isomers as expected in reversed phase high performance liquid chromatography. The selectivity between the dabsyl enantiomers with increasing acetonitrile concentration was little affected unlike the dansyl amino acids<sup>29</sup>. Therefore, it seems that there are some difference in the conformation of

**Table 1.** Capacity Ratio( $k'$ ) and Selectivity( $\alpha$ ) as a Function of pH

DABS-AA		pH 6.0		pH 6.5		pH 7.0		pH 7.5	
		$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$
Ser	D	1.03	1.08	0.89	1.06	0.96	1.05	1.10	1.04
	L	1.11		0.94		1.01		1.14	
Asn	D	1.00	0.98	0.92	0.93	1.03	0.89	1.29	0.84
	L	0.98		0.86		0.92		1.09	
Thr	D	1.58	0.92	1.31	0.85	1.30	0.85	1.50	0.77
	L	1.45		1.11		1.10		1.15	
Tyr	D	2.30	0.83	1.77	0.80	1.68	0.77	1.78	0.71
	L	1.90		1.42		1.29		1.26	
Met	D	3.38	1.07	2.61	1.04	2.53	1.06	2.77	1.00
	L	3.63		2.72		2.67		2.78	
Val	D	4.38	1.14	3.06	1.11	2.83	1.11	2.77	1.07
	L	5.01		3.39		3.15		2.96	
Phe	D	6.52	0.82	4.78	0.78	4.54	0.74	4.77	0.69
	L	5.33		3.71		3.36		3.31	
Leu	D	6.87	1.17	4.74	1.15	4.37	1.18	4.31	1.14
	L	8.06		5.44		5.15		4.92	

35% Acetonitrile and 65%  $\text{Cu(II)}$  chelate solution containing  $5 \times 10^{-3}$  M  $\text{Cu(II)-(BzPro)}_2$  is mobile phase. The ammonium acetate buffer concentration was  $2.5 \times 10^{-2}$  M. The flow rate is 1.0 ml/min.

**Table 2.** Capacity Ratio( $k'$ ) and Selectivity( $\alpha$ ) as a Function of Complex Concentration

DABS-AA		$1.0 \times 10^{-3}$ M		$5.0 \times 10^{-3}$ M		$10 \times 10^{-3}$ M	
		$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$
Ser	D	0.95	1.02	0.88	1.07	0.14	1.25
	L	0.97		0.94		1.43	
Asn	D	1.00	0.94	0.90	0.94	1.12	0.94
	L	0.94		0.85		1.05	
Thr	D	1.13	0.89	1.32	0.84	1.76	0.86
	L	1.01		1.11		1.52	
Tyr	D	1.65	0.81	1.80	0.81	2.62	0.76
	L	1.34		1.45		2.00	
Met	D	2.41	1.03	2.66	1.04	3.80	1.07
	L	2.49		2.76		4.05	
Val	D	2.19	1.02	3.12	1.10	5.29	1.25
	L	2.24		3.43		6.61	
Phe	D	4.12	0.81	4.94	0.78	7.77	0.74
	L	3.34		3.84		5.72	
Leu	D	3.54	1.04	4.91	1.14	8.26	1.29
	L	3.71		5.59		10.65	

35% Acetonitrile and 65% Cu(II) chelate solution containing  $5 \times 10^{-3}$  M Cu(II)-(BzPro)<sub>2</sub> is the mobile phase. The ammonium acetate buffer concentration was  $2.5 \times 10^{-2}$  M at pH 6.5. The flow rate is 1.0 ml/min.

the DABS-amino acids compared with those of DNS-amino acids due to the long dabsyl group.

Separation of the optical isomers of DABS-amino acids is also slightly dependent on the concentration of buffer in the mobile phase as shown in Table 4. The capacity factors of DABS-amino acids and their optical selectivity slightly decreased as the concentration of ammonium acetate in the mobile phase increased. This behavior can be explained that acetate ion as well as benzyl proline as a ligand is competitive to the binary chelate. So the decrease in the concentration of

**Table 4.** Capacity Ratio( $k'$ ) and Selectivity( $\alpha$ ) as a Function of Buffer Concentration.

DABS-AA		$1.0 \times 10^{-2}$ M		$2.0 \times 10^{-2}$ M		$3.0 \times 10^{-2}$ M	
		$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$
Ser	D	1.00	1.05	0.98	1.08	0.97	1.09
	L	1.05		1.06		1.06	
Asn	D	1.02	0.93	1.02	0.92	1.04	0.90
	L	0.95		0.94		0.94	
Thr	D	1.58	0.84	1.51	0.88	1.45	0.83
	L	1.33		1.33		1.20	
Tyr	D	2.23	0.68	2.21	0.76	2.00	0.78
	L	1.52		1.67		1.56	
Met	D	2.48	1.41	3.18	1.08	2.93	1.05
	L	3.50		3.43		3.07	
Val	D	4.10	1.16	3.99	1.15	3.39	1.09
	L	4.77		4.58		3.69	
Phe	D	6.08	0.72	6.23	0.75	5.59	0.75
	L	4.38		4.67		4.18	
Leu	D	6.32	1.25	6.27	1.20	4.99	1.22
	L	7.87		7.51		6.08	

35% Acetonitrile and 65% Cu(II) chelate solution containing Cu(II)-(BzPro)<sub>2</sub> at pH 6.5 is the mobile phase. The flow rate is 1.0 ml/min.

the ternary complexes resulting from increase in acetate ion decreases the retention and the optical selectivity as well.

Several models<sup>24,30-36</sup> which illustrate the separation mechanism of the optical isomers have been suggested. The separation of the optical isomers of the DABS-amino acids can be explained with a *cis-trans* reaction mechanism for the optical separation of DABS-amino acids. It can be suggested that two molecules of N-benzyl-L-proline be bounded to Cu(II) to form five membered ring<sup>32</sup> and the copper chelates resulted are well known as square planar<sup>37</sup> to have a *trans* or *cis* configuration. The *trans* configuration of

**Table 3.** Capacity Ratio( $k'$ ) and Selectivity( $\alpha$ ) as a Function of Acetonitrile Concentration


DABS-AA		32.5%		35.5%		37.5%		40.0%	
		$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$	$k'$	$\alpha$
Ser	D	1.02	1.03	0.91	1.14	0.67	1.07	0.53	1.08
	L	1.05		1.04		0.72		0.57	
Asn	D	1.08	0.90	0.95	0.94	0.72	0.92	0.56	0.96
	L	0.97		0.89		0.66		0.54	
Thr	D	1.56	0.83	1.26	0.87	0.95	0.89	0.73	0.89
	L	1.29		1.09		0.85		0.65	
Tyr	D	2.13	0.82	1.52	0.99	1.22	0.82	0.97	0.76
	L	1.73		1.51		1.00		0.74	
Met	D	3.22	1.02	2.80	1.18	1.98	1.06	1.46	1.07
	L	3.29		3.31		2.09		1.56	
Val	D	3.67	1.10	3.14	1.23	2.32	1.13	1.71	1.13
	L	5.98		5.35		3.58		2.54	
Phe	D	5.98	0.77	5.35	0.80	3.58	0.77	2.54	0.78
	L	4.59		4.26		2.74		1.98	
Leu	D	5.84	1.12	4.22	1.44	3.55	1.16	2.56	1.18
	L	6.56		6.09		4.13		3.03	

The added chelate solution is  $5 \times 10^{-3}$  M Cu(II)-(BzPro)<sub>2</sub>. The ammonium acetate buffer concentration was  $2.5 \times 10^{-2}$  M at pH 7.0. The flow rate is 1.0 ml/min.

DABS-amino acids and copper binary complexes is believed to have higher enantioselectivity than the *cis* form due to the symmetric configuration. So it can be assumed that the *trans* forms of the binary complexes are prevalent in the mobile phase.

The side chains in the dabsyl amino acids seem to have an intramolecular hydrophobic or hydrophilic interactions with the dabsyl group to form a dominant conformation<sup>29-31</sup>. The intramolecular interactions of the dabsyl amino acids seem to be more feasible than those of the dansyl amino acids due to the relatively long dabsyl group.

It has been believed that the ligand exchange reaction occurs as follows.



The equilibrium constants of D-isomers are not identical with those of L-isomers because of the difference in enantioselectivity for the ligand exchange reactions in the mobile phase. The retention behaviors of the DABS-amino acids can be explained by the proposed mechanism reported from the previous results<sup>21,29</sup>.

It can be suggested that the optical isomers of DABS-amino acids have an exchange reaction with the copper(II)-(BzPro)<sub>2</sub> chelate by a SN-2 reaction. The ligand exchange of the binary complexes (Cu(II)-(BzPro)<sub>2</sub>) with D- or L-DABS amino acids have an enantioselectivity to exhibit the separation of the enantiomers. L-DABS-amino acids are able to attack the Cu(II)-(BzPro)<sub>2</sub> chelate to produce the ternary complexes consisting of both *cis* and *trans* configurations without the steric hindrance, while D-DABS-amino acids are able to attack the Cu(II)-(BzPro)<sub>2</sub> chelate to form the ternary complexes of *cis* configuration but hardly to form those of *trans* configuration due to the steric hindrance of the five membered ring of L-benzyl proline with the dabsyl group of the DABS-amino acids. Therefore, the L-DABS amino acids are more retained than D-DABS amino acids because the ternary complexes of L-DABS amino acids retain much more than those of D-DABS amino acids.

However, the elution orders between D- and L-DABS-amino acids in case of DABS-Asn, DABS-Tyr, an DABS-Phe were reversed as known from the results of the dansyl amino acids<sup>29-31</sup>. It seems to have a greater hydrophobic interaction of the *cis* products resulted from L-forms because the prolyl group of the chelate has a same plane with the dabsyl group and side chain of the DABS-amino acids. It can be speculated that the bulky groups of the three rings on the same plane have a strong hydrophobic interaction with the reversed stationary phase.

**Acknowledgement.** The authors gratefully acknowledge the Korea Science and Engineering Foundation for support (1987) of this work.

## References

1. N. B. Levine and V. Nasinow, *J. Chromatogr.*, **286**, 207 (1984).

2. R. Charles, U. Beitler, B. Feibush, and E. Gil-Av., *J. Chromatogr.*, **112**, 121 (1975).
3. R. V. Snyder, R. J. Angelici, and R.B. Meck, *J. Amer. Chem. Soc.*, **94**, 2660 (1972).
4. B. Lefebvre, R. Audebert, and C. Quivoron, *J. Liq. Chromatogr.*, **1**, 761 (1978).
5. R. J. Baczuk, G. K. Landram, and R. J. Dubois, *J. Chromatogr.*, **60**, 351 (1971).
6. W. H. Pirkle and D. W. House, *J. Org. Chem.*, **44**, 1957 (1979).
7. L. R. Sousa, G. D. Sogah, D. H. Hoffman, and D. J. Crem, *J. Amer. Chem. Soc.*, **100**, 4569 (1978).
8. Y. Yuki, K. Saigo, K. Tachihana, and M. Hasegawa, *Chem. Lett.*, 1347 (1986).
9. V. A. Davankov and S. V. Roghozhin, *J. Chromatogr.*, **61**, 280 (1971).
10. V. A. Davankov and A. V. Semechkin, *J. Chromatogr.*, **141**, 313 (1977).
11. S. Lam and G. Malikin, *J. Chromatogr.*, **368**, 413 (1986).
12. S. Lam, F. Chow, and A. Karmen, *J. Chromatogr.*, **199**, 295 (1980).
13. T. Takeuchi, H. Asai, Y. Hashimoto, K. Watanabe, and D. Ishii, *J. Chromatogr.*, **331**, 99 (1985).
14. P. E. Hara and E. Gil-Av., *J. Chromatogr.*, **214**, 1226 (1979).
15. J. N. Lepage, W. Linder, G. Davaies, D. E. Seitz, and B. L. Karger, *Anal. Chem.*, **51**, 433 (1979).
16. Y. Tapuhi, N. Miller, and B. L. Karger, *J. Chromatogr.*, **205**, 325 (1981).
17. M. H. Engel and S. A. Mocko, *Anal. Chem.*, **56**, 2598 (1984).
18. S. Lam and A. Karmen, *J. Chromatogr.*, **255**, 41 (1983).
19. S. H. Lee, T. S. Oh, and K. S. Park, *J. Korean Chem. Soc.*, **30**, 216 (1986).
20. S. H. Lee, J. W. Ryu, and K. S. Park, *Bull. Korean Chem. Soc.*, **7**, 45 (1986).
21. C. Gilon, R. Leshem, and E. Grushika, *J. Chromatogr.*, **205**, 365 (1981).
22. N. Qi and H. Kitohara, *J. Chromatogr.*, **295**, 198 (1984).
23. E. Gil-Av., A. Tishbee, and P. E. Hara, *J. Amer. Chem. Soc.*, **102**, 5115 (1980).
24. J. K. Lin, and J. K. Chang, *Anal. Chem.*, **47**(9), 1634 (1975).
25. R. Knecht and J. Y. Chang, *Anal. Chem.*, **58**, 2375 (1986).
26. J. Jozefonvicz, D. Muller, and M. A. Petit, *J.C.S. Dalton*, 76 (1980).
27. J. March, "Advanced Organic Chemistry", 2nd Ed., 377(1977).
28. S. H. Lee, T. S. Oh, and B. E. Kim, *Bull. Korean Chem. Soc.*, **9**, 345 (1988).
29. S. H. Lee, T. S. Oh, and S. H. Bak, *Bull. Korean Chem. Soc.*, **10**, 491 (1989).
30. S. H. Lee, T. S. Oh, and B. J. Park, *J. Korean Chem. Soc.*, **34**, 76 (1990).
31. C. Gilon, R. Leshem, and E. Grushika, *Anal. Chem.*, **52**, 1206 (1980).
32. A. A. Kurganov and V. A. Davankov, *J. Chromatogr.*, **218**, 559 (1981).
33. C. Gilon, R. Leshem, and E. Grushika, *J. Chromatogr.*,

- 203, 365 (1981).  
 35. V. Beitler and B. Feibush, *J. Chromatogr.*, **123**, 149 (1976).  
 36. W. H. Pirkle and D. Sikkenga, *J. Chromatogr.*, **123**, 149 (1976).  
 37. V. A. Davankov and P. R. Mitchell, *J. C. S. Dalton*, 1012 (1972).

## Synthesis and Electrical Conductivities of Poly(1,4-phenylenevinylene-co-2,3,5,6-tetramethyl-1,4-phenylenevinylene)s

Jung-II Jin\*, Heung-Joong Kang, and Hong-Ku Shim†

Department of Chemistry, Korea University, Seoul 136-701

†Department of Chemistry, Korea Advanced Institute of Science and Technology, Taejon 305-701

Received May 15, 1990

A series of copolymers of poly(1,4-phenylenevinylene-co-2,3,5,6-tetramethyl-1,4-phenylenevinylene), poly(PV-co-TMPV), were prepared in film forms from the precursor polymer films. The sulfonium salt precursor polymers were synthesized by copolymerization of the mixtures of the respective bis(sulfonium salt) monomers. All of the copolymer films could be doped with  $\text{FeCl}_3$  to have high electrical conductivities and they showed good air stability. The maximum conductivity of the  $\text{FeCl}_3$ -doped films ranged  $10^{-3}$  to  $10^2 \text{Scm}^{-1}$  depending on the composition of the copolymer films. However, these copolymer films could not be doped with iodine. The coplanarity of PV and TMPV units in the main chain appears to be affected by steric effect of the methyl groups in the TMPV units.

### Introduction

The synthesis of poly(1,4-phenylenevinylene), PPV, and its derivatives and copolymers through the water-soluble precursor route<sup>1-7</sup> is one of the most promising methods to obtain the polymers with extended  $\pi$ -conjugated structures. The PPV derivatives are attracting much interest as new conductive materials and materials for non-linear optics<sup>8-11</sup> because they can be obtained as dense, tough and flexible films and show superior chemical stability. The precursor route has been applied to the preparations of poly(2,5-dimethoxy-1,4-phenylenevinylene)(PDMPV)<sup>4</sup>, poly(2,5-thienylenevinylene)(RTV)<sup>5</sup> and their copolymers.<sup>6</sup>

The PPV films obtained in this manner can be easily doped with strong oxidizing agents such as  $\text{AsF}_5$  to produce highly conducting materials, but they can not be effectively doped with  $\text{I}_2$ .<sup>2</sup> Doping PPV films with  $\text{AsF}_5$  results in conductivities of  $10\text{--}40 \text{Scm}^{-1}$  for unstretched samples and  $500\text{--}3000 \text{Scm}^{-1}$  for uniaxially stretched ones.<sup>2,3,12</sup> In contrast, the PPV derivatives such as PDMPV and its copolymers show easy dopability to  $\text{I}_2$  leading to high conductivities. For example, the  $\text{I}_2$ -doped PDMPV exhibits an electrical conductivity in the order of  $10^2 \text{Scm}^{-1}$  at room temperature. The high conductivities observed for doped PDMPV can be attributed to the electron-donating capacity of the methoxy groups on the phenylene group. They lower the oxidation potential of the conjugated polymers.<sup>13</sup>

In the preparation of these polymer films, water-soluble sulfonium salt precursor polymers of high molecular weight are first prepared and cast into films. They are then subjected to thermal elimination to the final polyconjugated polymers. If desired, the precursor films can be uniaxially

drawn before or during the thermal elimination reaction. Oriented films show a significantly enhanced conductivity on doping along stretched direction.<sup>14,15,20</sup>

Recently, we prepared poly(2-methoxy-5-methylthio-1,4-phenylenevinylene)(PMMPV) and its copolymers,<sup>14</sup> and poly(2-*n*-butoxy-5-methoxy-1,4-phenylenevinylene)(PBMPV)<sup>15</sup> via the corresponding water-soluble precursor polymers. These polyconjugated polymers could be readily doped with oxidizing dopants such as  $\text{I}_2$ ,  $\text{FeCl}_3$  and  $\text{Fe}(\text{ClO}_4)_3$  to produce materials having a wide range of conductivities depending on the nature of substituents and dopants. In the case of PBMPV,  $\text{I}_2$ -doped films exhibited conductivities as high as  $590 \text{Scm}^{-1}$  and  $\text{FeCl}_3$ -doped ones as high as  $2160 \text{Scm}^{-1}$ . The films could be stretched up to a draw ratio of 4-8 before the final elimination.

In this paper, we describe the synthesis of a series of poly(1,4-phenylenevinylene-co-2,3,5,6-tetramethyl-1,4-phenylenevinylene)s, poly(PV-co-TMPV), using different mole ratios of the two respective sulfonium salt monomers in polymerization. The copolymers in film forms were doped with  $\text{FeCl}_3$  and their electrical conductivities were measured. It was our purpose to learn more about the substituent effect on the electrical properties of PPV derivatives. The synthetic route and the structure of poly(PV-co-TMPV)s prepared and characterized in this investigation are shown below;

