

bons), 40.8(CH) minor isomer, 165.4(CO₂), 89.4, 87.3, 73.0, 62.4, 41.4, 33.3(ring carbons), 41.4(CH), 64.5(OCH₂), 13.8(OCH₂CH₃), 21.2(Me) ppm. EI-MS, *m/z*, 290(M⁺-2CO), 260(M⁺-2CO-NO), 245(M⁺-2CO-NO-Me), 234(M⁺-2CO-NO-Me-CN). 5d. major isomer, ¹H NMR(CDCl₃) δ 5.9(m, 4H), 4.20(q, 7.2 Hz, OCH₂), 3.69(d, 2.9 Hz, CH), 3.31(m, 2H), 2.35(m, 2H), 2.26(s, C(O)CH₃), 1.27(t, 7.2 Hz, CH₃), 0.85(d, 7.1 Hz, CH₃) ppm; minor, ¹H NMR(CDCl₃) δ 5.9(m, 4H), 4.19(q, 7.1 Hz, OCH₂), 3.75(d, 2.2 Hz, CH), 3.3(m, 2H), 2.35(m, 2H), 2.27(s, C(O)CH₃), 1.28(t, 7.1 Hz, CH₃), 1.28(t, 7.1 Hz, CH₃), 0.81(d, 7.1 Hz, CH₃) ppm; 5f(R=Me, Nu=SCH₂CH₂OH), ¹H NMR(CDCl₃) δ 5.60~6.03(m, 4H), 2.90(m, 1H), 2.73(m, 1H), 3.71(br, OH), 3.41(t, 3.5 Hz, OCH₂), 2.66(t, 2.2 Hz, SCH₂), 1.27(d, 9.1 Hz, Me). [{MeC₆H₄(CH(CN)CO₂Et)}Mn(CO)NO(P(OPh)₃)] yield 81.6%, IR ν_{CO} 1967 cm⁻¹, ν_{NO} 1735 cm⁻¹, ν_{CO₂} 1720 cm⁻¹, ¹³C NMR(CDCl₃) δ 229.3(CO), 150.1, 128.8, 124.1, 120.1(P(OPh)₃), 116.4(CN), major isomer, 165.1(CO₂), 88.2, 84.4, 71.9, 60.2, 40.2, 32.9 (ring carbons), minor isomer, 164.6(CO₂), 88.0, 84.0, 71.7, 60.4, 40.6, 32.9(ring carbons), 61.7(OCH₂), 42.5, 42.1(CH), 20.5(Me), 12.9(OCH₂CH₃).

6. Y. K. Chung and D. A. Sweigart, *J. Organomet. Chem.* **308**, 223 (1986).
7. L. A. P. Kane-Maguire, E. D. Honig and D. A. Sweigart, *Chem. Rev.*, **84**, 525 (1984).
8. W. A. Henderson and S. A. Buckler, *J. Am. Chem. Soc.*, **82**, 5794 (1960); W. E. McEwen, W. I. Shian, Y.-I. Yeh, D. N. Schulz, R. U. Pagilagan, J. B. Levy, C. Symmes, G. O. Nelson and I. Granth, *J. Am. Chem. Soc.*, **97**, 1782 (1975) and references therein.
9. R. G. Pearson, H. Sobel and J. Songstad, *J. Am. Chem. Soc.*, **90**, 319 (1968).
10. A. J. Pearson, S. L. Kole and B. Chen, *J. Am. Chem. Soc.*, **105**, 448 (1983); A. J. Pearson, S. L. Kole and T. Ray, *J. Am. Chem. Soc.*, **106**, 6060 (1984).
11. R. D. Pike, W. J. Ryan, G. B. Carpenter and D. A. Sweigart, *J. Am. Chem. Soc.*, **111**, 8535 (1989).

A Voltammetric Procedure for Determinations of Four Main Constituents in Ancient Coins

Myung-Zoon Czae*, Suw Yong Ly, and Chul Lee

Department of Chemistry, Hanyang University,
Seoul 133-791

Received August 27, 1990

Quite often analytical chemists are asked for to analyze several analytes in the same sample or in a set (large numbers) of samples, in many cases, for the exploratory purposes.^{1,2} In this laboratory a rapid simple method is required which can be applied to the multicomponent analysis of coin samples for the purposes of pattern recognition. Samples for




Figure 1. Typical DPAS voltammograms in 0.05 M, pH 8.0 sodium borate solution containing [Sn(IV)]= 2.5×10^{-6} M and [Pb(II)]= 2.7×10^{-7} M, showing the dependence of peak resolution on deposition potentials: A, -1.6V; B, -1.4V; C, -1.2V vs Ag/AgCl. Deposition time, 4 min. Scan rate, 10 mV/sec. Modulation amplitude, 5 mV. Pulse repetition time, 1 sec.

analysis were ancient coins (1423-1883 A.D.) having chemical compositions,³ 60-92% copper, 3-26% lead, 0.2-29% zinc, and 0.2-6% tin as the main constituents.

Recently there is a growing appreciation of the power of the pulse voltammetry for it provides multielement capability. It can offer the advantage of simultaneous multi-element determinations (four to six metals) at the lowest costs compared to other competing multianalyte techniques if a suitable experimental procedure is provided.⁴

The one-element-at-a-time nature of the most existing methods^{5,6} and the costs of ICP, X-ray fluorescence analysis⁷ and the limited availability of neutron activation analysis and their lacks of sensitivity (accordingly restriction to) certain metals suggest that the search for an electrochemical multimetal analysis scheme for the alloy samples is timely.

Ideally, a procedure employing only a single technique throughout the entire run is desirable, saying, that can detect the four main elements simultaneously and present their signals to the strip-chart recorder at a time with satisfactory resolutions. However it was not achievable because of the presence of stannic tin, which has peculiar behaviors and shows a great overlapping interference with lead.⁸ In this work, as the second best, the advantages of differential pulse polarography (DPP) for copper, lead, and zinc and anodic stripping voltammetry (ASV) for tin are combined.

In most cases (for both DPP and ASV), good chemistry is the proper solution to the resolution problem. One approach is to make choice of a supporting electrolyte medium. Preli-




Figure 2. Differential pulse polarograms of a coin sample (No. 49) in 0.05 M sodium borate supporting electrolyte at pH 2.5. Later estimated concentrations: $[Cu(II)] = 0.116$ mM, $[Pb(II)] = 12.6$ μ M, $[Zn(II)] = 20.9$ μ M, and $[Sn(IV)] = 2.64$ μ M. Conditions: scan rate 5 mV/sec, modulation amplitude 25 mV, and pulse repetition time 1 sec.

primary experiments were primarily intended to find out a such medium that (1) can be applicable to both techniques consistently without exchanging and with a minimum of sample solution manipulation, and (2) allows satisfactory separations, between lead and tin in ASV measurement, and in preceded (or subsequent) DPP measurements for the rest three metals. Although a citrate buffer has been shown⁸ to be a medium well suited to the purpose, it was excluded because of the complexity in sample solution manipulation (matrix exchange in the stripping step).

All voltammetric measurements were carried out with an 174A of EG&G PAR as described previously.⁹ Stock solutions of metal ions (1.000 g/l) and sample solutions (0.06–0.13 g/50 ml) were prepared by dissolving drilled chips of the pure metals and sample coins respectively in acids as directed.⁵ Temperature of the solution was laboratory ambient ($24 \pm 2^\circ\text{C}$).

A typical stripping voltammogram of a synthetic solution showing a good resolution of tin and lead peaks in sodium borate medium is presented in Figure 1. At low deposition potential (C), the two peaks are well resolved (more than 200 mV) with a lower sensitivity of tin which would be ascribed to a low efficiency of the reduction step, as in citrate buffer.⁸ In practice, however, more negative potential (B or A) was found to be more preferable owing to another decrease in sensitivity due to the formation of Cu-Sn intermetallic compound, the effect of which is alleviated by using the generalized standard addition method.¹⁰ Sodium borate medium (0.05 M, pH 2.0–2.5 this time) offers also proper DP polarograms with good separations between analytical peaks and especially copper peak from the sloping portion of the base line (the latter advantage was lost in the case of strongly acidic solution, 0.1 M HCl), demonstrated as in Figure 2.

As the result, the following procedure is proposed: pipet 10 ml of 0.05 M sodium borate solution and a 50 μ l aliquot of sample solution into the electrochemical cell. Adjust pH between 2.0–2.5 with dilute nitric acid (or sodium hydroxide). After de-oxygenation for 10 min, obtain DP polarograms selecting the following working parameters: initial potential, 0.2V; scan rate, 5 mV/sec; scan range, 1.5V; drop time, 1 sec; and modulation amplitude, 25 mV. Subsequently, just adjust pH again to 7.6–8.0 with NaOH for the ASV run.

After de-oxygenation for 2 min, start the stirrer and deposition step applying -1.4V for 4 min. After 30 sec resting period, anodic scan is started at the following conditions: scan rate 10 mV/sec; drop time 1 sec, modulation amplitude 5 mV.

For the quantitations, general standard addition method (GSAM) is recommended in ASV, whereas the external standards (using a calibration curve) are used in the DPP. Two standard additions of tin are made first and then two additions of copper are done keeping the coverage of the response surface in mind. Calculate the results by the total difference calculations—*g.s.a.m.* If the simple standard addition method is to be used in the DPP (as in the small samples case), it may be followed the ASV. Work is in progress in this laboratory to elaborate the method and for a deeper rationale behind it.

The method bias of the proposed procedure was evaluated by analyzing 10 samples and comparing the measurements by neutron activation analysis as the reference method. The *t*-values (at the 0.05 significant level) were 1.88, 1.60, 1.98 and 2.01 for Cu, Pb, Zn and Sn respectively showing no difference between the two methods.


In conclusions, a simple, rapid electrochemical method has been developed for the multianalyte (four) analysis of ancient coins. The use of sodium borate as a supporting electrolyte permits the simultaneous determination of copper, lead and zinc by DPP and subsequent ASV determination of tin consecutively without medium exchange. Good agreement with neutron activation analysis data confirms the accuracy, efficiency and the applicability of the proposed procedure to multicomponent analysis of copper-base alloy samples with a minimum of sample solution manipulation.

Acknowledgement. This work was generously supported by the Ministry of Education as a Basic Science Research Institute Program—1989.

References

- (a) M. A. Sharaf, D. L. Illman and B. R. Kowalski, "Chemometrics" John Wiley & Sons, p. 179 1986; (b) D. L. Massart, B. G. M. Vandeginste, S.N. Deming, Y. Michotte and L. Kaufman, "Chemometrics: a textbook," Elsevier, Chapt. 20–23 1988.
- (a) B. R. Kowalski, T. F. Schatzki, and F. H. Stross, *Anal. Chem.*, **44**, 217 (1972); (b) C. Jochum, P. Jochum, and B. R. Kowalski, *Anal. Chem.*, **55**, 85 (1981).
- Hyung Tae Kang, Ph.D. Thesis, Hanyang University, 1989.
- Joseph Wang, "Stripping Analysis Principles, Instrumentation and Applications," VCH Publishers, p. 2 1985.
- ASTM Designation: E 54–80, and E 62–76. in "Annual Book of ASTM Standards," Vol. 03. 05, p. 234, 1986.
- R. S. Young, "Encyclopedia of Industrial Chemical Analysis," Vol. 10, p. 598. F. D. Snell and L. S. Ettre, Eds., Interscience, New York, 1979.
- D. P. Stevenson, F. H. Stross and R. F. Heizer, *Archaeometry*, **13**, 17 (1971).
- E. Desimoni, F. Palmisano and L. Sabbatini, *Anal. Chem.*, **52**, 1889 (1989); S. Mannino, *Analyst*, **109**, 905 (1984).

9. T. Hong and M.-Z. Czae, *Bull. Korean Chem. Soc.*, **9**, 77 (1988).
 10. R. W. Gerlach and B. R. Kowalski, *Anal. Chim. Acta*, **134** (1982).



Scheme 1.

A New Route to N-Substituted Aminomethylphosphonates

Hyun-Joon Ha*, Gong-Sil Nam, and Kyong Pae Park

Division of Chemistry,

Korea Institute of Science and Technology, Seoul 136-791

Received August 27, 1990

Some of N-substituted aminomethylphosphonic acids and their esters are known to have herbicidal and fire-proofing applications.¹ Synthesis of their acids has been developed in a wide variety of methods.^{2,3} However, synthetic methodology of their esters is relatively rare. Only two methods were reported for the preparation of their esters: (1) substitution of chloromethylphosphonates³ or phosphonomethyltriflates⁴ with primary amines (2) rearrangement of P-(halomethyl)-N-phenylphosphonoamides.⁵ These are not generally applicable for diverse N-substituted aminomethylphosphonates.

We recently described that condensation of N-substituted N-methoxymethylamines (1) with trialkylphosphites affords N-substituted aminomethylphosphonates (3,4).⁶ The reaction of N-substituted amine with paraformaldehyde and sodium methoxide in methanol yields either N-substituted N-methoxymethylamine (1) or 1,3,5-trisubstituted-hexahydro-1,3,5-triazine⁷ (2) depending on the substituent R¹.⁸ The heating of 1 under reduced pressure results in a quantitative transformation into the corresponding 2. In this communication we present a new route to the synthesis of 3 and 4 involving the reaction of 2 with trialkylphosphites in the pre-

sence of TiCl₄.⁹ (Scheme 1)

Typical reaction procedure is as follows. To a stirred solution of 1,3,5-trisubstituted-hexahydro-1,3,5-triazine (2) (6 mmol) in 100 ml of CH₂Cl₂ under nitrogen was slowly added TiCl₄ (21 mmol) at 0°C. After being stirred for 10 min, trialkylphosphite (18 mmol) was added. The resulting solution was stirred at 0°C for 1-2 hr. After being observed that all starting material was consumed on TLC the reaction mixture was poured into water. The resulting solution was neutralized with sat. NaHCO₃ solution. The reaction product was extracted with CH₂Cl₂. Organic layer was washed successively with water and brine, dried over anhydrous MgSO₄, filtered, and concentrated under reduced pressure. The reaction product was further purified by column chromatography or short-path distillation. Results are summarized in the Table 1.

This method is widely applicable to the various N-substituted aminomethylphosphonates including simple alkyl (entry 1,2,3), bulky t-butyl (entry 4), allyl (entry 5), cyclohexyl (entry 6,7), phenyl (entry 8), benzyl (entry 9) as substituents. Furthermore, a commercial herbicide N-phosphonomethylglycine¹⁰ is also accessed by this method⁹ as its corresponding trialkylesters (entry 10,11).^{11,12}

The mechanism of this reaction is not clear. But, α-methyleneamine peaks in ¹H and ¹³C NMR were observed by mixing 1,3,5-trisubstituted-hexahydro-1,3,5-triazine and TiCl₄.¹³ The observed peaks were close to those of free N-substituted methyleneamine¹⁴ generated by flash vacuum thermolysis of the corresponding N-substituted aminoacetonitrile. Moreover, the color of this mixture became deep wine-red, which indicated that formation of α-methyleneamine complex with TiCl₄. On the basis of the observations we predict that this reaction is proceeded by

Table 1. N-Substituted Aminomethylphosphonates Prepared from the Reaction of 1,3,5-Trisubstitutedhexahydro-1,3,5-Triazines with Trialkylphosphites in the Presence of Titanium Tetrachloride

Entry	R ¹	R ²	Yield ^{a,b} (%)	¹ H NMR(CDCl ₃) ^c	IR (cm ⁻¹ , P=O)
1	CH ₃ CH ₂	C ₂ H ₅	49	3.16(d, 10 Hz)	1256
2	CH ₃ (CH ₂) ₂ CH ₂	CH ₃	65	3.17(d, 12 Hz)	1249
3	CH ₃ (CH ₂) ₂ CH ₂	C ₂ H ₅	69	3.00(d, 12 Hz)	1239
4	(CH ₃) ₃ C	C ₂ H ₅	45	3.83(dd, 10 Hz, 5 Hz)	1239
5	CH ₂ CHCH ₂	C ₂ H ₅	60	2.94(d, 10 Hz)	1255
6	Cyclo-C ₆ H ₁₁	CH ₃	45	3.06(d, 10 Hz)	1255
7	Cyclo-C ₆ H ₁₁	C ₂ H ₅	48	3.10(d, 10 Hz)	1253
8	Ph	C ₂ H ₅	80	3.54(dd, 12 Hz, 6 Hz)	1212
9	PhCH ₂	C ₂ H ₅	35	3.15(d, 10 Hz)	1257
10	C ₂ H ₅ O ₂ CCH ₂	CH ₃	92	3.17(d, 12 Hz)	1239
11	C ₂ H ₅ O ₂ CCH ₂	C ₂ H ₅	65	3.05(d, 12 Hz)	1238

^aYield of isolated pure product, not optimized except entry 10. ^bSatisfactory microanalyses obtained. ^cChemical shift of two methylene protons between nitrogen and phosphorus, referenced by TMS.