


Table 2. Oxidative Cleavage of Vicinal Diols with NiO₂-Al₂O₃ in Benzene at 50°C

Diols	Products	Time, h	Yield, % ^{a,b}
<i>trans</i> -1,2-cyclohexanediol	1,6-hexanedial	5	75
1-phenyl-1,2-ethanediol	benzaldehyde + formaldehyde	5	98
1,2; 5,6-di-O-isopropylidene-D-glycer-	1,2-O-isopropylidene-D-glycer-aldehyde	14	52
3	4 + formaldehyde	10	(43)

^a The yields were determined by GC. ^b The yields in parentheses are isolated ones.



indicates that NiO₂-Al₂O₃ system oxidizes benzylic and allylic alcohols to the corresponding aldehydes or ketones in shorter reaction time and in much higher yield than either activated MnO₂ or NiO₂ alone. For example, NiO₂-Al₂O₃ oxidizes geraniol to geranial in 85% in 6 h at room temperature whereas MnO₂ oxidizes geraniol to geranial in 50% in 90 h at 50°C.¹² Efficiency of NiO₂-Al₂O₃ system was more clearly shown in the oxidation of a carbohydrate allylic alcohol **1**. Oxidation of allylic alcohol **1** to ketone **2** was sluggish with activated MnO₂ or NiO₂ alone. Other various oxidizing agents such as PCC, PDC, and KMnO₄ under phase-transfer condition were not efficient and provided complex mixtures of products in certain cases. NiO₂-Al₂O₃ system efficiently oxidized compound **1** to its ketone **2** in 90% yield.

NiO₂-Al₂O₃ system also cleaved vicinal diols as shown in Table 2. 1-Phenyl-1,2-ethanediol was converted to benzaldehyde and formaldehyde in 98% yield in benzene at 50°C. At room temperature, however, quite long reaction time was required and the yield was substantially lowered. 1,2,5,6-Di-O-isopropylidene-D-mannitol and a carbohydrate diol **3** were also cleaved to the corresponding aldehydes in reasonable yields. The ease of using NiO₂-Al₂O₃ in organic solvents for cleavage of diols indicates that this solid oxidizing agent may prove to be a useful alternative to sodium metaperiodate or lead tetraacetate.

Acknowledgement. This work was supported by a grant from Yonsei University.

References

- (a) M. Z. Barakat, M. F. Abdel-Wahab, and M. M. El-Sadr, *J. Chem. Soc.*, 4685 (1959); (b) L. Crombie and J. Crossley, *J. Chem. Soc.*, 4983 (1963); (c) I. T. Harrison, *Proc. Chem. Soc.*, 110 (1964).
- E. Santaniello and P. Ferraboschi, *Synth. Commun.*, **10**, 75 (1980).

- F. S. Guziec and F. A. Luzzio, *J. Org. Chem.*, **47**, 1787 (1982).
- K. S. Kim, Y. K. Chang, S. K. Bae, and C. S. Hahn, *Synthesis*, 866 (1984).
- K. S. Kim, S. J. Kim, Y. H. Song, and C. S. Hahn, *Synthesis*, 1017 (1987).
- K. S. Kim, S. Chung, I. H. Cho, and C. S. Hahn, *Tetrahedron Lett.*, **30**, 2559 (1989).
- M. V. George, in "Organic Syntheses by Oxidation with Metal Compounds", W. J. Mijs and C. R. H. I. De Jonge eds., Plenum Press, New York, pp. 373-422, (1986).
- K. Nakagawa, R. Konaka, and T. Nakata, *J. Org. Chem.*, **27**, 1597 (1962).
- NiO₂ was prepared as described in reference 8.
- Al₂O₃ (W-200-B, Activity grade Super I) was purchased from ICN Pharmaceuticals, Inc. and used without further activation.
- The solid mixture was prepared by gentle grinding a mixture of NiO₂ and Al₂O₃ in a mortar for 2 min.
- M. Harfenist, A. Baviey, and W. Lazier, *J. Org. Chem.*, **19**, 1608 (1954).

Reaction and Coordination Chemistry of Ferrocenylphosphines with (η⁵-C₅H₅)Co(CO)₂-Crystal Structures of Two Ferrocenylphosphine Oxides

Tae-Jeong Kim*, Jung-Hee Lee, Soon-Chul Kwon, and Kee-Ha Kwon

*Department of Industrial Chemistry,
Kyungpook National University, Taegu 702-701*

Jae-Kook Uhm*, Hyuk Lee, and Sang-In Byun

*Department of Chemistry, Keimyung University,
Taegu 704-701*

Received February 11, 1990

There has been considerable interest in reactions catalyzed by ferrocenylphosphine derivatives of metals such as Rh, Ni, Pd, Pt, and Au¹⁻¹⁰. Of various chiral and achiral ferrocenylphosphine ligands those that have been most widely explored in connection with homogeneous catalysis are (η⁵-C₅H₄PPh₂)₂Fe(BPPF), (η⁵-C₅H₅)Fe(η⁵-C₅H₃(CHMeNMe₂)PPh₂-1,2)(PPFA), and (η⁵-C₅H₄PPh₂)Fe(η⁵-C₅H₃(CHMeNMe₂)PPh₂-1,2)(BPPFA). We have recently been interested in the synthesis of rhodium and iron complexes incorporating these ligands principally for use as catalyst precursors in the catalytic hydrogenation of polynuclear heteroaromatic compounds¹¹ and in the CO₂ activation for the synthesis of carbamates^{12,13}, respectively. During the course of our investigations we have noticed that the above ligands form a wide range of complexes with various coordination modes: a typical chelating bidentate, a monodentate diphosphine, or a bridging ligand in dimeric and trimeric species.




Figure 1. Molecular Structure of **2**. Selected bond lengths (Å) and angles (°) : Fe-C(1) 2.027(3), Fe-C(3) 2.058(4), Fe-C(5) 2.030(4), Fe-C(6) 2.030(4), Fe-C(8) 2.048(4), Fe-C(10) 2.027(4), O-P 1.477(3), C(11)-P 1.808(4), C(21)-P 1.806(3), C(1)-P-O 114.6(2), C(11)-P-O 112.4(2), C(11)-P-C(1) 105.9(2), C(21)-P-O 112.5(2), C(21)-P-C(1) 105.8(2), C(21)-P-C(1) 104.8(2), C(2)-C(1)-P 128.2(3), C(5)-C(1)-P 124.9(3).

Thus, in order to extend the coordination chemistry of these ligands, we had available $(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{Co})_2$ (**1**) which is a much-studied compound in connection with cyclization of olefins and acetylenes¹⁴⁻¹⁶. This paper documents some preliminary work on the reactions of BPPF, PPFA, and BPPFA with **1** and the structure determination of two ferrocenylphosphine oxides derived serendipitously from BPPF and PPFA.

BPPF (2.27 g, 4.1 mmol) and **1** (0.53 ml, 5.3 mmol) in toluene (75 ml) were stirred under reflux for 30 h in an argon atmosphere. As the reaction proceeded, the solution became darkbrown with deposit of black precipitates. Toluene was evaporated and residue was extracted with CH_2Cl_2 (10 ml) to be chromatographed on silica gel. Two orange bands were collected on elution with CH_2Cl_2 / acetone (1/2) to give $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4\text{P}(=\text{O})\text{Ph}_2)$ (**2**) (0.25 g, 25%) and $(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{P}(=\text{O})\text{Ph}_2)$ (**3**) (0.16 g, 11%) as orange crystals after usual workups.

The structures of **2** and **3** were established by analytical and spectroscopic methods¹⁷ and the X-ray crystal structure of **2** is shown in Figure 1¹⁸.

The anticipated coordination products of the type $\text{CpCo}(\text{BPPF})_n$ ($n=1,2$) have not been formed in an any observable amount.

The reaction of PPFA with **1** under the same reaction conditions as above has also led to the oxidation at phosphorus to give a chiral phosphine oxide $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4(\text{CHMeNMe}_2)\text{P}(=\text{O})\text{Ph}_{2-1,2})$ (**4**) (45%)¹⁹ as a sole product. Single crystals suitable for the X-ray study have been obtained by cooling an acetone solution of **4** at 0°C and the crystal structure is shown in Figure 2²⁰.

One of characteristic features of **4** is that, unlike the parent PPFA, free rotation of the two N-methyl groups around the C-N axis seems to be prohibited due to increased steric congestion by introduction of oxygen on phosphorus. This





Figure 2. Molecular Structure of **4**. Selected bond lengths (Å) and angles (°) : Fe-C(1) 2.029(3), Fe-C(9) 2.064(5), O-P 1.491(3), C(1)-P 1.785(4), C(11)-P 1.806(4), C(21)-P 1.809(4), C(1N)-N 1.456(8), C(2N)-N 1.433(8), C(5N)-N 1.454(5), C(2)-C(1) 1.437(6), C(5)-C(1) 1.431(6), C(8)-C(9) 1.422(8), C(21)-C(22) 1.367(6), C(22)-C(23) 1.391(8), C(1)-P-O 115.7(2), C(11)-P-O 112.8(2), C(21)-P-O 111.7(2), C(11)-P-C(1) 106.7(2), C(21)-P-C(1) 104.3(2), C(21)-P-C(11) 104.8(2), C(2)-P-C(1) 126.0(3), C(5)-P-C(1) 126.2(4), C(51)-C(5)-C(4) 128.4(4), C(51)-C(5)-C(1) 123.6(4), C(5)-C(51)-N 108.0(4), C(52)-C(51)-N 115.7(4), C(51)-N-C(2N) 115.7(5), C(2N)-N-C(1N) 111.9(5).

is evidenced by the ¹H-NMR which gives rise to two resonances for the NMe₂ groups at δ 1.63 and 1.44 ppm. Other ¹H and ³¹P-NMR patterns are as expected for the structure.

The reaction of BPPFA with **1** is somewhat different from those of BPPF and PPFA, yielding mostly the cobalt-ferrocenylphosphine complex $\text{CpCo}(\eta^1\text{-BPPFA-P})_2$ (**5**) (45%)²¹ as well as the oxidation product $(\eta^5\text{-C}_5\text{H}_4\text{P}(=\text{O})\text{Ph}_2)\text{Fe}(\eta^5\text{-C}_5\text{H}_3(\text{CHMeNMe}_2)\text{P}(=\text{O})\text{Ph}_{2-1,2})$ (**6**) (10%)²².



The complex formation in **5** is best confirmed by the presence of a sharp singlet at δ 3.92 ppm due to the unsubstituted Cp ring protons. The ³¹P-NMR is also supportive to give the different resonances at δ 26.84 (s) and -25.82 (s) ppm due to the coordinated and free phosphorus atoms, respectively. Remaining question concerning the site of coordination of BPPFA can be solved by assigning the high-field signal to the phosphine neighboring the CHMeNMe₂ group in the same ring. This can be achieved by comparing the ³¹P NMR spectra of BPPFA, PPFA, and **5**. The ¹H NMR pattern of **6** is straightforward. Thus, for example, AB protons of the monosubstituted Cp ring protons appear as two singlets at

δ 3.86 and 3.68 ppm while the disubstituted Cp ring gives rise to five broad signals expected for the ABC protons. The methine proton appears as a quartet at δ 4.26 ppm as usual.

Obviously, much work has yet to be done concerning the mechanisms for the formation of these unexpected phosphine oxides not alone their further uses as ligands for the preparation of other series of coordination compounds. The compound **5** is also interesting in that it may be used not only in the synthesis of heteronuclear metal clusters but also in homogeneous catalytic cyclization of some alkynes. These points are the subject of our future communication.

Acknowledgement. This work has been partially supported by the Ministry of Education through the Korea Research Foundation (1988) and the Korea Science and Engineering Foundation (Grant No. 88-0304-03). Dr. J. H. Cheong of Korea Institute of Science and Technology is gratefully acknowledged for the crystal structure determination.

References

- (a) T. Hayashi, K. Yamamoto, and M. Kumada, *Tetrahedron Lett.*, **15**, 4405 (1974). (b) T. Hayashi, T. Mise, S. Mitachi, K. Yamamoto, and M. Kumada, *ibid.*, **17**, 1133 (1976).
- M. Kumada, T. Hayashi, and K. Tamao, *Fundamental Research in Homogeneous Catalysis*, Plenum Press, New York, 175 (1982).
- T. Hayashi and M. Kumada, *Acc. Chem. Res.*, **15**, 395 (1982).
- W. R. Cullen, F. W. B. Einstein, T. Jones, and T. J. Kim, *Organometallics*, **2**, 741 (1983).
- W. R. Cullen, F. W. B. Einstein, T. Jones, and T. J. Kim, *Organometallics*, **4**, 346 (1985).
- T. G. Appleton, W. R. Cullen, S.V. Evans, T. J. Kim, and J. Trotter, *J. Organomet. Chem.*, **279**, 5 (1985).
- T. Hayashi, M. Konishi, Y. Kobori, M. Kumada, T. Higuchi, and K. Hirotsu, *J. Am. Chem. Soc.*, **106**, 158 (1984).
- W. R. Cullen, S. V. Evans, N. F. Han, and J. Trotter, *Inorg. Chem.*, **26**, 514 (1987).
- Y. Ito, M. Sawamura, and T. Hayashi, *J. Am. Chem. Soc.*, **108**, 6405 (1986).
- Y. Ito, M. Sawamura, and T. Hayashi, *Tetrahedron Lett.*, **28**, 6215 (1987).
- T. J. Kim and K. C. Lee, *Bull. Korean Chem.*, **10**, 279 (1989).
- T. J. Kim, K. H. Kwon, S. C. Kwon, J. O. Baeg, S. C. Shim, and D. H. Lee, *J. Organomet. Chem.*, **389**, 205 (1990).
- T. J. Kim, S. C. Kwon, K. H. Kwon, and N. H. Heo, manuscript in preparation.
- D. F. Harvey, B. M. Johnson, C. S. Ung, and K. P. C. Vollhardt, *Synlett.*, **15** (1989).
- K. P. C. Vollhardt, *Lect. Heterocycl. Chem.*, **9**, 59 (1987).
- K. P. C. Vollhardt, *Angew. Chem. Int. Ed. Engl.*, **23**, 539 (1984).
- For **2**: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.61-7.39(m, C_6H_5), 4.72(b, C_5H_4), 4.26(b, C_5H_3). $^{31}\text{P-NMR}$ (121.5 MHz, CDCl_3) δ 26.13 (s). Anal. Calcd for $\text{C}_{22}\text{H}_{19}\text{POFe}$: C, 68.42; H, 4.92. Found: C, 67.90; H, 4.70. For **3**: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.64-7.37(m, $\text{P}(\text{O})\text{Ph}_2$), 7.27(b, PPh_2), 4.57 and 4.36(b, $\text{C}_5\text{H}_4\text{P}(\text{O})\text{Ph}_2$), 4.21 and 4.03(b, $\text{C}_5\text{H}_4\text{PPh}_2$). $^{31}\text{P-NMR}$ (121.5 MHz, CDCl_3) δ 6.72 (s, $\text{P}(\text{O})\text{Ph}_2$), -19.37(s, PPh_2). Anal. Calcd for $\text{C}_{34}\text{H}_{28}\text{P}_2\text{OFe}$: C, 71.59; H, 4.91. Found: C, 70.70; H, 4.74.
- Crystal data for **2**: Space group $\text{P2}_1/c$, $a=14.172(2)$, $b=10.434(1)$, $c=11.894(2)$ Å, $\beta=90.53(1)^\circ$, $V=1758.7(5)$ Å³, $Z=4$, $D_{\text{calc}}=1.494$ gcm⁻³, Mo-K α radiation, $\lambda=0.71073$ Å, $\mu=2.29$ cm⁻¹. An Enraf-Nonius CAD-4 diffractometer, 2047 unique absorption corrected reflections with $1>3\sigma$ (1) in the range $3^\circ < 2\theta < 50^\circ$. The structure was solved by Patterson and Fourier Methods. Final $R=0.0317$, $R_w=0.0374$.
- For **4**: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.65-7.32 (m, C_6H_5), 4.41-4.25(m, C_5H_3), 4.18(s, C_5H_5), 3.89(q, $^3J=6$ Hz, CH), 1.63(s, NMe_2), 1.15(d, $^3J=6$ Hz, CMe), $^{31}\text{P-NMR}$ (121.5 MHz, CDCl_3) δ 25.03 (s). Anal. Calcd for $\text{C}_{26}\text{H}_{28}\text{NPOFe}$: C, 68.29; H, 6.13; N, 3.06. Found: C, 68.10; H, 6.18; N, 3.15.
- Crystal data for **4**: space group P2_1 , $a=9.3252(8)$, $b=11.342(1)$, $c=11.043(1)$ Å, $\beta=102.631(8)^\circ$, $V=1139.8(2)$ Å³, $Z=2$, $D_{\text{calc}}=1.331$ gcm⁻³, Mo-K α radiation, $\lambda=0.71073$ Å, $\mu=2.29$ cm⁻¹. An Enraf-Nonius CAD-4 diffractometer, 1797 unique absorption corrected reflections with $1>3\sigma$ (1) in the range $3^\circ < 2\theta < 50^\circ$. The structure was solved by Patterson and Fourier Methods. Final $R=0.0311$, $R_w=0.0344$.
- For **5**: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.59-7.14(m, C_6H_5), 4.36-4.13(m, C_5H_3), 3.92(s, C_5H_5), 3.84(q, $^3J=6$ Hz, CH), 1.76(s, NMe_2), 1.23(d, $^3J=6$ Hz, CMe). $^{31}\text{P-NMR}$ (121.5 MHz, CDCl_3) δ 26.64(s, PPh_2Co), -25.82(s, PPh_2). Anal. Calcd for $\text{C}_{81}\text{H}_{79}\text{N}_2\text{P}_4\text{Fe}_2\text{Co}$: C, 70.76; H, 5.75; N, 2.04. Found: C, 70.30; H, 5.51; N, 2.15.
- For **6**: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.55-7.32(m, C_6H_5), 4.81, 4.73, 4.59, 4.52 and 4.47(b, C_5H_3), 4.26(q, $^3J=9$ Hz, CH), 3.86 and 3.68 (b, C_5H_4), 1.56(s, NMe_2), 1.01(d, $^3J=9$ Hz, CMe), $^{31}\text{P-NMR}$ (121.5 MHz, CDCl_3) δ 26.17(s), 23.78 (s). Anal. Calcd for $\text{C}_{38}\text{H}_{37}\text{NP}_2\text{O}_2\text{Fe}$: C, 67.97; H, 5.51; N, 2.08. Found: C, 67.90; H, 5.61; N, 1.87.

Determination of Z-and E-2-Methylbut-1-en-1-ol

Chong Shik Chin,* Byeongno Lee, and Jongpil Chun

Department of Chemistry, Sogang University, Seoul 121-742

Received December 10, 1990

Fast double bond migration of allylic alcohols with metal complexes is an unique way of generating simple enols in non-aqueous solvents.^{1,2} The reaction of 2-ethylprop-2-en-1-ol with $[\text{Rh}(\text{CO})(\text{PPh}_3)_3] \text{ClO}_4$ generates a simple and relatively stable enol, 2-methylbut-1-en-1-ol, **1** containing both Z- and E-isomers according to $^1\text{H-NMR}$ data (Eq. 1).² Unambiguous assignments of the $^1\text{H-NMR}$ signals of **1**, however,

