

Articles

Electrochemical Study of Nickel(II) Complexes with Diaza-Macrocyclic Ligands in Acetonitrile

Moo-Lyong Seo*, Zun-Ung Bae†, and Tae-Myoung Park

Department of Chemistry, Gyeongsang National University, Chinju 660-701

†Department of Chemistry, Kyungpook National University, Taegu 702-701

Received February 19, 1989

The electrochemical behavior and the complex formation between Ni^{2+} and 1,7-diaza-15-crown-5 and 1,10-diaza-18-crown-6 in acetonitrile solution have been studied by DC polarography, differential pulse polarography and cyclic voltammetry. Nickel(II) complexes gave a single well-defined wave. The formation constants of their complexes were $10^{4.89}$ and $10^{3.86}$, respectively. Nickel(II) ion was found to form complexes of 1-to-1 composition with 1,7-diaza-15-crown-5 and 1,10-diaza-18-crown-6. In addition, reduction steps were irreversible and the reduction current were diffusion controlled. The electrochemical reduction mechanism of Ni(II)-macrocyclic diaza-crown complexes in acetonitrile solution is estimated.

Introduction

The alkaline and alkaline earth metal ion selectivity of polyether macrocyclic ligands have been well investigated¹⁻³. Along a given series of ligands, maximum stability is usually observed when optimum macrocyclic ring (or cavity) size is reached for a particular metal ion.

In contrast, such discrimination on the size of macrocyclic ring containing transition metal ions has been much less documented⁴⁻⁷. A major difficulty with these systems when the macrocyclic ligand contains an N-donor set is that traditional methods of determining stability constants (such as potentiometric titration) are not satisfactory with such complexes because thermodynamic equilibrium is often extremely slow to be established⁷⁻⁹.

Nevertheless, studies involving potentiometric determination of stability constants for nickel complexes of a number of mixed donor macrocyclic ligands have been carried out. Also, the electrochemical behavior of complexes with a number of macrocyclic ligands have been extensively studied in the aqueous solution¹⁰⁻¹². But relatively few studies on the electrochemical behavior have been reported in the non-aqueous solution.

Hence, this study is to investigate the complex formation and electrochemical behavior of nickel complexes of 1,7-diaza-15-crown-5 and 1,10-diaza-18-crown-6 as shown Figure 1 in acetonitrile solution by DC polarography, differential pulse polarography and cyclic voltammetry.

Experimental

Reagent. 1,7-Diaza-15-crown-5[DA15C5] and 1,10-diaza-18-crown-6[DA18C6] was obtained from Aldrich Co. and was used without further purification. Also, nickel nitrate used as a nickel ion source was obtained from Junsei Chemical Co. Acetonitrile (AN) used as solvent was purified by Coetzee method¹³ and tetraethylammonium perchlorate (TEAP)




Figure 1. Structural formulas of macrocyclic ligands.

used as supporting electrolyte was synthesized and purified by Kolthoff method¹⁴.

Polarographic and Cyclic Voltammetric Measurement. Polarographic and cyclic voltammetric measurements of complex are made with a EG & G PAR (Princeton Applied Research) Model 264A Polarographic Analyzer/Stripping Voltammeter using 1.0×10^{-3} M TEAP as supporting electrolyte. The voltammograms were recorded with X-Y recorder (Houston Series X-Y recorder).

The working electrode used for voltammetry was EG & G PARC Model 303A SMDE (static mercury drop electrode) system. The auxiliary electrode was platinum wire. The reference electrode with glass frit tip was Ag/Ag^+ [0.1 M $AgNO_3$ in AN] electrode. Then nonaqueous internal filling solution was 0.1 M $AgNO_3$ AN solution obtaining from EG & G PAR Co. All test solution were thoroughly degassed with nitrogen gas saturated with acetonitrile and a continuous stream of nitrogen gas was passed over the solution while measurement were being taken.

Results and Discussion

Polarographic and Voltammetric Behavior of Nickel Complexes. The Figure 2 shows typical DC polarograms and differential pulse polarograms of nickel(II) ion




Figure 2. DC and differential pulse polarograms of 1.0×10^{-3} M Nickel(II) and Nickel(II) complex with DA15C5 in 1.0×10^{-1} M TEAP-AN solution at 25°C . Curve (I) is DC polarographic residual current.

Table 1. Differential Pulse Polarographic Potentials (E_p) and Peak Currents (i_p) of 1.0×10^{-3} M Nickel(II) and Nickel(II) Complexes in 1.0×10^{-1} M TEAP-AN Solution at 25°C

Compounds	E_p (volts vs Ag/Ag^+ [0.1 M AgNO_3 in AN])	i_p (μA)
Ni(II)	-0.50	0.68
Ni(II)-DA15C5	-0.85	0.93
Ni(II)-DA18C6	-0.77	0.90

and its complex with DA15C5 in a AN solution containing 1.0×10^{-1} M tetraethyl-ammonium perchlorate (TEAP). Nickel(II) ion and nickel(II) complex exhibited one well-defined wave.

The differential pulse polarographic peak potentials (E_p) and peak currents (i_p) of each compound are given in Table 1. The peak currents for each complex is linearly varied at the concentration range of 2.5×10^{-4} M- 1.0×10^{-3} M.

Also, the typical cyclic voltammograms of two each complexes in acetonitrile solution at HMDE are shown in Figure 3. The cyclic voltammetric data at different scan rate (ν) for each complex are given in Table 2. The plot of peak current (i_p) versus the square root of scan rate ($\nu^{1/2}$) is linear for each complex. These results indicate that the waves are diffusion controlled.

For an irreversible system, peak potential (E_p) is shifted to a negative direction according to increase in scan rate (for a reduction process). Thus the plot of peak potential (E_p) versus $\log(\nu)$ is linear. We can obtain the transfer coefficients (α) from slope of plots. The transfer coefficients (α) determined from a slopes of these plots are given in Table 3¹⁵. In general, deviation of transfer coefficient (α) from unity is indicative of reversibility. Thus, the result of Table 3 indicate that each reduction step is irreversible.

Polarographic Determination of Stability Constants. The mole ratio of complex between Ni^{2+} and DA15C5 and DA18C6 are 1-to-1 in acetonitrile solution¹⁸. In the polarographic study the significant quantity for the stability of the complex is the half-wave potential difference of the complex and the free metal ion. The half-wave potentials of the polarographic wave of the complex are shifted to more nega-




Figure 3. Cyclic voltammograms of 1.0×10^{-3} M Nickel(II) complexes in 1.0×10^{-1} M TEAP-AN solution on HMDE at 25°C , scan rate 200 mV/sec.

Table 2. Cyclic Voltammetric Data of 1.0×10^{-3} M Nickel(II) Complexes in 1.0×10^{-1} M TEAP-AN Solution at 25°C and HMDE

Scan rates (mV/sec)	Ni(II)-DA15C5		Ni(II)-DA18C6	
	$-E_p$ (mV)	i_p (μA)	$-E_p$ (mV)	i_p (μA)
100	852	382	750	345
200	872	554	780	460
500	917	863	835	715
1,000	942	1,171	900	964

Table 3. Transfer Coefficients (α) of Nickel(II) Complexes at 25°C and HMDE

Compounds	Transfer coefficients (α)
Ni(II)-DA15C5	0.16
Ni(II)-DA18C6	0.10

tive potentials when the concentration of the ligand in solution is increased¹⁶. Particularly, in view of the irreversible process for the reduction step, the stability constant, K_{ML} can be obtained from following equation¹⁷.

$$\Delta E_{1/2} = -RT/(n\alpha F) \ln(1 + K_{ML}[L])$$

The values of logarithm of stability constant, $\log K_{ML}$ are 4.89 for DA15C5 and 3.86 for DA18C6 in 0.1 M tetraethylammonium perchlorate-acetonitrile solution, respectively. In acetonitrile solution nickel complexes of DA15C5 and DA18C6 have considerably higher stability than in protic solvent^{18,19}. This result is termed effect due to solvation²⁰. Thus, ligands exist as a protonated form in protic solvent. Also, comparing to stability of Ni(II)-DA15C5 complex is higher than those of Ni(II)-DA18C6 complex. This result can be evaluated from the relationship between ring of ligand and radius of metal ion²¹.

Change of Reduction Peak Potential of Complex by Adding Acid. To gain insight into the reduction mechanism of each complex, the change of reduction peak potential of complex by adding acid was examined. The change of reduction peak potential of differential pulse polarogram





Figure 4. The change of reduction peak potential of 1.0×10^{-3} M Ni(II)-DA15C5 complex in 1.0×10^{-1} M TEAP-AN solution by adding 1.0×10^{-5} M HClO₄ at 25°C. Amount of 1.0×10^{-5} M HClO₄ added; I=0.05, II=0.1, III=0.15 (m) respectively, Initial volume of AN solution: 5 ml.

of Ni(II)-DA15C5 complex by adding acid is shown in Figure 4. Then, 1.0×10^{-5} M HClO₄ aqueous solution was used as the adding acid. The peak potential of the differential pulse polarographic wave of the Ni(II)-DA15C5 complex is shifted to more positive potential when amounts adding acid in 5 ml AN Solution are increased. This result is due to the increase of concentration of protonated ligand with the increase of amounts of adding acid. Thus, electrochemical reduction mechanism of Ni(II)-macrocyclic diaza-crown complexes in acetonitrile solution is estimated as follows²².



Conclusion

Based on the electrochemical experimental results for Ni(II)-macrocyclic diaza-crown complexes in acetonitrile solution, reduction steps are diffusion controlled and irreversible. Formation constants of complexes were $10^{4.89}$ and $10^{3.86}$, respectively. Nickel(II) ion was found to form complexes of 1-to-1 composition with DA15C5 and DA18C6.

Acknowledgement. We are thankful for the financial supports by the Basic Research Institute program (1988), the Ministry of Education on Inorganic Analytical Chemistry.

References

1. J. J. Christensen, D. J. Eatough, and R. M. Izatt, *Chem. Rev.*, **74**, 351 (1974).
2. R. M. Izatt et al., *J. Am. Chem. Soc.*, **98**, 7620 (1976).
3. J. M. Lehn and J. P. Sauvage, *ibid.*, **97**, 6700 (1975).
4. D. H. Busch, *Acc. Chem. Res.*, **11**, 392 (1978).
5. T. E. Jones, L. L. Zimmer, L. L. Diaddario, D. B. Rorabacher, and L. A. Ochrymowycz, *J. Am. Chem. Soc.*, **97**, 7163 (1975).
6. M. Kodama and E. Kimura, *J. Chem. Soc. Dalton*, 1081 (1978).
7. A. Anichini, L. Fabbrizzi, P. Paolletti, and R. M. Clay, *ibid.*, 577 (1978).
8. F. P. Hinz and D. W. Margerum, *Inorg. Chem.*, **13**, 2941 (1974).
9. I. Tabushi and M. Fujiyoshi, *Tetrahedron Lett.*, 2157 (1978).
10. A. Hofmanova, J. Koryta, M. Brezina, and M. L. Mittal, *Inorg. Chim. Acta*, **28**, 73 (1978).
11. M. Kodama and E. Kimura, *J. Chem. Soc. Dalton*, 1473 (1977).
12. D. P. Zollinger and M. Bos, *Anal. Chim. Acta*, **161**, 83 (1984).
13. J. F. Coetzee, *Anal. Chem.*, **34**, 1139 (1962).
14. I. M. Kolthoff and J. F. Coetzee, *J. Am. Chem. Soc.*, **79**, 1870 (1957).
15. A. J. Bard and L. R. Faulkner, "Electrochemical method, Fundamental and Application" pp. 222-224, John Wiley and Sons, New York, 1980.
16. *Idem.*, *ibid.*, pp. 163-164, John Wiley and Sons, New York, 1980.
17. K. Angelish, M. Brezina, and J. Koryta, *Electroanal. Chem. and Interf. Electrochem.*, **45**, 504 (1973).
18. E. Luboch, A. Cygan, and J. F. Biernat, *Inorg. Chim. Acta*, **68**, 201 (1983).
19. F. Arnaud-Neu, B. Spiess, and M. J. Schwing-weill, *J. Helv. Chim. Acta*, **60**, 2633 (1970).
20. *Idem.*, *J. Am. Chem. Soc.*, **104**, 5641 (1982).
21. M. Kodama and E. Kimura, *J. Chem. Soc. Dalton*, 2341 (1976).
22. *Idem.*, *ibid.*, 104 (1978).