

Three Brackish Water Rotifers from Korea

Song, Min Ok and Kim, Won

(Department of Molecular Biology, College of Natural Sciences, Seoul National University, Seoul 151-742, Korea)

한국 기수산 윤형동물 3종

송 민 옥 · 김 원

(서울대학교 자연과학대학 분자생물학과)

적 요

1984년 4월에 강원도 일원의 경포호, 송지호, 영랑호 등의 기수역에서 채집된 윤형동물을 동정한 결과 *Notholca liepetterseni*, *Keratella cruciformis eichwaldi*, *Diplochlanis propatula propatula*의 2과 1종, 2아종이 확인되어 이들을 모두 재기재하였고 아울러 도판을 작성하였다. 이중 *Notholca liepetterseni*는 한국 미기록종이다.

Key words: taxonomy, brackish water rotifers, Korea.

INTRODUCTION

Since Hada (1936) had reported 14 species of Korean freshwater Rotifera, limnological and ecological studies of this taxon had been continued until Song and Kim (1989) carried out taxonomic study for the first time in Korea. After Song & Kim (1989), many taxonomic works have been performed (Chung et al., 1990, 1991a,b, 1992a,b; Kim et al., 1991). As a result, 134 species, 15 subspecies, 9 varieties, and 9 forms of rotifers have been recorded from Korea so far (Chung et al., 1992a), but all these records have been

concentrated on freshwater species. For a taxonomic study on brackish water rotifers, we examined specimens collected from three brackish water lakes on the east coast of Kangwon-do, Korea in April of 1989. As a result, *Notholca liepetterseni*, *Keratella cruciformis eichwaldi*, *Dipleuchlanis propatula propatula* were identified. Among them, *Notholca liepetterseni* is newly recorded from Korea and the other two species were not studied taxonomically and reported just in lists of limnological studies (Yamamoto, 1953; Cho, 1976; Mizuno et al., 1980). So, we redescribed and illustrated all of the identified species.

The classification system is based on Ruttner-Kolisko (1974), Koste (1978) and Bartoš (1959).

SYSTEMATIC ACCOUNT

Phylum Rotifera 윤형동물 문

Class Monogononta Plate, 1889 단성 강

Order Ploima Hudson and Gosse, 1886

Family Brachionidae Wesenberg-Lund, 1899

Genus *Notholca* Gosse, 1886

1. *Notholca liepetterseni* Björklund, 1972

(Fig. 1f)

Notholca liepetterseni Björklund, 1972 (p. 53, figs. 14E-H, table 4B); Koste, 1978 (p. 125, pl. 28, figs. 3E-H).

Material examined: 3 inds., Lake Yöngnang, April 22, 1989 (S. M. Yoon, G. S. Min and M. O. Song).

Description: Lorica oval, rounded caudally, 125-129 µm long and 85-87 µm wide. Anterodorsal margin with 6 spines; 2 median spines 9-18 µm long and slightly shorter than lateral spines; 2 submedian spines shortest and 3-8 µm long; 2 lateral spines 16-20 µm long. Anteroventral margin without spines and with wide, shallow and slightly serrated median notch. 2 sharp, long and mobile lateral spines present on posterior half of lorica and 22-29 µm long.

Remarks: In size and shape, *N. liepetterseni* is very similar to *N. striata* (O. F. Müller, 1786), but the mobile lateral spines, in mean, are markedly longer in *N. liepetterseni* (Björklund, 1972; Koste, 1978). Björklund (1972) performed biometrical analysis with samples from Norway and eastern USA and stated that there was no overlapping in spine length between these two species. *N. striata* has vertical stripes on dorsal plate, but *N. liepetterseni* does not (Koste, 1978). Hada (1939) recorded *N. striata* var. *biremis* (Ehrenberg, 1832) from marine and brackish waters in Japan, which is very similar to *N. liepetterseni* in sizes of lorica length and mobile lateral spines. We think that Hada (1939)'s *N. striata* var. *biremis* may be *N. liepetterseni* or a closely related species of it. *N. liepetterseni* is cold-stenothermic and dominates in winter and early spring, while *N. striata* dominates in the late spring and early summer (Björklund, 1972). *N. liepetterseni* is newly reported from Korea.

Genus *Keratella* Bory de St. Vincent, 1822

2. *Keratella cruciformis eichwaldi* (Levander, 1894)

(Fig. 1e)

Anuraea eichwaldi Levander, 1894 (p. 62, pl. 3, fig. 41).

Anuraea cruciformis var. *eichwaldi* Levander, 1901 (cited from Ahlstrom, 1943).

Keratella cruciformis var. *eichwaldi*: Ahlstrom, 1943 (p. 453, pl. 38, fig. 11); Yamamoto, 1952 (p. 28, fig. 72);




Fig. 1. a-d, *Dipleuchlanis propatula propatula* (Gosse, 1886): a, whole animal; b, whole animal with head; c, trophi (M: manubrium; U: unci; R: rami; F: fulcrum); d, cross section of lorica. e, *Keratella cruciformis eichwaldi* (Levander, 1894). f, *Notholca liepetterseni* Björklund, 1972. (Scales: a, b, c = 50 μm ; e, f = 10 μm)

Yamamoto, 1953 (p. 156, pl. 1, figs. 14, 15); Yamamoto, 1960 (p. 391, fig. 6A); Hada, 1939 (p. 498, fig. 1).

Keratella cruciformis f. *eichwaldi*: Koste, 1978 (p. 110, pl. 23, fig. 14).

Keratella eichwaldi: Koste & Shiel, 1987 (p. 1006, fig. 29: 3).

Material examined: 11 inds., Lake Yǒngnang, April 22, 1989 (S. M. Yoon, G. S. Min and M. O. Song); 20 inds., Lake Songji, April 21, 1989 (S. M. Yoon, G. S. Min and M. O. Song); 2 inds., Lake Kyǒngp'o, April 22, 1989 (S. M. Yoon, G. S. Min and M. O. Song).

Description: Lorica subovate, compressed dorsoventrally, 145-158 µm long and 88-100 µm wide. No posterior spines. Anterodorsal margin with 6 spines, which slightly divergent; 2 median spines longest. Dorsal plate slightly granulated and with 2 pairs of large plaques along median line, which extending about full length of lorica. Transverse ridge of antero- and posterocarinal plaques interrupted. Lateral and caudal margins of lorica with 9 pairs of lateral and marginal plaques and 1 posteromedian remnant. Anteroventral margin without spines, round and with shallow median notch.

Family Euchlanidae Bartoš, 1959

Genus *Dipleuchlanis* de Beauchamp, 1910

3. *Dipleuchlanis propatula propatula* (Gosse, 1886)

(Fig. 1a-d)

Diplois propatula Gosse in Hudson and Gosse, 1886 (vol. 2, p. 87, pl. 24, fig. 2).

Euchlanis subversa Bryce, 1890 (p. 77, figs. 44, 45).

Dipleuchlanis propatula: Collin et al., 1912 (P. 164, figs. 319, 320); Bartoš, 1959 (p. 405, pl. 72, figs. H, CH, J); Yamamoto, 1949 (p. 44, fig. 9); Mamaril and Fernando, 1978 (p. 127, fig. 62); Myers, 1930 (p. 380, pl. 22, figs. 5-7).

Dipleuchlanis propatula propatula: Koste, 1978 (p. 144, pl. 40, figs. 6a-e); Koste and Shiel, 1989 (p. 86, fig. 2: 2a-f).

Material examined: 15 inds., Lake Kyǒngp'o, April 22, 1989 (S. M. Yoon, G. S. Min and M. O. Song).

Description: Lorica oval in outline, truncate in front, and rounded posteriorly. Dorsal plate concave and smaller than ventral one all around; posterior portion gradually narrowing, obtusely pointed at posterior end; with width about 2/3 of ventral one, 150-163 µm long and 92-100 µm wide. Ventral plate convex, 150-170 µm long and 124-139 µm wide. Dorsal and ventral plates connected with deep sulci (Fig. 1d). Inner margins of sulci gradually narrowing, tapering abruptly at posterior portion and making spine like shape under microscope. Lateral edges of two plates almost parallel and somewhat elevated. Head connected with anterior margin of lorica by a cuticular membrane. Foot of 3 sections. Two cylindrical toes very long (75-86 µm), about 1/2 of dorsal plate length, slender, parallel sided, and tapering to short points, without swelling before tips. Trophi (Fig. 1c) modified malleate type; rami triangular in ventral view; unci consisting of 8 long teeth, which decreasing in size toward dorsal part and with swollen tips.

Remarks: In his original description, Gosse (in Hudson and Gosse, 1886) described the inner margins of lateral sulci as the edges of a cleft on dorsal plate and placed this species in the genus *Diplois* of family Salpinidae (= family Mytilinidae). He also erroneously stated that the smaller plate was ventral and there were three acute spines at the base of foot. Bryce (1890) corrected these errors. He found the smaller plate was dorsal and there was no cleft on it. He also clarified that Gosse's (in Hudson and Gosse, 1886) three spines at the base of foot were just optical presentments of acute angles formed by inner margins of sulci of lorica, posterior margin of dorsal plate and a side protection of a foot. We agree with Bryce (1890).

ABSTRACT

One species and two subspecies of rotifers inhabiting several brackish water lakes on the east coast of Korea are redescribed and illustrated. *Keratella cruciformis eichwaldi* and *Diplechlanis propatula propatula* are the known species, and *Notholca liepetterseni* is newly reported from Korea.

REFERENCES

- Ahlstrom, E. H., 1943. A revision of the rotatorian genus *Keratella*, with descriptions of three new species and five new varieties. Bull. Am. Mus. nat. Hist., **77**: 453.
- Bartoš, E., 1959. Výřínci-Rotatoria. Fauna ČSR, Praha, 15, pp. 1-965.
- Björklund, B. G., 1972. Taxonomic and ecological studies of species of *Notholca* (Rotatoria) found in sea- and brackish water, with description of a new species. Sarsia, **51**: 53-57.
- Bryce, D., 1890. Two new species of Rotifera. Sci. Gossip, London, **26**: 77-79.
- Cho, K. S., 1976. Limnological studies on the Asan and Namyang Reservoirs to be varying as freshwater. Reports of Science Education. no. **2**: 74.
- Chung, C. E., H. B. Yoo and S. Y. Kim, 1990. A new record of the 3 species on freshwater Rotifera in Korea. Korean J. Limnol., **23**, **4**: 279-285.
- Chung, C. E., H. B. Yoo and S. Y. Kim, 1991a. Freshwater Rotifera of Korea. I. Family Lecanidae (Rotifera: Monogononta). Korean J. Limnol., **23**: 207-225.
- Chung, C. E., H. B. Yoo and S. Y. Kim, 1991b. Freshwater Rotifera of Korea. II. Family Colurellidae (Rotifera: Monogononta). Korean J. Syst. Zool., **7**, **2**: 241-256.
- Chung, C. E., H. B. Yoo and S. Y. Kim, 1992a. Rotifera from Korean inland waters IV. *Brachionus* and *Platyias* of Brachionidae (Rotifera: Monogononta). Korean J. Syst. Zool., **8**, **1**: 35-56.
- Chung, C. E., H. B. Yoo and S. Y. Kim, 1992b. Freshwater Rotifera of Korea III. Family Notommatidae (Rotifera: Monogononta). Korean J. Limnol., **25**, **1**: 1-8.
- Collin, A., H. Dieffenbach, R. Sachse and M. Voigt, 1912. Rotatoria und Gastrotricha. Süßwasserfauna Deutschlands, H.**14**: 164.
- Hada, Y., 1936. The plankton of Lake Seiko, Suigen in autumn. Chosen Nat. Hist. Soc. J., **21**: 1-11.
- Hada, Y., 1939. Some Rotatoria from sea and brackish water. Zool. Mag., **51**: 496-503.
- Hudson, C. T. and P. H. Gosse, 1886. The Rotifera or wheel-Animalcules. London, Longmans, Green, & Co., I: VI + 128pp.; II: 144pp.
- Kim, W., S. Y. Moon, and M. O. Song, 1991. The systematic study on the freshwater Rotifera of Korea. Korean J. Zool., **34**, **4**: 548-556.
- Koste, W., 1978. Die Räderterre Mitteleuropas. Überordnung Monogononta. Begründet von M. Voigt. I. Textbd. VIII + 673pp.; II. Tafelbd. II + 476pp. mit 234 Tafeln, Stuttgart.
- Koste, W. and R. J. Shiel, 1987. Rotifera from Australian inland waters. II. Epiphanidae and Brachionidae (Rotifera: Monogononta). Invertebr. Taxon., **7**: 1006.

- Koste, W. and R. J. Shiel, 1989. Rotifera from Australian inland waters. III. Euchlanidae, Mytilinidae and Trichotriidae (Rotifera: Monogononta). *Trans. R. Soc. S. Aust.*, **113**: 86.
- Levander, K. M., 1894. Materialien zur Kenntniss der Wasserfauna in der Umgebung von Helsingfors, mit besonderer Berücksichtigung der Meeresfauna. II. Rotatorien. *Acta Soc. Fauna Flora Fenn.*, **12**, **3**: 62-63.
- Mamaril, A. C. and C. H. Fernando, 1978. Freshwater zooplankton of the Philippines (Rotifera, Cladocera, and Copepoda). *Natural and Applied Science Bulletin*, **30**, **4**: 127.
- Mizuno, T., S. W. Kang and K. S. Cho, 1980. A faunal study on freshwater plankton from Korea. *Investigation of Scientific Study Abroad*. The Ministry of Education, pp. 71-72 (in Japanese).
- Myers, F. J., 1930. The Rotifer fauna of Wisconsin. V. The genera *Euchlanis* and *Monommata*. *Trans. Wis. Acad. Sci. Arts Lett.*, **25**: 380-381.
- Ruttner-Kolisko, A., 1974. Plankton rotifers: biology and taxonomy (translated by G. Kolisko). *Binnengewässer*, **26**, **1** (Suppl.): 1-146.
- Song, M. O. and H. S. Kim, 1989. Monogonont Rotifers (Monogononta: Rotifera) inhabiting several lowland swamps in Kyōngsangnam-do, Korea. *Korean J. Syst. Zool.*, **5**, **2**: 139-157.
- Yamamoto, K., 1949. The rotifer fauna of Japanese inland waters. I. *Jpn. J. Limnol.*, **14**: 44.
- Yamamoto, K., 1952. The rotifer fauna of Japanese inland waters. VII. *Jpn. J. Limnol.*, **16**: 28-29.
- Yamamoto, K., 1953. Preliminary studies on the rotatorian fauna of Korea. *Pacif. Sci.*, **7**, **2**: 156-157.
- Yamamoto, K., 1960. Plankton Rotatoria in Japanese inland waters. *Hydrobiologia*, **16**, **4**: 391.

RECEIVED: 28 OCTOBER 1992

ACCEPTED: 14 NOVEMBER 1992