

rum (KBr pallet, cm^{-1}) 3070 w, 3020 w, 2920 w, 2890 w, 2570 s, 2360 w, 1600 w, 1495 m, 1450 w, 1380 w, 1260 w, 1070 w, 1030 s, 1000 w, 980 w, 970 w, 940 w, 900 w, 860 m, 800 w, 755 m, br, 740 m, br, 700 s, 670 w, 650 w, 620 w, 600 w, 530 w, 485 w, 470 w.

10. ^{13}C NMR (160.5 MHz, ppm, CD_3CN) 11.3 (d, $J_{\text{BH}}=160$ Hz), 8.4 (d, $J_{\text{BH}}=145$ Hz), -7.0 (d, $J_{\text{BH}}=145$ Hz), -8.8 (d, $J_{\text{BH}}=145$ Hz), -16.8 (d, $J_{\text{BH}}=130$ Hz), -21.5 (s), -42.6 (d, $J_{\text{BH}}=130$ Hz).

Stereochemical Process in the 1,4-Addition of a Hydroxyl Group to α,β -Unsaturated Carboxylic Esters

Hyo Won Lee*, Ihl-Young Choi Lee†, and Il Hyun Park‡


Department of Chemistry, Chungbuk National University,
Cheongju, Chungbuk 360-763

†Korea Research Institute of Chemical Technology,
Taejeon 305-606

‡Department of Biochemistry, Chungnam National University,
Taejeon 305-764

Received December 9, 1991

As a part of our synthetic effort toward tetronasin¹ we had a plan of the construction of tetrahydrofuran fragment utilizing the intramolecular 1,4-addition reaction of a hydroxyl group to α,β -unsaturated esters. And it was necessary for us to investigate the stereochemical process about this kind of reaction because the stereochemical outcome of this reaction was crucial in our synthetic scheme. Herein we report stereochemical process in the synthesis of tetrahydrofuran rings *via* 1,4-addition of a hydroxyl group to α,β -unsaturated carboxylic esters.




During our synthetic routes to tetronasin, Wittig reaction² of tetrahydrofuranoid hemiacetal **1a-1c** gave the esters of (*E*)-6-hydroxy-2-enoic acids (**2a-2e**), which were subsequently converted to the corresponding tetrahydrofuran rings **3a-3e** *via* intramolecular 1,4-addition. To see the stereochemical outcome for the formation of the tetrahydrofuran rings, we examined the role of γ -substituents in the stereochemical induction and the results are outlined in Table I. The stereochemical assignment of the tetrahydrofuran rings was determined by ^1H -NMR spectroscopy^{3,4}. Furthermore, in order

Table 1. Cyclization of α,β -Unsaturated Esters

Entry	α,β -Unsaturated Esters	Method ^a	Products	Yield ^b (ratio ^c)
1		A		94% (2:1)
2		A		92% (3:1)
3		B		93% (4:5)
4		A		84% (2:3)
5		B		62% (1:3)

^aMethod A: NaOEt (2 eq). Method B: NaO^tBu (2 eq.). ^bYields refer to isolated products after chromatographic purification. ^cRatios refer to isomeric products and are based on 300 MHz ^1H -NMR analysis of reaction mixtures.

to determine the stereochemistry of the compound **3a**, the compound **2a** was converted to the reduced alcohol derivative of **3a** *via* a series of reactions including epoxidation-cyclization process⁵.



It is noteworthy that all of the compounds **2a-2e** ended as 2,3-*trans* derivatives of tetrahydrofuran rings. The *trans* relationship between C-4 substituent and side chain of the tetrahydrofuranyl ester at C-1 can be accounted by the cyclization *via* more stable conformation as shown Figure 1, where R_M and R_L stand for medium and large substituents at γ -position of the α,β -unsaturated ester, respectively.

Relating to the stereochemical role of the hydroxyl substituents at γ -position of the α,β -unsaturated esters, Felkin's model predicts that the nucleophile would attack the face opposite to the allylic oxygen function because of the anti-bonding effects of secondary orbital⁶. Thus, conformations of **2a-2e** contributing to the cyclization to tetrahydrofuran rings in our system are shown in the Figure 1. Accordingly,




Figure 1.


the stereochemistry at C-4 of α,β -unsaturated esters governs the stereochemical outcome for the formation of tetrahydrofuran esters. The rationale was also supported by the fact that even when there was no γ -substituent for α,β -unsaturated carboxylic ester such as the case for ring closure reaction toward nonactic acid, the predominant product was known to have 2,3-*cis* relationship⁷. Also similar stereochemical result can be found for the intramolecular conjugate addition of carbamate group in the α,β -unsaturated esters⁸. But it is interesting that when we prepared *Z* isomer of **2a** and carried out the cyclization to examine the effect of γ -substituent on stereoselectivity, we had the same stereochemical results as that of *E* isomer of **2a**. On the contrary to the previous report that the stereoselectivity for the conjugate addition of nucleophile to Michael acceptors depends upon *Z/E* configurations of Michael acceptors⁹, our results imply that the stereochemical induction for Michael acceptor was insensitive to *Z/E* configurations in our system. Perhaps $A^{1,3}$ strain due to the ester group was almost same as that due to 2-methyl group and thus the steric effect of γ -substituent became dominant in the process of stereochemical induction. However, it is still open to the question that the how much the steric effect of α -substituent of *Z/E* isomers can contribute to the stereochemical outcome along with that of competing γ -substituent in the intramolecular Michael reactions leading to tetrahydrofuran rings.

Acknowledgment. We would like to thank Korea Ministry of Science and Technology for financial support of this work.

References

- H. W. Lee, I.-Y. C. Lee, and S. K. Kim, *Tetrahedron Lett.*, **31**, 7637 (1990).
- (a) H. Ohri, G. H. Jones, J. G. Moffatt, M. L. Madox, A. T. Christensen, and S. K. Byram, *J. Am. Chem. Soc.*, **97**, 4602 (1975); (b) A. G. M. Barrett and H. G. Sheth, *J. Org. Chem.*, **48**, 5017 (1983).
- (a) P. A. Bartlett, J. D. Meadows, and E. Ottow, *J. Am. Chem. Soc.*, **106**, 5304 (1984); (b) P. A. Bartlett and C. Chapuis, *J. Org. Chem.*, **51**, 2799 (1986).
- R. D. Walkup and G. Park, *Tetrahedron Lett.*, **29**, 5505 (1988).
- The structure of **3a** was determined by the chemical conversion of **2a** to tetrahydrofuran **4** through Sharpless asymmetric epoxidation-cyclization and the deoxygenation at

tertiary alcohol according to Corey's procedure (E. J. Corey and P. B. Hopkins, *Tetrahedron Lett.*, **23**, 1979 (1982).) and comparison of ¹H-NMR spectrum of **4** with those of the reduced forms of **3a**.



- (a) M. Chérest, H. Felkin, and N. Prudent, *Tetrahedron Lett.*, 2199 (1968); (b) P. Caramella, N. G. Rondan, M. N. Paddon-Row, and K. N. Houk, *J. Am. Chem. Soc.*, **103**, 2438 (1981); (c) K. N. Houk, M. N. Paddon-Row, N. G. Rondan, Y.-D. Wu, F. K. Brown, D. C. Spellmeyer, J. T. Metz, Y. Li, and R. J. Loncharich, *Science*, **231**, 1108 (1986).
- von H. Gerlach and H. Wetter, *Helv. Chim. Acta*, **57**, 2306 (1974).
- M. Hirma, T. Shigemoto, and S. Ito, *J. Org. Chem.*, **52**, 3342 (1987).
- Y. Yamamoto, *Acc. Chem. Res.*, **20**, 243 (1987).

Synthesis of Specifically Deuterated DNA Hexamer

Jin-Young Park, Ki-Hang Choi, Bok-Luel Lee[†], and Byong-Seok Choi*

Department of Chemistry, Korea Advanced Institute of Science and Technology, Taejeon 305-701

[†]College of Pharmacy, Kyungshung University, Pusan 608-736

Received December 9, 1991

The quinoxaline family of antibiotics of which echinomycin is a member are powerful antimicrobial and antitumor agents. The single-crystal X-ray study¹ of a echinomycin complex with d(CGACG) has shown that, surprisingly, the two central A-T base pairs are of the Hoogsteen type (Figure 1). Two-dimensional NMR studies of echinomycin complexes with d(ACGT) and d(ACGTACGT) duplexes have been reported^{2,3}. The van der Waals contacts detected in X-ray crystallographic analysis of are echinomycin-oligonucleotide co-

*To whom correspondence should be addressed.