

**CONFIDENCE INTERVALS IN THREE-FACTOR
NESTED VARIANCE COMPONENT MODEL(FOR
THE RATIO OF TOTAL VARIANCES)**

KWAN-JOONG KANG

1. Introduction

Consider the three-factor nested variance component model given by

$$(1.1) \quad y_{ijklm} = \mu + A_i + B_{ij} + C_{ijk} + \varepsilon_{ijklm},$$

where $i = 1, 2, \dots, I$, $j = 1, 2, \dots, J$, $k = 1, 2, \dots, K$ and $m = 1, 2, \dots, M$. And A_i , B_{ij} , C_{ijk} and ε_{ijklm} are independent unobservable random variables and $A_i \sim N(0, \sigma_A^2)$, $B_{ij} \sim N(0, \sigma_B^2)$, $C_{ijk} \sim N(0, \sigma_C^2)$, $\varepsilon_{ijklm} \sim N(0, \sigma_\varepsilon^2)$, μ is an unknown parameter, and the y_{ijklm} are observable random variables(Graybill).

The problem of confidence intervals on linear combinations of more than two variances is suggested by Smith in 1936. And in 1946, Satterthwaite studied and expanded the method and the result of his studies has been known as "Satterthwaite procedure".

Lately, the procedure has been widely used and developed by many authors. Especially, Howe (1974), Graybill and Wang (1979), Leiva and Graybill (1986) got a good approximation. The precisions of confidence interval are satisfactory, but it is the result mostly found out in one or two factor model. In 1993, Kang is to use and expand of Graybill and Wang's (1979) approximation, the approximation $1 - \alpha$ lower and upper confidence intervals on σ_A^2 , σ_B^2 , σ_C^2 , σ_ε^2 and the ratio of these of (1.1) in three-factor.

The purpose of this paper, there is no method available for setting exact $1 - \alpha$ confidence intervals on the ratio of total variances of these, so by using this approximate $1 - \alpha$ confidence intervals and the precisions will be treated in Section 2,3 and 4.

2. Three-factor nested variance component model

In the model (1.1), the computations are displayed in Table 1, where the sums are over i, j, k and m as specified in the model.

Source	D.F.	S.S.	M.S.	E.M.S.
Total	$IJKM$	$\sum \sum \sum \sum y_{ijklm}^2$		
Mean	1	$y^2/(IJKM)$		
Factor A	n_1	$\sum - \sum \sum \sum (\bar{y}_{i...} - \bar{y}_{....})^2$	S_1^2	θ_1
B within A	n_2	$\sum \sum \sum \sum (\bar{y}_{ij..} - \bar{y}_{i...})^2$	S_2^2	θ_2
C within B within A	n_3	$\sum \sum \sum \sum (\bar{y}_{ijk.} - \bar{y}_{ij..})^2$	S_3^2	θ_3
Error	n_4	$\sum \sum \sum \sum (y_{ijklm} - \bar{y}_{ijk.})^2$	S_4^2	θ_4

where $n_1 = I - 1, n_2 = I(J - 1), n_3 = IJ(K - 1), n_4 = IJK(M - 1), \theta_1 = \sigma_e^2 + M\sigma_C^2 + KM\sigma_B^2 + JKM\sigma_A^2, \theta_2 = \sigma_e^2 + M\sigma_C^2 + KM\sigma_B^2, \theta_3 = \sigma_e^2 + M\sigma_C^2, \theta_4 = \sigma_e^2$. The upper α probability point of Snedecor's F distribution and chi-square distribution are denoted by $F_{\alpha; \dots}$ and $\chi_{\alpha; \dots}^2$, respectively, where $\alpha_1 + \alpha_2 = \alpha$, often $\alpha_1 = \alpha_2 = \frac{\alpha}{2}$. And

$$P \left[\frac{S_i^2}{\theta_i} \leq \frac{\chi_{\alpha; n_i}^2}{n_i} = F_{\alpha; n_i, \infty} \right] = 1 - \alpha, i = 1, 2, 3, 4.$$

3. The confidence intervals on the ratio of variances

3.1. Confidence intervals on $\sigma_A^2/(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2)$

At first, we want to find out the form of the $1 - \alpha$ upper confidence interval on $\frac{\sigma_A^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2}$. Then, the form $\frac{\sigma_A^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2}$ is

$$\frac{JKM\sigma_A^2}{JKM(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2)} = \frac{\theta_1 - \theta_2}{\theta_1 + (J-1)\theta_2 + J(K-1)\theta_3 + JK(M-1)\theta_4}.$$

In this, we consider, the lower limit of $1 - \alpha$ upper confidence interval on $\frac{J\sigma_A^2}{\sigma_B^2 + \sigma_C^2 + \sigma_e^2}$. Then it is a function (Graybill and Wang) of S_1^2, S_2^2, S_3^2 and

S_4^2 say $q(S_1^2, S_2^2, S_3^2, S_4^2)$, such that $P \left[q \leq \frac{\theta_1 - \theta_2}{\theta_2 + (K-1)\theta_3 + K(M-1)\theta_4} \right] = 1 - \alpha$. We want the confidence interval to be an exact $1 - \alpha$ interval.

1) When $\sigma_C^2 = \sigma_\epsilon^2 = 0$ (i.e. $\theta_3 = 0, \theta_4 = 0$), we get

$$\frac{\theta_1 - \theta_2}{\theta_2 + (K - 1)\theta_3 + K(M - 1)\theta_4} = \frac{\theta_1}{\theta_2} - 1$$

and

$$P \left[\frac{S_1^2}{S_2^2 F_{\alpha; n_1, n_2}} - 1 \leq \frac{\theta_1}{\theta_2} - 1 \right] = 1 - \alpha.$$

2) When the hypothesis $H_0 : \sigma_A^2 = 0$ is accepted for a size α test, we want the corresponding $1 - \alpha$ confidence interval to include zero, and when H_0 is rejected we want q to be an increasing function of S_1^2/S_2^2 . This condition is required because $\frac{\theta_1 - \theta_2}{\theta_2 + (K - 1)\theta_3 + K(M - 1)\theta_4} = \theta^*$ in $P[q \leq \theta^*] = 1 - \alpha$ is an increasing function of $\frac{\theta_1}{\theta_2}$, and if $\frac{\theta_1}{\theta_2}$ increases then its estimate would tend to increase. Hence if q does not increase as $\frac{S_1^2}{S_2^2}$ increases, clearly the confidence coefficient would deviate a large amount from $1 - \alpha$ as $\frac{\theta_1}{\theta_2}$ gets large. For a size α test of $H_0 : \sigma_A^2 = 0$ versus $H_a : \sigma_A^2 > 0$, the hypothesis H_0 is accepted if and only if $\frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}$. Thus, we want $q(S_1^2, S_2^2, S_3^2, S_4^2) = 0$ when $S_1^2 \leq S_2^2 F_{\alpha, n_1, n_2}$, $q(S_1^2, S_2^2, S_3^2, S_4^2) > 0$ is increasing in $\frac{S_1^2}{S_2^2}$ when $S_1^2 > S_2^2 F_{\alpha, n_1, n_2}$.

3) When $J \rightarrow \infty$ (hence n_2, n_3 , and $n_4 \rightarrow \infty$), we want the confidence interval to have an exact confidence coefficient $1 - \alpha$. When $J \rightarrow \infty$, it follows that $n_2 \rightarrow \infty, n_3 \rightarrow \infty$ and $n_4 \rightarrow \infty$ and from this it follows that $S_2^2 \rightarrow \theta_2, S_3^2 \rightarrow \theta_3$ and $S_4^2 \rightarrow \theta_4$ in probability also $S_2^2 + (K - 1)S_3^2 + K(M - 1)S_4^2 \rightarrow \theta_2 + (K - 1)\theta_3 + K(M - 1)\theta_4$. From these results, we get $P \left[\frac{S_1^2}{F_{\alpha, n_1, \infty}} \leq \theta_1 \right] = P \left[\frac{S_1^2}{F_{\alpha, n_1, \infty}} - S_2^2 \leq \theta_1 - \theta_2 \right] = 1 - \alpha$. Then divide the left and right sides respectively, by $S_2^2 + (K - 1)S_3^2 + K(M - 1)S_4^2$ and its equivalent value $\theta_2 + (K - 1)\theta_3 + K(M - 1)\theta_4$ to obtain $1 - \alpha = P \left[\frac{S_1^2 - S_2^2 F_{\alpha, n_1, \infty}}{\{S_2^2 + (K - 1)S_3^2 + K(M - 1)S_4^2\} F_{\alpha, n_1, \infty}} \leq \frac{\theta_1 - \theta_2}{\theta_2 + (K - 1)\theta_3 + K(M - 1)\theta_4} \right]$. Hence, when $J \rightarrow \infty$, we obtain

$$q = \begin{cases} \frac{S_1^2 - S_2^2 F_{\alpha, n_1, \infty}}{\{S_2^2 + (K - 1)S_3^2 + K(M - 1)S_4^2\} F_{\alpha; n_1, \infty}} & \text{when } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, \infty} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, \infty} \end{cases}$$

The maximum likelihood estimate of

$$\frac{\theta_1 - \theta_2}{\theta_2 + (K-1)\theta_3 + K(M-1)\theta_4} \left(= \frac{J\sigma_A^2}{\sigma_B^2 + \sigma_C^2 + \sigma_e^2} \right)$$

is the form $\frac{a_0 S_1^2 + a_1 S_2^2}{a_2 S_2^2 + a_3 S_3^2 + a_4 S_4^2}$, and determine the constants a_0, a_1, a_2, a_3 and a_4 by imposing the conditions 1), 2) and 3) of 3.1, we get $a_0 = 1$, $a_2 = -a_1 = -F_{\alpha; n_1, n_2}$. Thus,

$$q = \begin{cases} \frac{S_1^2 - S_2^2 F_{\alpha; n_1, n_2}}{S_2^2 F_{\alpha; n_1, n_2} + a_3^* S_3^2 + a_4^* S_4^2} & \text{when } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}, \end{cases}$$

where $\lim_{J \rightarrow \infty} a_3^* = (K-1)F_{\alpha; n_1, \infty}$, $\lim_{J \rightarrow \infty} a_4^* = K(M-1)F_{\alpha; n_1, \infty}$. Therefore the lower limit of a $1 - \alpha$ upper confidence interval on $\frac{\sigma_A^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2}$ is

$$(3.1.1) \quad L = \begin{cases} \frac{S_1^2 - S_2^2 F_{\alpha; n_1, n_2}}{S_1^2 + (J-1)S_2^2 F_{\alpha; n_1, n_2} + A_\alpha} & \text{when } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}, \end{cases}$$

where $A_\alpha = \{J(K-1)S_3^2 + JK(M-1)S_4^2\}F_{\alpha; n_1, \infty}$. And the upper limit of a $1 - \alpha$ lower confidence interval is obtained from (3.1.1) by replacing α with $1 - \alpha$ in the tabulated F 's. We get $P \left[\frac{\sigma_A^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_e^2} \leq U \right] = 1 - \alpha$, where

$$(3.1.2) \quad U = \begin{cases} \frac{S_1^2 - S_2^2 F_{1-\alpha; n_1, n_2}}{S_1^2 + (J-1)S_2^2 F_{1-\alpha} + A_{1-\alpha}} & \text{when } \frac{S_1^2}{S_2^2} > F_{1-\alpha; n_1, n_2} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{1-\alpha; n_1, n_2}. \end{cases}$$

If we choose $a_3^* = (K-1)F_{\alpha; n_1, n_3}$ and $A_4^* = JK(M-1)F_{\alpha; n_1, n_4}$, then the lower limit of a $1 - \alpha$ upper confidence interval(L) and the upper

limit of a $1 - \alpha$ lower confidence interval(U) on $\frac{\sigma_A^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2}$ are

$$(3.1.3) \quad L = \begin{cases} \frac{S_1^2 - S_2^2 F_{\alpha; n_1, n_2}}{S_1^2 + (J - 1) S_2^2 F_{\alpha; n_1, n_2} + A_\alpha} & \text{when } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}, \end{cases}$$

where $A_\alpha = J(K - 1) S_3^2 F_{\alpha; n_1, n_3} + JK(M - 1) S_4^2 F_{\alpha; n_1, n_4}$

$$(3.1.4) \quad U = \begin{cases} \frac{S_1^2 - S_2^2 F_{1-\alpha; n_1, n_2}}{S_1^2 + (J - 1) S_2^2 F_{1-\alpha; n_1, n_2} + A_{1-\alpha}} & \text{when } \frac{S_1^2}{S_2^2} > F_{1-\alpha; n_1, n_2} \\ 0 & \text{when } \frac{S_1^2}{S_2^2} \leq F_{1-\alpha; n_1, n_2}. \end{cases}$$

3.2. Confidence intervals on $\sigma_B^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$

Similar to that of Section 3.1, we want to find out the upper confidence interval on

$$\frac{KM\sigma_B^2}{JKM(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)} = \frac{\theta_2 - \theta_3}{\theta_1 + (J - 1)\theta_2 + J(K - 1)\theta_3 + JK(M - 1)\theta_4}.$$

We consider a function of S_1^2, S_2^2, S_3^2 and S_4^2 say q such that

$$P \left[q \leq \frac{\sigma_B^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2} \leq 1 \right] = 1 - \alpha.$$

We impose the three conditions on q ;

1) When $\sigma_C^2 = \sigma_\epsilon^2 = 0$ (i.e. $\theta_3 = \theta_4 = 0$), we want the confidence coefficient to be exactly $1 - \alpha$. When $\theta_3 = \theta_4 = 0$, it follows that $\frac{\theta_2}{\theta_1 + (J - 1)\theta_2}$, and its inverse is $\frac{\theta_1 + (J - 1)\theta_2}{\theta_2} = \frac{\theta_1}{\theta_2} + (J - 1)$. If $\theta_3 = \theta_4 = 0$ implies $S_3^2 = S_4^2 = 0$ with probability 1. Thus we get

$$1 - \alpha = P \left[\frac{S_1^2}{S_2^2 F_{\alpha; n_1, n_2}} + (J - 1) \leq \frac{\theta_1}{\theta_2} + (J - 1) \right] = P \left[\frac{S_2^2 F_{1-\alpha; n_1, n_2}}{S_1^2 + (J - 1) S_2^2 F_{1-\alpha; n_1, n_2}} \leq \frac{\theta_2}{\theta_1 + (J - 1)\theta_2} \right].$$

2) We require the confidence coefficient to be exactly $1 - \alpha$. When $J \rightarrow \infty$, it follows that $n_2 \rightarrow \infty$, $n_3 \rightarrow \infty$ and $n_4 \rightarrow \infty$, $S_2^2 \rightarrow \theta_2$, $S_3^2 \rightarrow \theta_3$ and $S_4^2 \rightarrow \theta_4$ in probability. The exact $1 - \alpha$ lower confidence interval on θ_1 is $\theta_1 \leq \frac{S_1^2}{F_{1-\alpha, n_1, \infty}}$. And we get $\theta_1 + (J-1)\theta_2 + J(K-1)\theta_3 + JK(M-1)\theta_4 \leq \frac{S_1^2}{F_{1-\alpha, n_1, \infty}} + (J-1)S_2^2 + J(K-1)S_3^2 + JK(M-1)S_4^2$. Thus, q is

$$q = \frac{S_2^2 - S_3^2}{\frac{S_1^2}{F_{1-\alpha, n_1, \infty}} + (J-1)S_2^2 + J(K-1)S_3^2 + JK(M-1)S_4^2} \leq \frac{\theta_2 - \theta_3}{\theta_1 + (J-1)\theta_2 + J(K-1)\theta_3 + JK(M-1)\theta_4}$$

3) When $\theta_2 = \theta_3$, we require the lower limit of the confidence interval to be zero when the α level test of $H_0 : \theta_2 = \theta_3$ is accepted. Thus,

$$q(S_1^2, S_2^2, S_3^2, S_4^2) \begin{cases} > 0 & \text{if } \frac{S_2^2}{S_3^2} > F_{\alpha; n_2, n_3} \\ = 0 & \text{if } \frac{S_2^2}{S_3^2} \leq F_{\alpha; n_2, n_3}, \end{cases}$$

we get $q = \frac{S_2^2 - S_3^2 F_{\alpha, n_2, n_3}}{\frac{S_1^2}{F_{1-\alpha, n_1, n_2}} + (J-1)S_2^2 + J(K-1)S_3^2 F_{\alpha, n_2, n_3} + JK(M-1)S_4^2}$. We choose

the following function for $q(S_1^2, S_2^2, S_3^2, S_4^2)$; $q = \frac{S_3^2 \left[\frac{S_2^2}{S_3^2} + a_1 + a_2 \left(\frac{S_2^2}{S_3^2} \right)^{-1} \right]}{a_3 S_1^2 + S_2^2 + a_4 S_3^2 + a_5 S_4^2}$. The constant a_1, a_2, a_3 and a_4 are determined by forcing q to satisfy conditions 1), 2) and 3) of 3.2. From $F_{\alpha; n_2, n_3} + a_1 + \frac{a_2}{F_{\alpha, n_2, n_3}} = 0$, we get $a_2 = -F_{\alpha, n_2, n_3}(a_1 + F_{\alpha; n_2, n_3})$, $a_3 = \frac{1}{F_{1-\alpha, n_1, n_2}}$, $a_1 = -F_{\alpha; n_2, \infty}$, $a_5 = JK(M-1)$ and $a_4 = J(K-1)F_{\alpha, n_2, \infty}$. Thus

$$q = \frac{S_3^2 \left[\frac{S_2^2}{S_3^2} - F_{\alpha; n_2, \infty} - F_{\alpha; n_2, n_3} (F_{\alpha; n_2, n_3} - F_{\alpha; n_2, \infty}) \left(\frac{S_2^2}{S_3^2} \right)^{-1} \right]}{\frac{S_1^2}{F_{1-\alpha, n_1, n_2}} + S_2^2 + J(K-1)S_3^2 F_{\alpha; n_2, \infty} + JK(M-1)S_4^2}$$

Let

$$B_\alpha = F_{\alpha; n_2, n_3} (F_{\alpha; n_2, \infty} - F_{\alpha; n_2, n_3}) \frac{S_3^4}{S_2^2},$$

$$C_\alpha = J(K-1)S_3^2 F_{\alpha; n_2, \infty} + JK(M-1)S_4^2 F_{\alpha; n_2, \infty},$$

the lower limit of a $1 - \alpha$ upper confidence interval on $\frac{\sigma_B^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2}$ is

$$(3.2.1) \quad L = \begin{cases} \min \left[1, \frac{J(S_2^2 - S_3^2 F_{\alpha, n_2, \infty} + B_\alpha)}{\frac{S_1^2}{F_{1-\alpha, n_1, n_2}} + (J-1)S_2^2 + C_\alpha} \right] & \text{if } \frac{S_2^2}{S_3^2} > F_{\alpha; n_2, n_3} \\ 0 & \text{if } \frac{S_2^2}{S_3^2} \leq F_{\alpha; n_2, n_3}. \end{cases}$$

And the upper limit of a $1 - \alpha$ lower confidence interval is obtained from (3.2.1), by replacing α and $1 - \alpha$ with $1 - \alpha, \alpha$ in (3.2.1). We get

$$(3.2.2) \quad P \left[\frac{\sigma_B^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2} \leq U \right] = 1 - \alpha, \text{ where}$$

$$U = \begin{cases} \min \left[1, \frac{J(S_2^2 - S_3^2 F_{1-\alpha; n_2, \infty} + B_{1-\alpha})}{\frac{S_1^2}{F_{\alpha, n_1, n_2}} + (J-1)S_2^2 + C_{1-\alpha}} \right] & \text{if } \frac{S_2^2}{S_3^2} > F_{1-\alpha; n_2, n_3} \\ 0 & \text{if } \frac{S_2^2}{S_3^2} \leq F_{1-\alpha; n_2, n_3}. \end{cases}$$

We can choose other different cases of value of a_4 and a_5 , L and U are as follows;

$$(3.2.3) \quad \begin{cases} L \text{ and } U \text{ in (3.2.1), (3.2.2)} \\ \text{when } C_\alpha = J(K-1)S_3^2 + JK(M-1)S_4^2, \end{cases}$$

$$(3.2.4) \quad \begin{cases} L \text{ and } U \text{ in (3.2.1), (3.2.2)} \\ \text{when } B_\alpha = F_{\alpha, n_2, n_3}(F_{\alpha; n_2, \infty} - F_{\alpha, n_2, n_3}). \end{cases}$$

3.3. Confidence intervals on $\sigma_C^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$

The lower limit of a $1 - \alpha$ upper confidence intervals on $\sigma_C^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$. We can shows the similar to those of Section 3.2, putting

$$D_\alpha = J(K-1)S_3^2 F_{1-\alpha; n_1, n_3} + JK(M-1)S_4^2 F_{1-\alpha, n_3, n_4},$$

and

$$q = \frac{S_3^2 - S_4^2 F_{1-\alpha; n_3, n_4}}{\frac{S_1^2}{F_{1-\alpha, n_1, n_3}} + \frac{(J-1)S_2^2}{F_{1-\alpha, n_2, n_3}} + D_\alpha}.$$

Therefore, the lower limit of $1 - \alpha$ upper confidence interval on $\sigma_C^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\varepsilon^2)$ is

$$(3.3.1) \quad L = \begin{cases} \min \left[1, \frac{JK(S_3^2 - S_4^2 F_{\alpha; n_3, n_4})}{\frac{S_1^2}{F_{1-\alpha, n_1, n_3}} + \frac{(J-1)S_2^2}{F_{1-\alpha, n_2, n_3}} + D_\alpha} \right] & \text{if } \frac{S_3^2}{S_4^2} > F_{\alpha; n_3, n_4} \\ 0 & \text{if } \frac{S_3^2}{S_4^2} \leq F_{\alpha; n_3, n_4}. \end{cases}$$

And the upper limit of a $1 - \alpha$ lower confidence interval is obtained from L by replacing α and $1 - \alpha$ with $1 - \alpha, \alpha$ in the tabulated F 's. Thus we get

$$P \left[\frac{\theta_3 - \theta_4}{\theta_1 + (J-1)\theta_2 + J(K-1)\theta_3 + JK(M-1)\theta_4} \leq U \right],$$

where

$$(3.3.2) \quad U = \begin{cases} \min \left[1, \frac{JK(S_3^2 - S_4^2 F_{1-\alpha; n_3, n_4})}{\frac{S_1^2}{F_{\alpha, n_1, n_3}} + \frac{(J-1)S_2^2}{F_{1-\alpha, n_2, n_3}} + D_{1-\alpha}} \right] & \text{if } \frac{S_3^2}{S_4^2} > F_{1-\alpha; n_3, n_4} \\ 0 & \text{if } \frac{S_3^2}{S_4^2} \leq F_{1-\alpha; n_3, n_4}. \end{cases}$$

We can choose other different case of valued of D_α , L and U are as follows;

$$(3.3.3) \quad \begin{cases} L \text{ and } U \text{ in (3.3.1), (3.3.2),} \\ \text{when } D_\alpha = J(K-1)S_3^2 + JK(M-1)S_4^2. \end{cases}$$

3.4. Confidence intervals on $\sigma_\epsilon^2/(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$

We can begin with

$$\frac{JKM(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)}{\sigma_\epsilon^2} = \frac{\theta_1 + (J-1)\theta_2 + J(K-1)\theta_3 + JK(M-1)\theta_4}{\theta_4},$$

we want to determine a function of S_1^2 , S_2^2 and S_4^2 say $q(S_1^2, S_2^2, S_4^2)$ such that $P\left[q \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4}\right]$ is approximately equal to $1 - \alpha$. Scale invariance and large sample theory suggested that $q = \frac{S_2^2}{S_4^2} h\left(\frac{S_1^2}{S_2^2}\right)$, where the function $h(\cdot)$ is determined by requiring it to satisfy the following three conditions

1) When the hypothesis $\text{test } H_0 : \theta_1 = \theta_2$ is accepted by using a size α test (i.e. $\frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}$) we require the corresponding $1 - \alpha$ confidence interval on $\frac{\theta_1 + (J-1)\theta_2}{\theta_4}$ to be exact. The exact $1 - \alpha$ confidence interval on $\frac{(J-1)\theta_2}{\theta_4}$ is $\frac{S_2^2(J-1)\left(n_1 \frac{S_1^2}{S_2^2} + n_2\right)}{S_4^2(n_1 + n_2)F_{\alpha, n_1 + n_2, n_4}} \leq \frac{(J-1)\theta_2}{\theta_4}$.

2) The confidence coefficient for the interval on $\frac{\theta_1 + (J-1)\theta_2}{(J-1)\theta_4}$ is to be exact as $J \rightarrow \infty$. To satisfy this we let $h\left(\frac{S_1^2}{S_2^2}\right) = \frac{S_1^2}{S_2^2} F_{\alpha, n_1, \infty} + J - 2 + g\left(\frac{S_1^2}{S_2^2}\right)$, where $\lim_{J \rightarrow \infty} \frac{1}{Jg\left(\frac{S_1^2}{S_2^2}\right)} = 0$.

3) The confidence coefficient is to be exact as $\sigma_A^2 \rightarrow \infty$; when $\sigma_A^2 \rightarrow \infty$ the quantity $\frac{\sigma_\epsilon^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2}$ is dominated by $\frac{\theta_1}{\theta_4}$ so we want an exact $1 - \alpha$ confidence interval on $\frac{\theta_1}{\theta_4}$. Thus we get $h\left(\frac{S_1^2}{S_2^2}\right) = \frac{S_1^2}{S_2^2} F_{\alpha; n_1, n_4}$ when $\sigma_A^2 \rightarrow \infty$ (and hence $S_1^2 \rightarrow \infty$). If we impose conditions 1), 2) and 3) of 3.4, we get the following for the approximate upper $1 - \alpha$ confidence interval on $\frac{\theta_1 + (J-1)\theta_2}{\theta_4}$ is

$$\frac{\theta_1 + (J-1)\theta_2}{\theta_4} \leq \begin{cases} \frac{S_2^2}{S_4^2 F_{1-\alpha, n_1, n_4}} \left(\frac{S_1^2}{S_2^2} + E_{1-\alpha}\right) & \text{if } \frac{S_1^2}{S_2^2} > F_{1-\alpha; n_1, n_2} \\ \frac{JS_2^2\left(\frac{n_1 S_1^2}{S_2^2} + n_2\right)}{(n_1 + n_2)S_4^2 F_{1-\alpha, n_1 + n_2, n_4}} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{1-\alpha; n_1, n_2}, \end{cases}$$

where $E_{1-\alpha} = \frac{J(n_1 F_{1-\alpha; n_1, n_2} + n_2) F_{1-\alpha; n_1, n_4}}{(n_1 + n_2) F_{1-\alpha; n_1 + n_2, n_4}} - F_{1-\alpha; n_1, n_2}$. By a simulation study, we determined that the confidence coefficients for the above interval is too large, so the following function was considered: $h\left(\frac{S_1^2}{S_2^2}\right) = b_0 + b_1\left(\frac{S_1^2}{S_2^2}\right) + b_2\left(\frac{S_1^2}{S_2^2}\right)^{-1}$, which would tend to decrease the above confidence coefficients. We now impose conditions 1), 2) and 3) to evaluate b_0 , b_1 and b_2 . So we get

$$\frac{\theta_1 + (J-1)\theta_2}{\theta_4} \leq \begin{cases} \frac{S_2^2}{S_4^2 F_{1-\alpha; n_1, n_4}} \left\{ \frac{S_1^2}{S_2^2} + Y + F_{1-\alpha; n_1, n_2} \right. \\ \left. (E_{1-\alpha} - Y) \frac{S_2^2}{S_1^2} \right\} & \text{if } \frac{S_1^2}{S_2^2} > F_{1-\alpha; n_1, n_2} \\ \frac{S_2^2 \left(\frac{n_1 S_1^2}{S_2^2} + n_2 \right)}{(n_1 + n_2) S_4^2 F_{1-\alpha; n_1 + n_2, n_4}} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{1-\alpha; n_1, n_2} \end{cases}$$

where $Y = \frac{J F_{1-\alpha; n_1, \infty}^2}{F_{1-\alpha; n_1, n_2}} - \frac{F_{1-\alpha; n_1, n_4}}{F_{1-\alpha; n_1 + n_2, n_4}}$. Thus, the $1 - \alpha$ upper confidence interval on $\frac{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_D^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_D^2}$ is

$$(3.4.1) \quad L = \begin{cases} \min\{1, S_{1-\alpha}\} & \text{if } \frac{S_1^2}{S_2^2} > F_{1-\alpha; n_1, n_2} \\ \min\{1, T_{1-\alpha}\} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{1-\alpha; n_1, n_2} \end{cases}$$

where

$$S_{1-\alpha} = \frac{JKMS_4^2}{\frac{S_2^2}{F_{1-\alpha; n_1, n_4}} \left[\frac{S_1^2}{S_2^2} + Y + F_{1-\alpha; n_1, n_2} (E_{1-\alpha} - Y) \frac{S_2^2}{S_1^2} \right] + D}$$

$$D = J(K-1)S_3^2 + JK(M-1)S_4^2,$$

$$T_{1-\alpha} = \frac{KMS_4^2}{V_{1-\alpha} + (K-1)S_3^2 + K(M-1)S_4^2}$$

The upper confidence interval on $\frac{\theta_1 + (J-1)\theta_2}{\theta_4}$ is obtained from equation 3.4-2) by replacing $1 - \alpha$ with α in the tabulated F 's. Thus we get

$$(3.4.2) \quad \begin{cases} \frac{S_2^2}{S_4^2 F_{\alpha; n_1, n_4}} \left[\frac{S_1^2}{S_2^2} + E_\alpha \right] \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ \frac{JS_2^2 \left(\frac{n_1 S_1^2}{S_2^2} + n_2 \right)}{(n_1 + n_2) S_4^2 F_{\alpha; n_1 + n_2, n_4}} \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2} \end{cases}$$

Then the probability of equation (3.4.2) were too lage. As a second approximation, let $h(\frac{S_1^2}{S_2^2}) = c_0 + c_1(\frac{S_1^2}{S_2^2}) + c_2(\frac{S_1^2}{S_2^2})^{-1}$, in 3.4-2) we choosen the above satisfies the limiting conditions. thus we get

$$(3.4.3) \quad \left\{ \begin{array}{ll} \frac{S_2^2}{S_4^2 F_{\alpha; n_1, n_4}} \left[\frac{S_1^2}{S_2^2} + (J-1)F_{\alpha; n_1, \infty} + F_{\alpha; n_1, n_2} \{E_\alpha - (J-1)F_{\alpha; n_1, \infty}\} \frac{S_2^2}{S_1^2} \right] \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} F_{\alpha; n_1, n_2} \\ \frac{-JS_2^2(\frac{n_1 S_1^2}{S_2^2} + n_2)}{(n_1 + n_2)S_4^2 F_{\alpha; n_1 + n_2, n_4}} \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} < F_{\alpha; n_1, n_2} \end{array} \right.$$

The probabilities of equation (3.4.2) were too small. Hence an upper confidence interval for $\frac{\theta_1 + (J-1)\theta_2}{\theta_4}$ is obtained by averaging the lower confidence interval on $\frac{\theta_1 + (J-1)\theta_2}{\theta_4}$ is

$$(3.4.4) \quad \left\{ \begin{array}{ll} \frac{S_2^2}{S_4^2 F_{\alpha; n_1, n_4}} \left[\frac{S_1^2}{S_2^2} + \frac{1}{2} \{ (J-1)F_{\alpha; n_1, \infty} + E_\alpha \} + \frac{1}{2} F_{\alpha; n_1, n_2} \{E_\alpha - (J-1)F_{\alpha; n_1, \infty}\} \frac{S_2^2}{S_1^2} \right] \leq \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ \frac{JS_2^2(\frac{n_1 S_1^2}{S_2^2} + n_2)}{(n_1 + n_2)S_4^2 F_{\alpha; n_1 + n_2, n_4}} > \frac{\theta_1 + (J-1)\theta_2}{\theta_4} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}. \end{array} \right.$$

Therefore $1 - \alpha$ upper confidence interval on $\frac{\sigma_\epsilon^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2}$ is

$$P \left[\frac{\sigma_\epsilon^2}{\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2} \leq U \right] = 1 - \alpha,$$

where

$$(3.4.5) \quad U = \begin{cases} \min\{1, S_\alpha\} & \text{if } \frac{S_1^2}{S_2^2} > F_{\alpha; n_1, n_2} \\ \min\{1, T_\alpha\} & \text{if } \frac{S_1^2}{S_2^2} \leq F_{\alpha; n_1, n_2}, \end{cases}$$

$$\begin{aligned}
 S_\alpha &= \frac{JKMS_4^2}{\frac{S_2^2}{F_{\alpha; n_1, n_4}} \left[\frac{S_1^2}{S_2^2} + \frac{1}{2} \{ (J-1)F_{\alpha; n_1, \infty} + E_\alpha \} + G \right] + D}, \\
 G &= \frac{1}{2} F_{\alpha; n_1, n_2} \{ E_\alpha - (J-1)F_{\alpha; n_1, \infty} \} \frac{S_2^2}{S_1^2}, \\
 D &= J(K-1)S_3^2 + JK(M-1)S_4^2, \\
 E_\alpha &= \frac{J(n_1 F_{\alpha; n_1, n_2} + n_2) F_{\alpha; n_1, n_4}}{(n_1 + n_2) F_{\alpha; n_1 + n_2, n_4}} - F_{\alpha; n_1, n_2}, \\
 T_\alpha &= \frac{KMS_4^2}{\frac{S_2^2 \left(n_1 \frac{S_1^2}{S_2^2} + n_2 \right)}{(n_1 + n_2) F_{\alpha; n_1 + n_2, n_4}} + (K-1)S_3^2 + K(M-1)S_4^2}.
 \end{aligned}$$

We can choose another value,

$$(3.4.6) \quad \begin{cases} \text{replace } D \text{ with } J(K-1)S_3^2 F_{1-\alpha; n_1, n_3} \\ \quad + JK(M-1)S_4^2 F_{1-\alpha; n_1, n_4} \text{ of } S_{1-\alpha}, T_{1-\alpha} \text{ in } L, \\ \text{replace } D \text{ with } J(K-1)S_3^2 F_{\alpha; n_1, n_3} \\ \quad + JK(M-1)S_4^2 F_{\alpha; n_1, n_4} \text{ of } S_\alpha, T_\alpha \text{ in } U. \end{cases}$$

4. Discussion

In this chapter, I would like to discuss these precisions.

(1) The method of Graybill and Wang approximation was used in finding out the confidence interval on $\sigma_A^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ of (1.1). Graybill and Wang confidence intervals of the two-factor nested variance model, when $1 - \alpha = 0.95$, the precisions of the confidence intervals L and U lie between 0.950-0.966, 0.950-0.953 respectively. Accordingly the precisions of the above-mentioned intervals will be similar to those of this.

(2) The confidence intervals on $\sigma_B^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ and $\sigma_C^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ are similar to that of Graybill and Wang's confidence intervals on $\sigma_B^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2)$. Thus, the precisions of the confidence intervals on $\sigma_B^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ and $\sigma_C^2 / (\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ similar to that of this, when $1 - \alpha = 0.95$, the precisions of the confidence interval L and U lie between 0.947-0.986, 0.950-0.968 respectively.

(3) The precisions of the confidence intervals on $\sigma_\epsilon^2/(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ is similar to that of Graybill and Wang's precisions of the confidence intervals on $\sigma_C^2/(\sigma_A^2 + \sigma_B^2 + \sigma_C^2)$. Thus the precisions of the confidence intervals on $\sigma_\epsilon^2/(\sigma_A^2 + \sigma_B^2 + \sigma_C^2 + \sigma_\epsilon^2)$ is similar to that of od this, when $1 - \alpha = 0.95$, the precisions of the confidence interval L and U lie between 0.947-0.979, 0.946-0.957 respectively.

REFERENCES

1. Arteage, C , Jeyaratnam, S. and Graybill, F. A , *Confidence intervals for proportions of total variance in two-way cross component of variance model*, Communication in Statistics Theory and Methods 11 (1982), 1943-1958.
2. Burdik, R K and Graybill, F. A , *Confidence intervals on linear combinations of variance components in the unbalanced one-way classification*, Technometrics 26 (1984), 131-136
3. Burdik, R. K. and Graybill, F. A., *Confidence intervals on the total variance in an unbalanced two-fold nested classification with equal subsampling*, Communication in Statistics Theory and Methods 14(4) (1985), 761-774.
4. Graybill, F A , *Theory and application of the linear model*, North Scituate, Mass., Duxbury Press, 1976
5. Graybill, F. A and Wang, C. H., *Confidence intervals for proportion of variability in two-factor nested variance components models*, JASA 74 (1979), 368-374.
6. Howe, W G., *Approximate confidence limits on the mean of $X + Y$ where X and Y are two tabled independent random variables*, JASA 69 (1974), 789-794
7. Kang, Kwan-Joong, *Confidence intervals in three-factor nested variance component model*, J Korean Stat. Soc. 22(1) (1993), 39-54
8. Leiva, Ricardo A. and Graybill, F. A., *Confidence intervals for variance components in the balanced two-way model with interaction*, Communication in Statistics Simula 15(2) (1986), 301-322
9. Satterthwaite, F E , *An approximate distribution of estimates of variance components*, Biometrics Bulletin 2 (1946), 110-114.
10. Smith, H. Fairfield, *The problem of comparing the result of two experiments with unequal errors*, J Council of Scientific and Industrial Research 9 (1936), 211-212
11. Wang, C. and Graybill, F A., *Confidence intervals on a ratio of variances in the two-factor nested components of variance model*, Communication in Statistics Theory and Methods A 10(14) (1981), 1357-1368

Department of Mathematics
 Dong-A University
 Pusan 604-714, Korea