

## ON THE PETTIS DECOMPOSABILITY AND DUNFORD DECOMPOSABILITY

CHUN-KEE PARK

### 1. Introduction

Huff[7] studied Pettis integrability using the associated operator from a Banach space into  $L_1(\mu)$ . Bator[1] introduced the notion of a Pettis decomposable function from a finite measure space into a dual Banach space which is an extension of a Pettis integrable function.

In this paper, in order to generalize the Huff's results we study Pettis decomposability using the associated operator. Furthermore, we introduce the notion of a Dunford decomposable function which is an extension of a Pettis decomposable function, and obtain some properties of the Dunford decomposable function.

### 2. Preliminaries

Throughout this paper, let  $(\Omega, \Sigma, \mu)$  be a finite measure space and let  $X$  and  $Y$  be real Banach spaces with duals  $X^*$  and  $Y^*$ , respectively. We denote the closed unit ball of  $X$  by  $B_X$ . The adjoint of a continuous linear operator  $T$  from  $X$  to  $Y$  will be denoted by  $T^*$ .

If  $f : \Omega \rightarrow X^*$  is a Dunford integrable function, then we define  $T_f : X^{**} \rightarrow L_1(\mu)$  by  $T_f(x^{**}) = x^{**} \circ f$  for each  $x^{**} \in X^{**}$  and let  $\tilde{T}_f = T_f|_X$ , where  $T_f$  is called the associated operator with  $f$ . Note that if  $f : \Omega \rightarrow X^*$  is a Dunford equi-integrable function then  $\tilde{T}_f^{**}(X^{**}) \subseteq L_1(\mu)$ .

We use the following theorems in order to obtain our results.

---

Received December 21, 1993.

Supported by Kangwon National University Research Grant, 1993.

**THEOREM 2.1** [1]. *A weak\* scalarly bounded function  $f : \Omega \rightarrow X^*$  has the RS-property if and only if for every  $\epsilon > 0$  there exists  $E \in \Sigma$  with  $\mu(\Omega \setminus E) < \epsilon$  such that the set  $\{(x \circ f)\chi_E : x \in B_X\}$  is weakly pre-compact.*

**THEOREM 2.2** [9]. *If a bounded weakly measurable function  $f : \Omega \rightarrow X^*$  has the RS-property, then there exists a Pettis integrable function  $g : \Omega \rightarrow X^*$  such that  $x \circ f = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ .*

**THEOREM 2.3** [6]. *A bounded linear operator  $T : X \rightarrow Y$  is weakly compact if and only if  $T^* : Y^* \rightarrow X^*$  is  $(\omega^*, \omega)$ -continuous.*

**THEOREM 2.4** [7]. *A Dunford integrable function  $f : \Omega \rightarrow X$  is Pettis integrable if and only if the operator  $T_f : X^* \rightarrow L_1(\mu)$  defined by  $T_f(x^*) = x^* \circ f$  is  $(\omega^*, \omega)$ -continuous.*

*In particular, if  $f$  is Pettis integrable then  $T_f$  is a weakly compact operator.*

### 3. Pettis Decomposability

In this section, we investigate some properties of the Pettis decomposable function using the associated operator.

**DEFINITION 3.1** [1]. *A function  $f : \Omega \rightarrow X^*$  is said to be  $\mu$ -weak\* scalarly null if  $x \circ f = 0$   $\mu$ -a.e. for every  $x \in X$ .*

*A bounded weakly measurable function  $f : \Omega \rightarrow X^*$  is called  $\mu$ -Pettis decomposable if there exist a Pettis integrable function  $g$  and a  $\mu$ -weak\* scalarly null function  $h$  such that  $f = g + h$ .*

**THEOREM 3.1.** *If  $X$  is a separable Banach space, then a bounded weakly measurable function  $f : \Omega \rightarrow X^*$  is Pettis decomposable if and only if  $f$  is Pettis integrable.*

*proof.* If  $f : \Omega \rightarrow X^*$  is Pettis integrable, then clearly  $f$  is Pettis decomposable.

Conversely, if  $f : \Omega \rightarrow X^*$  is Pettis decomposable, then there exist a Pettis integrable function  $g$  and a  $\mu$ -weak\* scalarly null function  $h$  such that  $f = g + h$ . Let  $(x_i)_{i=1}^\infty$  be dense in  $X$ . Since  $h$  is  $\mu$ -weak\* scalarly null,  $x \circ h = 0$   $\mu$ -a.e. for every  $x \in X$ . For each  $i$ , let  $N_i$  be  $\mu$ -null sets

such that  $x_i \circ h(t) = 0$  if  $t \in \Omega \setminus N_i$ . Then for each  $i$ ,  $x_i \circ h(t) = 0$  if  $t \in \Omega \setminus (\cup_{i=1}^{\infty} N_i)$ . Let  $x \in X$  be arbitrary. Since  $(x_i)_{i=1}^{\infty}$  is dense in  $X$ , there exists a subsequence  $(x_{i_j})_{j=1}^{\infty}$  of  $(x_i)_{i=1}^{\infty}$  such that  $\lim_{j \rightarrow \infty} x_{i_j} = x$ . Hence  $x \circ h(t) = (\lim_{j \rightarrow \infty} x_{i_j}) \circ h(t) = \lim_{j \rightarrow \infty} (x_{i_j} \circ h(t)) = 0$  if  $t \in \Omega \setminus (\cup_{i=1}^{\infty} N_i)$ . Therefore  $h(t) = 0$  if  $t \in \Omega \setminus (\cup_{i=1}^{\infty} N_i)$ . Since  $\mu(\cup_{i=1}^{\infty} N_i) = 0$ ,  $h = 0$   $\mu$ -a.e. Thus  $f = g$   $\mu$ -a.e. Since  $g$  is Pettis integrable,  $f$  is also Pettis integrable.

**THEOREM 3.2.** *Let  $f : \Omega \rightarrow X^*$  be a bounded weakly measurable function. Then  $f : \Omega \rightarrow X^*$  is Pettis decomposable if and only if the operator  $\tilde{T}_f^{**} : X^{**} \rightarrow L_1(\mu)$  is  $(\omega^*, \omega)$ -continuous.*

*proof.* Suppose that  $f : \Omega \rightarrow X^*$  is Pettis decomposable. Then there exist a Pettis integrable function  $g$  and a  $\mu$ -weak\* scalarly null function  $h$  such that  $f = g + h$ . Since  $g$  is Pettis integrable, the operator  $T_g : X^{**} \rightarrow L_1(\mu)$  defined by  $T_g(x^{**}) = x^{**} \circ g$  is  $(\omega^*, \omega)$ -continuous by Theorem 2.4. Hence  $\tilde{T}_g^{**} = T_g$  ([4], p685). Since  $h$  is  $\mu$ -weak\* scalarly null,  $x \circ f = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ . Hence  $\tilde{T}_f = \tilde{T}_g$ , and so  $\tilde{T}_f^{**} = \tilde{T}_g^{**}$ . Thus  $\tilde{T}_f^{**} = T_g$ . Therefore  $\tilde{T}_f^{**}$  is  $(\omega^*, \omega)$ -continuous.

Conversely, suppose that the operator  $\tilde{T}_f^{**} : X^{**} \rightarrow L_1(\mu)$  is  $(\omega^*, \omega)$ -continuous. Then  $\tilde{T}_f^{**}$  is weakly compact by Theorem 2.3. Hence  $\tilde{T}_f$  is also weakly compact, and so  $\tilde{T}_f(B_X) = \{\tilde{T}_f(x) : x \in B_X\} = \{x \circ f : x \in B_X\}$  is weakly sequentially compact. Therefore every sequence in  $\{x \circ f : x \in B_X\}$  has a weakly convergent subsequence. Hence  $\{x \circ f : x \in B_X\}$  is weakly pre-compact. By Theorem 2.1,  $f : \Omega \rightarrow X^*$  has the RS-property. By Theorem 2.2, there exists a Pettis integrable function  $g : \Omega \rightarrow X^*$  such that  $x \circ f = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ . Let  $h = f - g$ , then  $h$  is  $\mu$ -weak\* scalarly null. Therefore  $f = g + h$  is Pettis decomposable.

**THEOREM 3.3.** *Let  $f : \Omega \rightarrow X^*$  be a bounded weakly measurable function. Then  $f : \Omega \rightarrow X^*$  is Pettis integrable if and only if  $T_f = \tilde{T}_f^{**}$ .*

*proof.* Suppose that  $f : \Omega \rightarrow X^*$  is Pettis integrable. Then  $T_f : X^{**} \rightarrow L_1(\mu)$  defined by  $T_f(x^{**}) = x^{**} \circ f$  is  $(\omega^*, \omega)$ -continuous by Theorem 2.4. Let  $x^{**} \in X^{**}$  be arbitrary. Since  $X$  is weak\* dense in  $X^{**}$ , there exists a net  $(x_\alpha)$  in  $X$  such that  $x_\alpha \xrightarrow{\omega^*} x^{**}$ .  $T_f(x_\alpha) = \tilde{T}_f(x_\alpha) = \tilde{T}_f^{**}(x_\alpha)$  for each  $\alpha$ . Since  $T_f$  is  $(\omega^*, \omega)$ -continuous,

$T_f(x_\alpha) \xrightarrow{\omega} T_f(x^{**})$ . Since  $f$  is Pettis integrable,  $T_f$  is weakly compact by Theorem 2.4. Hence  $\tilde{T}_f$  is also weakly compact, and so  $\tilde{T}_f^*$  is weakly compact. By Theorem 2.3,  $\tilde{T}_f^{**}$  is  $(\omega^*, \omega)$ -continuous. Hence  $\tilde{T}_f^{**}(x_\alpha) \xrightarrow{\omega} \tilde{T}_f^{**}(x^{**})$ . Therefore  $T_f(x^{**}) = \tilde{T}_f^{**}(x^{**})$  for all  $x^{**} \in X^{**}$ . Thus  $T_f = \tilde{T}_f^{**}$ .

Conversely, suppose that  $T_f = \tilde{T}_f^{**}$ . Then  $\tilde{T}_f$  is weakly compact since  $\tilde{T}_f^{**}(X^{**}) \subseteq L_1(\mu)$ . Hence  $\tilde{T}_f^*$  is also weakly compact, and so  $\tilde{T}_f^{**}$  is  $(\omega^*, \omega)$ -continuous by Theorem 2.3. By hypothesis,  $T_f$  is  $(\omega^*, \omega)$ -continuous. Therefore  $f$  is Pettis integrable by Theorem 2.4.

#### 4. Dunford Decomposability

In this section, we introduce the new concept of a Dunford decomposable function which is an extension of Pettis decomposable function and investigate properties of the Dunford decomposable function.

**DEFINITION 4.1.** A Dunford integrable function  $f : \Omega \rightarrow X^*$  is called  $\mu$ -Dunford decomposable if there exist a Dunford equi-integrable function  $g$  and a  $\mu$ -weak\* scalarly null function  $h$  such that  $f = g + h$ .

**LEMMA 4.1.** Let  $f : \Omega \rightarrow X$  be a Dunford integrable function. Then  $f : \Omega \rightarrow X$  is Dunford equi-integrable if and only if the operator  $T_f : X^* \rightarrow L_1(\mu)$  defined by  $T_f(x^*) = x^* \circ f$  is weakly compact.

*proof.*  $f : \Omega \rightarrow X$  is Dunford equi-integrable if and only if  $\{x^* \circ f : x^* \in B_{X^*}\}$  is uniformly integrable in  $L_1(\mu)$ .  $T_f(B_{X^*}) = \{x^* \circ f : x^* \in B_{X^*}\}$  is uniformly integrable in  $L_1(\mu)$  if and only if  $T_f(B_{X^*})$  is relatively weakly compact in  $L_1(\mu)$ . Thus  $f$  is Dunford equi-integrable if and only if  $T_f$  is weakly compact.

**THEOREM 4.2.** If  $f : \Omega \rightarrow X^*$  is Dunford decomposable, then  $\tilde{T}_f^{**} : X^{**} \rightarrow L_1(\mu)^{**}$  is weakly compact.

*proof.* If  $f : \Omega \rightarrow X^*$  is Dunford decomposable, then there exist a Dunford equi-integrable function  $g$  and a  $\mu$ -weak\* scalarly null function  $h$  such that  $f = g + h$ . Since  $h$  is  $\mu$ -weak\* scalarly null,  $x \circ f = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ . Hence  $\tilde{T}_f = \tilde{T}_g$ , and so

$\tilde{T}_f^{**} = \tilde{T}_g^{**}$ . Since  $g : \Omega \rightarrow X^*$  is Dunford equi-integrable,  $T_g : X^{**} \rightarrow L_1(\mu)$  defined by  $T_g(x^{**}) = x^{**} \circ g$  is weakly compact by Lemma 4.1. Hence  $\tilde{T}_g : X \rightarrow L_1(\mu)$  is weakly compact, and so  $\tilde{T}_g^{**}$  is also weakly compact. Therefore  $\tilde{T}_f^{**}$  is weakly compact.

**THEOREM 4.3.** *If  $f : \Omega \rightarrow X^*$  is Dunford integrable then the following are equivalent:*

- (i)  $f : \Omega \rightarrow X^*$  is Dunford decomposable.
- (ii) There exists a Dunford equi-integrable function  $g$  such that for every  $x \in X$ ,  $T_f(x) = x \circ g$  in  $L_1(\mu)$ .
- (iii) For every  $\epsilon > 0$ , there exist  $E \in \Sigma$  and a Dunford equi-integrable function  $g$  such that  $\mu(\Omega \setminus E) < \epsilon$  and  $(x \circ f)\chi_E = x \circ g$   $\mu - a.e.$  for every  $x \in X$ .

*proof.* (i) $\Rightarrow$ (ii). If  $f : \Omega \rightarrow X^*$  is Dunford decomposable, then there exist a Dunford equi-integrable function  $g$  and a  $\mu - weak^*$  scalarly null function  $h$  such that  $f = g + h$ . Since  $h$  is  $\mu - weak^*$  scalarly null,  $x \circ f = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ . Therefore  $T_f(x) = x \circ g$  in  $L_1(\mu)$  for every  $x \in X$ .

(ii) $\Rightarrow$ (iii). Suppose that (ii) holds. Then there exists a Dunford equi-integrable function  $g$  such that  $x \circ f = x \circ g$   $\mu - a.e.$  for every  $x \in X$ . Let  $E = \Omega$ . Then (iii) holds.

(iii) $\Rightarrow$ (i). Suppose that (iii) holds. By the exhaustion principle ([5], p70), there exists a Dunford equi-integrable function  $g; \Omega \rightarrow X^*$  such that  $x \circ f = x \circ g$   $\mu - a.e.$  for every  $x \in X$ . Let  $h = f - g$ . Then  $h$  is  $\mu - weak^*$  scalarly null. Hence  $f = g + h$  is Dunford decomposable.

### References

- [1] E.M. Bator, *A decomposition of bounded scalarly measurable functions taking their ranges in dual Banach spaces*, Proc. Amer. Math. Soc. **102** (1988), 850-854.
- [2] E.M. Bator, *Pettis integrability and the equality of the norms of the weak\* integral and the Dunford integral*, Proc. Amer. Math. Soc. **95** (1985), 265-270.
- [3] E.M. Bator, *Pettis decomposition for universally scalarly measurable functions*, Proc. Amer. Math. Soc. **104** (1988), 795-800.

- [4] E. M. Bator, P. W. Lewis and D. Race, *Some connections between Pettis integration and operator theory*, Rocky Mountain Math. J. **17** (1987), 683–695.
- [5] J. Diestel and J. J. Uhl, Jr., *Vector measures*, *Math. Surveys*, vol. 15, Amer. Math. Soc., Providence, R.I., 1977.
- [6] N. Dunford and J. T. Schwarz, *Linear operators, Part I*, Interscience, New York, 1958.
- [7] R. E. Huff, Remarks on Pettis integrability, Proc. Amer. Math. Soc. **96** (1986), 402–404.
- [8] L. H. Riddle, E. Saab and J. J. Uhl, Jr., *Sets with the weak Radon-Nikodym property in dual Banach spaces*, Indiana Univ. Math. J. **32** (1983), 527–541.
- [9] M. Talagrand, *Pettis integral and measure theory*, Mem. Amer. Math. Soc. Vol. 51, no. 307, 1984.

Chun-Kee Park  
Department of Mathematics  
Kangwon National University  
Chuncheon 200-701, Korea