

## ON WEAK-MIXING PROPERTY

YONG SUN CHO AND HYUN WOO LEE

### 1. Introduction.

Let  $(X, \mathcal{B}, m)$  be a probability space and let  $T : X \rightarrow X$  a measure-preserving transformation. We introduce some results related ergodicity and mixing property of  $T$ . We try to extend 1-fold ergodicity and 1-fold mixing to  $r$ -fold ergodicity and  $r$ -fold mixing. We shall use the result, about sequences of real numbers, to obtain formulations of  $r$ -fold weak-mixing. We show that for a measure-preserving transformation  $T$ ,  $r$ -fold weak-mixing properties of  $T$  are equivalent to  $r$ -fold strong-mixing properties of  $T \times T$ . Now, we shall introduce the concept of  $r$ -fold weak-mixing property and  $r$ -fold strong-mixing property. We show that basic properties of 1-fold mixing keep same properties in the case of  $r$ -fold mixing. We shall use that  $U$  is defined on functions by  $Uf = f \circ T, \forall f \in L^p(X, \mathcal{B}, m), p \geq 1$ .

### 2. $r$ -fold mixing.

DEFINITION 1. Let  $T$  be a measure-preserving transformation of a probability space  $(X, \mathcal{B}, m)$  and let  $r \geq 1$ .

(a)  $T$  is  $r$ -fold ergodic if  $\forall A_0, A_1, \dots, A_r \in \mathcal{B}$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n m(A_0 \cap T^{-i} A_1 \cap \dots \cap T^{-ri} A_r) = m(A_0)m(A_1) \cdots m(A_r).$$

(b)  $T$  is  $r$ -fold weak-mixing if  $\forall A_0, A_1, \dots, A_r \in \mathcal{B}$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n |m(A_0 \cap T^{-i} A_1 \cap \dots \cap T^{-ri} A_r) \\ - m(A_0)m(A_1) \cdots m(A_r)| \\ = 0. \end{aligned}$$

---

Received November 2, 1994.

(c)  $T$  is  $r$ -fold strong-mixing if  $\forall A_0, A_1, \dots, A_r \in \mathcal{B}$ ,

$$\lim_{n_1, \dots, n_r \rightarrow \infty} m(A_0 \cap T^{-n_1} A_1 \cap \dots \cap T^{-(n_1 + \dots + n_r)} A_r) = \prod_{j=0}^r m(A_j).$$

**THEOREM 1.** Let  $T : (X, \mathcal{B}, m) \rightarrow (X, \mathcal{B}, m)$  be a measure-preserving transformation.

(i) The following are equivalent :

- (1)  $T$  is  $r$ -fold ergodic.
- (2) For all  $f_0, f_1, \dots, f_r \in L^{r+1}(X, \mathcal{B}, m)$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \int (f_0 U^i f_1 \dots U^{ri} f_r) dm = \prod_{j=0}^r \int f_j dm.$$

(ii) The following are equivalent :

- (1)  $T$  is  $r$ -fold weak-mixing.
- (2) For all  $f_0, f_1, \dots, f_r \in L^{r+1}(X, \mathcal{B}, m)$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \left| \int (f_0 U^i f_1 \dots U^{ri} f_r) dm - \prod_{j=0}^r \int f_j dm \right| = 0.$$

(iii) The following are equivalent :

- (1)  $T$  is  $r$ -fold strong-mixing.
- (2) For all  $f_0, f_1, \dots, f_r \in L^{r+1}(X, \mathcal{B}, m)$ ,

$$\lim_{n_1, \dots, n_r \rightarrow \infty} \int (f_0 U^{n_1} f_1 \dots U^{n_1 + \dots + n_r} f_r) dm = \prod_{j=0}^r \int f_j dm.$$

*Proof.* (i),(ii) and (iii) are proved by using similar methods of [4].

**REMARK.** Every  $r$ -fold strong-mixing transformation implies  $r$ -fold weak-mixing and every  $r$ -fold weak-mixing transformation implies  $r$ -fold ergodic.

This is because if  $\{a_n\}$  is a sequence of real numbers then

$$\lim_{n \rightarrow \infty} a_n = 0$$

implies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} |a_i| = 0$$

and this second condition implies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} a_i = 0.$$

We shall use the following result, about sequences of real numbers, to obtain other formulations of  $r$ -fold weak-mixing.

**THEOREM 2.** *If  $\{a_n\}$  is a bounded sequence of real numbers then the following are equivalent :*

(i)

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} |a_i| = 0.$$

(ii) *There exists a subset  $J$  of  $\mathbb{Z}^+$  of density zero*

$$\left( \text{i.e., } \left( \frac{\text{cardinality}(J \cap \{0, 1, \dots, n-1\})}{n} \right) \rightarrow 0 \right),$$

*such that  $\lim_{n \rightarrow \infty} a_n = 0$  provided  $n \notin J$ .*

(iii)

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} |a_i|^2 = 0.$$

*Proof.* See [7].

**THEOREM 3.** *If  $T$  is a measure-preserving transformation of a probability space  $(X, \mathcal{B}, m)$ , the followings are equivalent:*

- (i)  $T$  is weak-mixing.  
(ii) For all  $A_0, A_1, \dots, A_r$  of  $\mathcal{B}$  there is a subset  $J(A_0, A_1, \dots, A_r)$  of  $\mathbb{Z}^+$  of density zero such that

$$\lim_{\substack{n \notin J(A_0, A_1, \dots, A_r) \\ n \rightarrow \infty}} m(A_0 \cap T^{-n} A_1 \cap \dots \cap T^{-rn} A_r) = m(A_0)m(A_1) \cdots m(A_r).$$

- (iii) For all  $A_0, A_1, \dots, A_r$  of  $\mathcal{B}$  we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n |m(A_0 \cap T^{-i} A_1 \cap \dots \cap T^{-ri} A_r) \\ - m(A_0)m(A_1) \cdots m(A_r)|^2 \\ = 0. \end{aligned}$$

*Proof.* Apply Theorem 2 with

$$a_n = m(A_0 \cap T^{-n} A_1 \cap T^{-2n} A_2 \cap \dots \cap T^{-rn} A_r) - m(A_0)m(A_1) \cdots m(A_r).$$

The following theorem gives a way of checking the mixing properties for examples by reducing the computations to a class of sets we can manipulate with.

**THEOREM 4.** *Let  $(X, \mathcal{B}, m)$  be a probability space and let  $\mathcal{T}$  be a semi-algebra that generates  $\mathcal{B}$ . Let  $T : X \rightarrow X$  be a measure-preserving transformation. Then  $T$  is  $r$ -fold strang-mixing iff*

$$\forall C_0, C_1, \dots, C_r \in \mathcal{T},$$

$$\begin{aligned} \lim_{n_1, \dots, n_r \rightarrow \infty} m(C_0 \cap T^{-n_1} C_1 \cap \dots \cap T^{-(n_1 + \dots + n_r)} C_r) \\ = m(C_0)m(C_1) \cdots m(C_r). \end{aligned}$$

*Proof.* See [4].

### 3. $r$ -fold weak-mixing.

We have main result. This result connects  $r$ -fold weak-mixing of  $T$  with  $r$ -fold ergodicity of  $T \times T$ . Also, this result relates to  $r$ -fold weak-mixing of  $T$  with  $r$ -fold weak-mixing of  $T \times T$ .

**THEOREM 5.** *If  $T$  is a measure-preserving transformation on a probability space  $(X, \mathcal{B}, m)$  then the followings are equivalent:*

- (i)  $T$  is  $r$ -fold weak-mixing .
- (ii)  $T \times T$  is  $r$ -fold ergodic .
- (iii)  $T \times T$  is  $r$ -fold weak-mixing .

*Proof.* ((i)  $\Rightarrow$  (iii)). Let  $A_0, A_1, \dots, A_r \in \mathcal{B}$  ,  $B_0, B_1, \dots, B_r \in \mathcal{B}$ . There exist subsets  $J_1(A_0, A_1, \dots, A_r)$ ,  $J_2(B_0, B_1, \dots, B_r)$  of  $\mathbb{Z}^+$  of density zero such that

$$\lim_{\substack{n \notin J_1 \\ n \rightarrow \infty}} m(A_0 \cap T^{-n}A_1 \cap \dots \cap T^{-rn}A_r) = m(A_0)m(A_1) \cdots m(A_r)$$

and

$$\lim_{\substack{n \notin J_2 \\ n \rightarrow \infty}} m(B_0 \cap T^{-n}B_1 \cap \dots \cap T^{-rn}B_r) = m(B_0)m(B_1) \cdots m(B_r).$$

Then

$$\begin{aligned} & \lim_{\substack{n \notin J_1 \cup J_2 \\ n \rightarrow \infty}} (m \times m)\{(A_0 \times B_0) \cap (T \times T)^{-n}(A_1 \times B_1) \cap \dots \\ & \quad \cap (T \times T)^{-rn}(A_r \times B_r)\} \\ &= \lim_{\substack{n \notin J_1 \cup J_2 \\ n \rightarrow \infty}} m(A_0 \cap T^{-n}A_1 \cap \dots \cap T^{-rn}A_r) \cdot \\ & \quad m(B_0 \cap T^{-n}B_1 \cap \dots \cap T^{-rn}B_r) \\ &= m(A_0)m(A_1) \cdots m(A_r)m(B_0)m(B_1) \cdots m(B_r) \\ &= (m \times m)(A_0 \times B_0)(m \times m)(A_1 \times B_1) \cdots (m \times m)(A_r \times B_r). \end{aligned}$$

By Theorem 2 we know

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n |(m \times m)[(A_0 \times B_0) \cap (T \times T)^{-i}(A_1 \times B_1) \cap \cdots \\ & \quad \cap (T \times T)^{-ri}(A_r \times B_r)] - \prod_{j=0}^r (m \times m)(A_j \times B_j)| \\ & = 0. \end{aligned}$$

Since the measurable rectangles form a semi-algebra that generates  $\mathcal{B} \times \mathcal{B}$ , Theorem 4 asserts that  $T \times T$  is  $r$ -fold weak-mixing.

((iii)  $\Rightarrow$  (ii)). It is clear that (iii) implies (ii).

((ii)  $\Rightarrow$  (i)). Let  $A_0, A_1, \dots, A_r \in \mathcal{B}$ . We shall show

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \{m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r) - m(A_0)m(A_1) \cdots m(A_r)\}^2 = 0.$$

We have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r) \\ & = \frac{1}{n} \sum_{i=1}^n (m \times m)\{(A_0 \times X) \cap (T \times T)^{-i}(A_1 \times X) \cap \cdots \\ & \quad \cap (T \times T)^{-ri}(A_r \times X)\} \\ & \rightarrow (m \times m)(A_0 \times X)(m \times m)(A_1 \times X) \cdots (m \times m)(A_r \times X) \\ & = m(A_0)m(A_1) \cdots m(A_r). \end{aligned}$$

Also

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \{m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r)\}^2 \\
&= \frac{1}{n} \sum_{i=1}^n (m \times m) \{(A_0 \times A_0) \cap (T \times T)^{-i}(A_1 \times A_1) \cap \cdots \\
&\quad \cap (T \times T)^{-ri}(A_r \times A_r)\} \\
&\quad \rightarrow (m \times m)(A_0 \times A_0)(m \times m)(A_1 \times A_1) \cdots \\
&\quad (m \times m)(A_r \times A_r) \\
&\quad = m(A_0)^2 m(A_1)^2 \cdots m(A_r)^2.
\end{aligned}$$

Thus

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \{m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r) - m(A_0)m(A_1) \cdots m(A_r)\}^2 \\
&= \frac{1}{n} \sum_{i=1}^n \{(m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r))^2 \\
&\quad - 2m(A_0 \cap T^{-i}A_1 \cap \cdots \cap T^{-ri}A_r)m(A_0)m(A_1) \cdots m(A_r) \\
&\quad + m(A_0)^2 m(A_1)^2 \cdots m(A_r)^2\} \\
&\quad \rightarrow 2m(A_0)^2 m(A_1)^2 \cdots m(A_r)^2 - 2m(A_0)^2 m(A_1)^2 \cdots m(A_r)^2 \\
&\quad = 0.
\end{aligned}$$

Therefore  $T$  is  $r$ -fold weak-mixing by Theorem 3.

### References

- [1] Bedford, T., Keane, M. and Series, C., *Ergodic Theory, Symbolic Dynamics and Hyperbolic Spaces*, Oxford University press, 1991.
- [2] Bratteli, O. and Robinson, D. W., *Operator Algebras and Quantum Statistical Mechanics I*, Springer-Verlag, 1979.
- [3] Cornfeld, I. P., Fomin, S. V. and Sinai, Ya. G., *Ergodic Theory*, Springer-Verlag, 1982.
- [4] Inhae, N. and Hyunwoo, L., *A Note on Mixing Property*, J. of Basic Sci., Sungshin Women's Univ. **12** (1994.).
- [5] Mane, R., *Ergodic Theory, and Differentiable Dynamics*, Springer-Verlag, 1987.

- [6] Sinai, Ya. G., *Dynamical Systems II, Encyclopedia of Mathematical Sciences*, vol. 2, Springer-Verlag, 1989.
- [7] Walters, P., *An Introduction to Ergodic Theory*, Springer-Verlag, 1982.

Department of Mathematics  
Sungshin Women's University  
Seoul, 136-742, Korea