

ON FUZZY s -CONTINUOUS FUNCTIONS

WON KEUN MIN

ABSTRACT. We introduce the concepts of fuzzy s -continuous functions. And we investigate several properties of the fuzzy s -continuous function. In particular, we study the relation between fuzzy continuous functions and fuzzy s -continuous functions.

1. INTRODUCTION

Fuzzy topological spaces were first introduced in the literature by Chang [1] who studied a number of the basic concepts including fuzzy continuous maps and compactness. And fuzzy topological spaces are a very natural generalization of topological spaces. In 1983, A.S. Mashhour et al.[3] introduced supra topological spaces and studied s -continuous functions and s^* -continuous functions. In 1987, M.E. Abd El-Monsef et al.[2] introduced the fuzzy supra topological spaces and studied fuzzy supra-continuous functions and characterized a number of basic concepts. Also fuzzy supra topological spaces are generalizations of supra topological spaces. In this paper, we introduce fuzzy s -continuous function and establish a number of characterizations. Let X be a set and let $I = [0, 1]$. Let I^X denote the set of all mapping $a: X \rightarrow I$.

A member of I^X is called a fuzzy subset of X . And unions and intersections of fuzzy sets are denoted by \vee and \wedge respectively and defined by

$$\begin{aligned}\vee a_i &= \sup\{a_i(x) \mid i \in J \text{ and } x \in X\}, \\ \wedge a_i &= \inf\{a_i(x) \mid i \in J \text{ and } x \in X\}.\end{aligned}$$

Received December 26, 1995.

1991 *Mathematics Subject Classification.* 54A40.

Key words and phrases. fuzzy s -continuous, fuzzy continuous, fuzzy supra topological space, fuzzy supracontinuous.

DEFINITION 1.1[1]. A fuzzy topology T on X is a collection of subsets of I^X such that

- (1) $0, 1 \in T$,
- (2) if $a, b \in T$, then $a \wedge b \in T$,
- (3) if $a_i \in T$ for all $i \in J$, then $\bigvee a_i \in T$.

(X, T) is called a fuzzy topological space. Members of T are called fuzzy open sets in (X, T) and complement of a fuzzy open set is called a fuzzy closed set.

DEFINITION 1.2. Let f be a mapping from a set X into a set Y . Let a and b be the fuzzy sets of X and Y , respectively. Then $f(a)$ is a fuzzy set in Y , defined by

$$f(a)(y) = \begin{cases} \sup_{z \in f^{-1}(y)} a(z), & \text{if } f^{-1}(y) \neq \emptyset, y \in Y \\ 0, & \text{otherwise,} \end{cases}$$

and $f^{-1}(b)$ is a fuzzy set in X , defined by $f^{-1}(b)(x) = b(f(x))$, $x \in X$.

DEFINITION 1.3[3]. A subfamily T^* of I^X is said to be a fuzzy supra-topology on X if

- (1) $1 \in T^*$,
- (2) if $a_i \in T^*$ for all $i \in J$, then $\bigvee a_i \in T^*$.

(X, T^*) is called a fuzzy supra topological space. The elements of T^* are called fuzzy supra open sets in (X, T^*) . And a fuzzy set a is supra closed iff $co(a) = 1 - a$ is a fuzzy supra open set. And the fuzzy supra topological spaces T^* is denoted by fsts.

DEFINITION 1.4[3]. The supra closure of a fuzzy set a is denoted by $scl(a)$, and given by

$$scl(a) = \bigwedge \{s \mid s \text{ is a fuzzy supra closed set and } a \leq s\}.$$

The supra interior of a fuzzy set a is denoted by $si(a)$ and given by

$$si(a) = \bigvee \{t \mid t \text{ is a fuzzy supra open set and } t \leq a\}.$$

2. Fuzzy s -continuous function

DEFINITION 2.1[2]. Let (X, T) be a fuzzy topological space and T^* be a fuzzy supra-topology on X . We call T^* a fuzzy supra-topology associated with T if $T \subset T^*$.

DEFINITION 2.2[2]. Let $f: (X, T^*) \rightarrow (Y, S^*)$ be a mapping between two fuzzy supratopological spaces. f is a fuzzy supracontinuous function if $f^{-1}(S^*) \subseteq T^*$.

DEFINITION 2.3. Let (X, T) and (Y, S) be fuzzy topological spaces and T^* be an associated fuzzy supra-topology with T . A function $f: X \rightarrow Y$ is a fuzzy s -continuous function if the inverse image of each fuzzy open set in Y is T^* -fuzzy supra open in X .

THEOREM 2.4. Let (X, T) and (Y, S) be fts. Let f be a function from X into Y . Let T^* be an associated fuzzy supra-topology with T . Then the followings are equivalent :

- (1) f is fuzzy s -continuous.
- (2) The inverse image of each fuzzy closed set in Y is T^* -fuzzy supra closed.
- (3) $scl(f^{-1}(a)) \leq f^{-1}(cl(a))$ for every fuzzy set a in Y .
- (4) $f(scl(a)) \leq cl(f(a))$ for every fuzzy set a in X .
- (5) $f^{-1}(int(b)) \leq si(f^{-1}(b))$ for every fuzzy set b in Y .
- (6) For each fuzzy set a in X and each fuzzy neighborhood b of $f(a)$, there is a fuzzy supra neighborhood c of a such that $f(c) \leq b$.

PROOF. (1) \Rightarrow (2). Let a be fuzzy closed set in Y . Since f is a fuzzy s -continuous, $f^{-1}(1 - a) = 1 - f^{-1}(a)$ is fuzzy supra open in X . Therefore $f^{-1}(a)$ is a fuzzy supra closed set in X .

(2) \Rightarrow (3). Since $cl(a)$ is fuzzy closed for every fuzzy set a in Y , $f^{-1}(cl(a))$ is T^* -fuzzy supra closed. Therefore,

$$f^{-1}(cl(a)) = scl(f^{-1}(cl(a))) \geq scl(f^{-1}(a)).$$

(3) \Rightarrow (4). Let a be fuzzy subset in X and let $f(a) = b$. Then $f^{-1}(scl(b)) \geq scl(f^{-1}(b))$. So $f^{-1}(cl(f(a))) \geq scl(f^{-1}f(a)) \geq scl(a)$, and hence $cl(f(a)) \geq f(scl(a))$.

(4) \Rightarrow (2). Let b be a fuzzy closed set in Y and be $a = f^{-1}(b)$. Then $f(scl(a)) \leq cl(f(a)) = cl(f(f^{-1}(b))) \leq cl(b) = b$. Since $scl(a) \leq f^{-1}(f(scl(a))) \leq f^{-1}(b) = a$, then a is T^* -fuzzy supra closed.

(2) \Rightarrow (1). Obvious.

(1) \Rightarrow (5). Let b be a fuzzy subset in Y . Since $f^{-1}(int(b))$ is T^* -fuzzy supra open set in X , $f^{-1}(int(b)) \leq si(f^{-1}(int(b))) \leq si(f^{-1}(b))$.

(5) \Rightarrow (1). Let a be a fuzzy open set in Y . Since $f^{-1}(a) \leq si(f^{-1}(a)) \leq f^{-1}(a)$, $f^{-1}(a)$ is T^* -fuzzy supra open.

(6) \Rightarrow (1). Let b be any fuzzy open set in Y and let $f^{-1}(b) = a$. Then b is a fuzzy neighborhood of $f(a) = f(f^{-1}(b))$. There exists a fuzzy supra neighborhood c of $a = f^{-1}(b)$ such that $f(c) \leq b$. Thus $c \leq f^{-1}f(c) \leq f^{-1}(b)$. Therefore, $f^{-1}(b)$ is a fuzzy supra neighborhood of $f^{-1}(b)$. And $f^{-1}(b)$ is a fuzzy supra open set in X , by [6, Theorem 2.2].

(1) \Rightarrow (6). Obvious. \square

REMARK. Every fuzzy continuous function is fuzzy s -continuous. But the converse of this implication is not true, as following example shows.

EXAMPLE 2.1. Let a_1, a_2 , and a_3 be fuzzy subsets of $X = I$, defined as

$$a_1(x) = \begin{cases} 0, & \text{if } 0 \leq x \leq 1/2, \\ 2x - 1, & \text{if } 1/2 \leq x \leq 1; \end{cases}$$

$$a_2(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq 1/4, \\ -4x + 2, & \text{if } 1/4 \leq x \leq 1/2, \\ 0, & \text{if } 1/2 \leq x \leq 1; \end{cases}$$

$$a_3(x) = \begin{cases} 1, & \text{if } \leq x \leq 1/2, \\ -2x + 2, & \text{if } 1/2 \leq x \leq 1. \end{cases}$$

Consider the fuzzy space $T_1 = \{0, a_1, a_2, a_1 \vee a_2, 1\}$ and an associated supra fuzzy space $T_1^* = \{0, a_1, a_2, a_3, a_1 \vee a_2, a_1 \vee a_3, 1\}$. Let $g: X \rightarrow X$ be defined by $g(x) = (1/2)x$. Clearly, we have $g^{-1}(0) = 0$, $g^{-1}(1) = 1$, $g^{-1}(a_1 \vee a_2) = a_3$, $g^{-1}(a_2) = a_3$, and $g^{-1}(a_1) = 0$. $co(a_1) = a_3$ is a fuzzy supra open in (X, T_1^*) but it is not fuzzy open in (X, T_1) . Hence the fuzzy mapping g is fuzzy s -continuous but not fuzzy continuous.

REMARK. In general, the composition of two fuzzy s -continuous functions need not be fuzzy s -continuous.

EXAMPLE 2.2. Let $X = I$. Consider the fuzzy sets

$$a(x) = \begin{cases} 1, & \text{if } 0 \leq x < 1/3 \\ 1/2, & \text{if } 1/3 \leq x \leq 2/3 \\ 0, & \text{if } 2/3 < x \leq 1, \end{cases}$$

$$b(x) = 1/2, \text{ if } 0 \leq x \leq 1,$$

$$c(x) = 1/3, \text{ if } 0 \leq x \leq 1.$$

Let $T_1 = \{0, a, 1\}$ and $T_1^* = \{0, a, b, a \vee b, 1\}$. Let $T_2 = \{0, c, 1\}$ and $T_2^* = \{0, a, c, a \vee c, 1\}$. Let $f: (X, T_1) \rightarrow (X, T_1)$ be a fuzzy mapping defined by $f(x) = (x + 1)/3$. Let $g: (X, T_2) \rightarrow (X, T_1)$ be a fuzzy mapping defined by $g(x) = (1/3)x$. Clearly, f and g are fuzzy s -continuous. But $(f \circ g)$ is not fuzzy s -continuous, since a is a fuzzy open set in (X, T_1) but $(f \circ g)^{-1}(a) = b$ is not fuzzy supra open in T_2^* .

THEOREM 2.5. If a fuzzy mapping $f: (X, T_1) \rightarrow (Y, T_2)$ is fuzzy s -continuous and $g: (Y, T_2) \rightarrow (Z, T_3)$ is fuzzy continuous, then $(g \circ f)$ is fuzzy s -continuous.

PROOF. The proof is clear by the definitions of fuzzy s -continuous functions and fuzzy continuous functions. \square

THEOREM 2.6. Let (X, T) and (Y, S) be fts, T^* and S^* be two associated fuzzy supra-topologies with T and S , respectively. If $f: X \rightarrow Y$ is a fuzzy mapping, and one of the followings;

- (1) $f^{-1}(si(a)) \leq int(f^{-1}(a))$ for each fuzzy set a in (Y, S) ,
- (2) $cl(f^{-1}(a)) \leq f^{-1}(scl(a))$ for each fuzzy set a in (Y, S) ,
- (3) $f(cl(b)) \leq scl(f(b))$ for each fuzzy set b in (X, T) , holds, then f is fuzzy continuous.

PROOF. If the condition (2) is satisfied, let b be a fuzzy closed set in Y , then $cl(f^{-1}(b)) \leq f^{-1}(scl(b)) = f^{-1}(b)$. Therefore $f^{-1}(b)$ is a fuzzy closed set in X .

If the condition (3) is satisfied, let b be a fuzzy subset in Y , then $f^{-1}(b)$ is a fuzzy subset in X and $f(cl(f^{-1}(b))) \leq scl(f(f^{-1}(b)))$. Thus $cl(f^{-1}(b)) \leq f^{-1}(scl(b))$. Therefore, since the condition (2) is satisfied, f is a fuzzy continuous function.

Similarly, we can prove in the case (1). \square

LEMMA [4]. Let $g: X \rightarrow X \times Y$ be the graph of a fuzzy mapping $f: X \rightarrow Y$. Then, if a is a fuzzy set in X and b is a fuzzy set in Y , $g^{-1}(a \times b) = a \wedge f^{-1}(b)$.

THEOREM 2.7. Let $f: (X, T) \rightarrow (Y, S)$ be a fuzzy mapping and T^* be an associated supra-topology with T . Let $g: X \rightarrow X \times Y$, given by $g(x) = (x, f(x))$ be its graph mapping. If g is fuzzy s -continuous, then f is fuzzy s -continuous.

PROOF. Suppose that g is a fuzzy s -continuous and a is a fuzzy open set in (Y, S) . Then $f^{-1}(a) = 1 \wedge f^{-1}(a) = g^{-1}(1 \times a)$. Therefore, $f^{-1}(a)$ is a fuzzy supra open set in (X, T^*) . \square

References

1. Chang, C.L, *Fuzzy topological spaces*, J.Math. Anal. Appl. **24**(1968), 182–190.
2. Abd El-Monsef, M.E and Ramadan, A.E, *On fuzzy supra topological spaces*, Indian J. Pure and Appl. Math. **18**(4)(1987), 322–329.
3. Mashhour, A.S., Allam, A.A., Mahmoud, F.S. and Khedr, F.H., *On supra topological spaces*, Indian J. Pure and Appl. Math. **14**(4)(1983), 502–510.
4. Singal, M.K. and Singal, A.R., *Fuzzy alpha-set and alpha-continuous maps*, Fuzzy Sets and Systems **48**(1992), 383–390.
5. Zadeh, L.A., *Fuzzy sets*, Inforn. and Control **8**(1965), 338–353.

Department of Mathematics
Kangwon National University
Chuncheon, 200-701, Korea