

New Species of Two *Psammocinia* Horny Sponges (Dictyoceratida: Irciniidae) from Korea

Chung Ja Sim and Kyung Jin Lee

(Department of Biology, Hannam University, Daejeon 300-791, Korea)

ABSTRACT

Two new species of the family Irciniidae, *Psammocinia wandoensis* n. sp. and *P. samyangensis* n. sp., are described. They were collected from the South Sea, Korea. *P. wandoensis* n. sp. closely resembles *P. rugosa* (Lendenfeld, 1889) from Australia in morphology, but new species differs from *P. rugosa* by the filaments. Though *P. samyangensis* n. sp. is very similar to *P. jejuensis*, but our species differs from *P. Jejuensis* by the fibre shape.

Key words: new species, horny sponge, *Psammocinia*, Korea

INTRODUCTION

To date, the genus *Psammocinia* has been poorly studied so far. Only four species are reported by Lendenfeld (1889), and one is recorded by Bergquist (1995) from New Caledonia. Three species are reported by Cook and Bergquist (1996) from New Zealand. Three *Psammocinian* species are recorded from Korea by Sim (1998).

The characteristic of this genus is that it has large quantities of sand grains throughout its fibres, matrices and surface, and also has many fine collagenous filaments which are special characters of the family Irciniidae (Bergquist, 1980). Bergquist (1995) noted that filaments are separated from the sponging fibre skeleton. In this study we found that the filaments extend from the fibres, which are permeated by many pores.

The present studies were supported by the Basic Science Reserch Institute Program, Ministry of Education, 1997, Procjct No. BSRI-97-4428.

In this study, the specimens from Wando Island were collected by skin diving, Chejudo Island specimens were collected by fishing nets. The characters used in the identification of the *Psammocinia* species are external morphology, fibre type, skeletal pattern and filament size. Light microscopy and SEM (AKASHI, ISI-40) were used for the fibre arrangement and the filaments morphology.

Type specimens are deposited in the Natural History Museum, Hannam University, Daejeon, Korea.

RESULT

Phylum Porifera Grant, 1836 해면동물 문
 Class Demospongiae Sollas, 1885 보통해면 강
 Order Dictyoceratida Minchin, 1900 망각해면 목
 Family Irciniidae Gray, 1867 가는실해면 과

1. *Psammocinia samyangensis* n. sp. 삼양모래해면 (신칭) (Fig. 1 A-J)

Type specimen. Holotype (Por. 30, NHM, Hannam Univ.), Samyang 1 dong (Chejudo Island), 9 July 1991, (SCUBA).

Description. Specimens irregular mass. Size up to 8 cm long, 3 cm thick and 7 cm wide, attached to rock substrate. Surface filamentous membrane mixed with large sand grains. Low, sharp conules, 1.5-2 mm high, 2-4 mm apart. Colour beige in spirit. Texture soft.

Skeleton: Cortex with many primary fibres, 140-400 μm in diameter, with 10-20 μm sand core. Secondary fibres, 50-230 μm in diameter, mixed with small sand grains. In choanosomes, primary fibres difficult to see due to attached sand (Fig. 1E). Sometimes large oxea mixed with sand are associated with secondary fibres. Sometimes thin secondary fibres connected with large sand like bridge (Fig. 1F). Filaments, 3-8 μm , emerge from pores of secondary fibres (Fig. 1G-I). Terminal knob, 10-17.5 μm in diameter.

Remarks. This new species is similar to *P. jejuensis* (Sim, 1998) but our species differs from the latter by the fibre shape, size, and that secondary fibres rarely contain sand. Hard texture of *P. jejuensis* is also difference with new species.

Etymology. The specific name, *samyangensis* is named after the type locality.

2. *Psammocinia wandoensis* n. sp. 완도모래해면 (신칭) (Fig. 2 A-I)

Type specimen. Holotype (Por. 31, NHM, Hannam Univ.), Wando Island, 27 July 1995, 10 m, (sea woman).

Description. Irregular lobate sponge. Cylindrical digitate processes arise. Size up to 9 cm long, 2 cm thick and 13 cm wide, attached to rock substrate. Surface covered with low sharp conules, 1-2 mm high, 3-5 mm apart. Oscule on the top, 0.5-1 mm diameter. Numerous pores. Texture firm and compressible. Live specimen dark grey colour.

Skeleton: Cortex, 1 mm thick sand crust mixed with black colour sand, 20-120 μm in diameter. Many filaments arranged under sand crust. Stout fasciculate columns up to 75-150 μm diameter, fibres simple not complex. Sand attached inside the fibres (Fig. 2D). In choanosomes, primary




Fig. 1. *Psammocinia samyangensis* n. sp.: A, top-side view showing surface; B, surface of specimen (conules); C, endosome of specimen (transitional section, SEM); D, Cortex fibre (SEM); E, cortex secondary fibre mixed with sand; F, choanosome secondary fibre (SEM); G-H, filaments started from secondary fibre (SEM); I, terminal knob comes out from the hole of secondary fibre (SEM). Scale bars = 1 cm (A), 5 cm (B), 300 mm (C-D), 200 mm (E), 300 mm (F), 10 mm (G-I).

columns measure 190–600 μm in diameter. Secondary fibres difficult to see. Collagen filaments, fine 2–6 μm in diameter, terminal knob 12–16 μm in diameter (Fig. 2H, G).

Remarks. This new species stands out among the *Psammocinia* by the peculiar trellis like column mixed with many small sand grain inside of the fibres. Filaments emerge from the primary and secondary fibres. Around the pore is very rough, sometimes larger than filament diameter. The




Fig. 2. *Psammocinia wandoensis* n. sp.: A, side view showing surface; B, surface of specimen (conules); C, endosome of specimen (transitional section, SEM); D, cortex fibre cored with sand; E, fibre of conule (SEM); F, choanosome fibre; G, Cortex primary fibre; H-I, filament comes out from the hole in fibre; J, terminal knob. Scale bars = 2 cm (A), 5 cm (B), 300 μ m (C), 200 μ m (D), 100 μ m (E), 160 μ m (F), 100 μ m (G), 10 μ m (H-J).

filament pore are difficult to see. This species is very closely related to *P. rugosa* (Lendenfeld, 1889) in external morphology.

Etymology. The specific name, *wandoensis* is named after the type locality.

ACKNOWLEDGEMENT

The authors are grateful to Dr. James T. Wetzel, Department of Biology, Presbyterian College, for review of this manuscript.

REFERENCES

- Bergquist, P. R. 1980. A revision of the supraspecific classification of the orders Dictyoceratida, Dendroceratida and Verongida (Class Demospongiae). *N. Z. J. Zool.*, **7**: 443-503.
- Bergquist, P. R. 1995. Dictyoceratida, dendroceratida and verongida from the New Caledonia Lagoon (Porifera: Demospongiae). *Mem. Queensland Mus.*, **38**(1): 1-51.
- Cook, S. de C. and Bergquist, P. R. 1996. New species of dictyoceratid sponges (Porifera: Demospongiae: Dictyoceratida) from New Zealand. *N. Z. J. Mar. Fresh. Res.*, **30**:19-34
- Lendenfeld, R. von. 1889 A monograph of the horny sponges. London: Pub. Roy. Soc. Trübner and Co, Ludgate Hill, E C, **2**: 468-936.
- Sim, C. J. 1998. Three new horny sponges of the genus *Psammocinia* (Dictyoceratida: Irciniidae) from Korea. *Korean J. Syst. Zool.*, **14**(1): 35-42.

RECEIVED: 7 June 1998

ACCEPTED: 28 October 1998

한국 모래해면 속 (망각해면 목: 가는실해면 과)의 2신종

심 정 자 · 이 경 진
(한남대학교 이과대학 생물학과)

요 약

한국의 남해에서 채집된 모래해면 속의 2신종, 삼양모래해면 (*Psammocinia samyangensis* n. sp.)과 완도모래해면 (*P. wandoensis* n. sp.)을 명명하고 기재하였다. 완도모래해면은 호주의 *P. rugosa* (Lendenfeld, 1889)와 외형에서 많이 유사하지만 filaments에서 차이를 나타내며, 삼양모래해면은 제주모래해면 (*P. jejuensis*)과 유사하나 섬유의 형태에서 차이를 나타낸다.