

1988 - 1997

1.

가가

1960 , 1978 (1994)
17.5%, 80 64.7%, 90 15.5% 80 , 60 2.3%, 70
84.5%

, 1971 1991 11.3% (3
, 1992), 1992 1996 , 15.7%가 (,
1997).

(iatrogenic disease)

가

(issue) , 가

Chesney & Ozar(1995) (1) ; , ,
, , (2) ; , (가 , ,
, (3) ; , , (4)

2.

1) 10

2)

3)

1.

	1988	1997	
,	('89),	,	('90),
('89),	('90),	,	('91-'94),
('95),	('95),	,	('92),
('94)	1010	.	
	158	10	
2070		264	

CD ROM

232

1988	1997	3080
390	(12.7%)	
390		8

1.

1

, , , , , , (issue),
 , 8
 , , , , ,
 (1).

< 1 >

1.	(34) :	(3),	(9),	(1),	(6),	(15)
2.	(58) :	(12) -	(4),	(5),	(2),	(1)
		(15) -	(8),	() (2),		(2),
		(31) -	(1),	(2)		
				(14),		(9),
			(5),		(2),	
			(1)			
3.	(30) :	(15),	(15)			
4.	(82) :	(36),	(19),	(14),	(6),	(7)
5.	(32) :	() (8),	(15),	.	.	(9)
6.	() (72)					
	(13) -		(2),	92),	(1),	(1),
		(1),	(1)	(4)		
	(59) -		(14),	(11),	(11),	() (8),
		(6),	가	(7),	(1),	가
					(1)	
7.	(50) :	()	(12),	(8),	(7),	(7),
		(2),	(4),			(2),
		(2),	(4) -	,	,	,
		-	91),			(1)
8.	(32) :		(27),			(5)

2.

가 70.8% 가
 , 가 14.4% ,
 (Q)가 14.1% (2).
 0.7% 32 가 37.5% (12)
 () 19
 가 .

3.

1.

34 (8.7%) . (3) ,
 (9) , (1) 10 20-45 가
 . 40-60 (6

), (15) (3).

< 2>

	24(6.2)	3(0.7)	7(1.8)	-	34(8.7)
	47(12.1)	7(1.8)	4(1.0)	-	58(14.9)
	15(3.8)	8(2.1)	7(1.8)	-	30(7.7)
	55(14.1)	10(2.6)	15(3.9)	2(0.5)	82(21.1)
	29(7.4)	2(0.5)	1(0.2)	-	32(8.2)
	51(13.1)	19(4.9)	2(0.5)	-	72(18.5)
	32(8.2)	5(1.3)	12(3.1)	1(0.2)	50(12.8)
	23(5.9)	2(0.5)	7(1.8)	-	32(8.2)
	276(70.8)	56(14.4)	55(14.1)	3(0.7)	390(100.0)

< 3>

10	• ; , 가 , •	•	6
20	• , •	•	7
	(Triangulation)		
	9	2	2 13
40-60	• , , •	•	21
	15	1	5 21

(1)

(13) , 1988 - 1991

(8)가 , ; / , , ;

가 .

(, 1988)

1992 - 1997 10

20

(2)가 triangulation 가 가 (2) 40-60 (2), Colaizzi Van Kaam Q (2), (1996) 가 1 (1997). 15 (3), (3), (2), (2), (2) (1), (1), (1) Kraines 가 Neugarten Rosenberg Rockliff Walker (: ,) 가

가
가
가
(: , 가)

2)

) 58 (14.9%) (4) .

< 4>

				10
				2
	9	1	2	12
				6
				7
				2
	14	1		15
				31
	24	5	2	31

가 가
가
(1) 12 ,
(4) (4-6)
(2) , 8
(1) ,
(1)가 . 2

4

(, 1991)

(process)

(outcome)

(8)

(,)

(4),

(1),

(1) .

2

(2)

가

, 15

6

가

가 2

14

3 ,

2

1

Staintons

Walker

Barnard

Feeding and

Teaching Scale

Price

Mother-Infant Sensitivity Scale, Cohler

Broussard

Neonatal Perception Inventory

가

가

가

가

가
(3) 가
31
가 14 ,
가 9 ,
(5),
(2), 가 1
가
가
) , 가 가 , 가 가 (가 가
가 가
5 가
가
가
가
3)
가 15 ,
가 15 30 (7.7%) , 30, 40 60
(5).

< 5 >

·	·	·	·
·	()	·	15
·	·	()	
·	·	· ()	15
·	·	·	
	15	8	7 30

가

가

가

4)

82 (21.1%)

(1)

36 , 가 32 가 ,

(2), (2) 가 26 ,

5 , 가 5 (6).

< 6 >

가				1
60				1
	.			
	(, ,)			
	.			
	.			
	.		.	32
	.	가	.	
	.	()		
	.			
	.			
	.			
				1
				1
	26	5	5	36

(6), 가 (6), (5)
 . 가 ,
 2 .
 , , , , ,
 ,
 가 .
 (2)
 19 . 가 14 가
 , (4) . 가 12 ,
 Van Kaam Giorgi 가 4 , 가 3 .
 ,
 가 . 가 가

, 가

가

33.76 ,

35.39

가

< 7>

.	.	.	(
.가	.	.)	14
.	.	.		
.	.	.		
.가	.	.		4
.	.	.		
.	.	.		1
	12	3	4	19

(3)

14 (8). (8), (3
)가 . 가 8 가 , 가
(3), (2)가 . 가
, 가
가 가

(8).

(4)

6

Spradley

(5)

7

가 2 . 가 , ,
가 가 , ,
가 .

29
 2 Olson's 3-dimensional circumplex model B. Neuman's health systems model Derogatis (SCL-90-R)가 7

6) ()

72 (18.5%)

(1)

13 '88-'89

가 . 90

, 60

가

가

10

가

(2) ()

50 (10).

10

60

가 38

가

가 19

가 2

(8)

Cox

Anderson

가

가

가

Pender

14

. BSE

BSE

< 10 >

. ' ' 11
 가 8
 가 3 . 40
 ' 가 ' (7), () 8 , , 가
 1 .
 7)
 가
 50 (12.8%) .
 , , , , , , , , ,
 , , (12)
 , , (4)
 (2)가 . , ,
 , , , .
 가
 (4) 10 50
 (3)
 가 .
 가 (20-45) (Colaizzi),
 (2)
 , 가 .
 가 ,
 (8) 10
 , , 가 7 가 ,
 1 . 2
 , 가 가
 , , , , , 가

(7) , 가 .

가 (7) , 10 가

가 가 . 가

가 가 , , Q , 48.5%

4 가 .

(4) ,

Heidegger .

(2) , 가 , ,

가 (2) ,

2

(21-50) , 가 .

가 가 , 가 ,

가 가 . 가 ,

가 가

(11) .

	88	89	90	91	92	93	94	95	96	97	
(. . .),						1	2	1		8	12
				1	1	3			1	2	8
		2		1		1		1		2	7
,			1						5	1	7
,						1			1		2
					1		1		1	1	4
								1	1	2	4
,	1					1					2
					1					1	2
(,)							1	1			2
	1	2	1	2	3	7	4	4	9	17	50

8)

가

가

가

가

32 (8.2%)

(12).

< 12 >

											26
5-8											1
		22				2			3		27
											5
									4		5

(1)

가

27

1988

10

1988 1997 9 1010

2070 3080 390

(12.7%)

1. (34 , 8.7%), (58 , 14.9%), (30 , 7.7%), (76 , 19.5%), (32 , 8.2%), (72 , 18.5%), (50 , 12.8%), (32 , 8.2%) 8

2. 가 70.8% 가 가 14.1%

가 14.1%

3. 10-20 , 가 24 , 5 , 3

4. (12), (15), (31)

가 47

5. (15) (15)

가 15 , 8 , 7

6. 가 21.1% 가 (36), (19), (14), (6), (7)

7. (29)가 가 21

, 10

6

8. (13) (59) , , , 51 , 19 , 2

Pender 가 15

9. , , , , , , 가

10.

1.

2. 가

3. 가 , 가

가

4. 가 .
5. 가 .
6. 가 .
- , , , (1992).
 , 4, 5-16,
- , , , (1994). (
- : 1960-1991.8). , 24(1), 58-69.
- , , , (1997). _____
- , 3(2), 205-214.
- , , , (1996). _____
- , 2(1).
 (1996). : , _____, 5(1), 56-70.
- , , , (1995). _____
- , 1(2), 222-233.
 (1977). _____, 27(4), 933-942.
- , , , , , (1992).
 _____, 22(3), 271-296.
- Chesney, M. A. & Ozar, E. M. (1995). Women & Health : In search of a paradigm. Women's Health Research & Gender Behavior and Policy, 1(1), 3-26.
- Griffith-Kenney, J. (1986). *Contemporary Women's Health*, Addison-Wesley.
- McElmurry, B. J., Noor, K. E. & Parker, R. S. (1993). Women's Health status across the globe : Michigan Initiative for Women's Health, Univ. of Michigan.
- Trippet, S. E. & Bryson, M. R. (1995). A model of women's health nursing, *Health Care for Women International*, 16, 31-41.

- Abstract -

Key concept : Women's Health, Nursing Research

**Analysis of Published Nursing
 Research on Women's Health :
 1988-1997**

Lee, Sook Hee · Park, Young Sook

The purpose of this study was to identify the research contents and designs of nursing papers on women's health.

1010 articles published in 9 Korean nursing journals and 2070 master's theses and doctoral dissertations in nursing science for 10 years from 1988 through 1997 were reviewed.

The percent of nursing papers on women's health was 12.7% (390) out of a total of 3080 papers.

Major findings of 390 nursing papers analyzed were as following :

1. The contents of nursing papers on women's health were classified by 8 categories such as childbirth(19.5%), women's health state and behavior(18.5%), motherhood(14.9%), cultural and social issues(12.8%), menstruation(8.7%), mental health(8.2%), caring activities(8.2%), and surgery(7.7%).
2. The principal designs used in nursing papers on women's health were descriptive and correlational survey(70.8%), experimental design(14.4%), and qualitative research design(14.1%).