

New Records of Four Hydroids (Cnidaria, Hydrozoa) in Korea

Jung Hee Park

(Division of Life Science, College of Natural Sciences,
The University of Suwon, Kyonggi-do 445-743, Korea)

ABSTRACT

Some marine hydroids were collected from the eastern sea of Korea during the period from June 1989 to July 1997. Among the identified species four hydroids: *Bimeria annulata* (Nutting, 1901), *Bougainvilla ramosa* (van Beneden, 1844), *Eudendrium boreale* Yamada, 1954 and *Plumularia undulata* Yamada, 1950 were turned out to be new to the Korean fauna.

Key words: new record, hydroids, Korea

INTRODUCTION

So far that the Korean marine hydroids known by the previous works(Kamita and Sato, 1941; Rho, 1967; 1969; Rho and Chang, 1972; 1974; Rho and Park, 1979; 1980; 1983; 1984; 1986; Park and Rho, 1986; Park, 1988; 1990; 1991; 1992; 1993; 1995; 1997; 1998), are seven subspecies, 130 species of 18 families in two orders.

Some hydroid specimens were collected from off the eastern sea of Korea with fishing nets or scuba divers during the period from June 1989 to July 1997. Among them four hydroids: *Bimeria annulata* (Nutting, 1901), *Bougainvilla ramosa* (van Beneden, 1844), *Eudendrium boreale* Yamada, 1954 and *Plumularia undulata* Yamada, 1950 were turned out to be new to the Korean fauna. The redescriptions and illustrations on the species were given. So the Korean hydroid fauna identified up to date consists of seven subspecies, 134 species of 18 families in two

This work was supported by the Korea Science and Engineering Foundation(Project No. 971-0510-050-2).

orders.

DESCRIPTION

Order Athecatae 민컵하드라총목

Family Bougainvilliidae 보우겐빌하드라과

1. *Bimeria annulata* (Nutting, 1901) 고리마디히드라 (신칭) (Fig. 1A-D)

Garveia annulata Nutting, 1901, p. 166, pl. 15, figs. 1-2; Fraser, 1911, p. 22; Fraser, 1914, p. 117, pl. III, fig. 6.

Bimeria annulata: Torrey, 1902, p. 28, pl. I, figs. 1-3; Hirohito, 1988, p. 92, fig. 32d-g.

Material examined. Sokcho harbour, 19 Jul. 1997, J. H. Park.

Description. Hydrorhiza branched stolonial and arranged densely. Stem arising from stolon, monosiphonic in common but sometimes polysiphonic composed of 2-3 stems on lower part, unbranched or branched one or two times, smooth throughout or annulated irregularly and reached to below 15 mm. Hydranth formed on apical portion of stem, with 10-11 filiform tentacles and arranged in a whole surrounding conical hypostome. Periderm more or less thick and extending to base of tentacles of hydranth forming pseudohydrotheca. Gonophores arising from hydrorhizae, round oval shaped and with short stalks covered with periderm which expanding to cup-like its terminals.

Remarks. The specimens of *Bimeria annulata* from Korea are more or less different from the original one of Nutting (1901) whose stem and branch have the regular and distinct annulations. But Torrey (1902) examined many specimens including the type specimen of Nutting and pointed out that stems are not so annulated as in the figures of Nutting.

Distribution. Korea, Japan (Sagami Bay), Pacific coast of North America.

2. *Bougainvillia ramosa* (van Beneden, 1844) 분지보우겐빌하드라 (신칭) (Fig. 1E-J)

Eudendrium ramosum van Beneden, 1844, p. 56, pl. 4.

Bougainvillia ramosa: Hincks, 1868, p. 109, pl. 19, fig. 2; Stechow, 1913, p. 60; Stechow, 1919, p. 26; Stechow, 1925, p. 411; Briggs, 1931, p. 281; Broch, 1933, p. 11; Vervoort, 1946, p. 135, figs. 52a, 53; Yamada, 1959, p. 24; Vannucci and Rees, 1961, p. 82; Millard, 1975, p. 97, fig. 33E-H; Hirohito, 1988, p. 97, figs. 34b-f.

Bougainvillia benedeni Bonnevie, 1898, p. 484, pl. 26, figs. 34-35.

Bougainvillia vanbenedeni: Stechow, 1919, p. 25.

Material examined. Jumunjin, 18 Jul. 1997, J. H. Park; Sokcho harbour, 19 Jul. 1997, J. H. Park.

Description. Colony relatively large, reached 50 mm high and much branched profusely or irregularly. Stem and branch polysiphonic or monosiphonic. Periderm undistinct, smooth or undulating, or annulated partially or throughout, and expanding to base of tentacles forming pseudohydrotheca. Hydranth with a whole of 6-8 filiform tentacles. Medusa buds arising from stem and branch, oval-shaped and with short pedicel.

Remarks. According to Millard (1975) *B. ramosa* is variable in its growth form. One is short,




Fig. 1. A-D, *Bimeria annualata*. A, hydranth; B, hydrorhiza; C, D, gonophores. E-J, *Bougaivillia ramosa*. E, F, polysiphonic stems with pseudohydrothecae and annulated branches; G-H, hydranth; I, gonophore; J, part of colony. Scale bars = 0.4 mm (A-J).

reaching 15 mm long, unfascicled and little branched. The other is tall, as our specimens, reaching 50 mm, fascicled and much branched.

Distribution. Korea, Japan (Sagami Bay), North Sea, North Atlantic, South Atlantic, Europe, Mediterranean, Australia, South Africa, Indonesia.

Family Eudendriidae 풀하드라과

3. *Eudendrium boreale* Yamada, 1954 북방산풀하드라 (신칭) (**Fig. 2A-E**)

Eudendrium boreale Yamada, 1954, p. 3, fig. 2a-f.

Material examined. Sokcho harbour, 20 Jul. 1997, J. H. Park.

Description. Colony relatively large, reached 8–10 cm, and hydrorhiza forming small mass with twisted stolons. Stem monosiphonic but polysiphonic at basal portion and irregularly branched one or two times more and then forming branchlet bundle at apical portion. Stem and branch not in one plane, distinctly, regularly and very closely annulated throughout. Periderm expanded to base of hydranth, forming small cup. Hydranth with about 20 filiform tentacles. No gonophores examined.

Remarks. According to Yamada (1954) male gonophores (Fig. 2D) two-chambered in a whole on the base of the hydranth that is not aborted. Female gonophores (Fig. 2B-C) oval, in dense clusters around the body of hydranth which are usually not aborted.

Distribution. Korea, Japan (Hokkaido).




Fig. 2. A-D, *Eudendrium boreale*. A, E, annulated stems and branches with hydranths; B-C, female gonophores (from Yamada, 1954); D, male gonophore (from Yamada, 1954). Scale bars = 0.1 mm (A, E).




Fig. 3. *Plumularia undulata*. A, hydrorhiza with thick periderm and internal projections; B, a part of hydrocladial axis with hydrothecae; C, E, large gonothecae arising from hydrocaulus; D, hydrocaulus with hydrocladial and nematothecae. Scale bars = 0.3 mm (B-E), 0.5 mm (A).

Order Thecatae 컵히드라총목

Family Plumulariidae 깃히드라과

4. *Plumularia undulata* Yamada, 1950 파도깃히드라 (신칭) (Fig. 3A-E)

Plumularia undulata Yamada, 1950, p. 17, pl. 1, figs. 24-26; Yamada, 1958, p. 80.

Material examined. Sokcho harbour. 28 Jun. 1989, J. H. Park; Jukbyon, 21 Jul. 1997, J. H. Park.

Description. Colonies small, about 1-2 cm long, simple or pinnate, and arising from hydrorhiza creeping on seaweeds. Hydrorhiza flattened, with very thick periderm and internal projections. Stem monosiphonic, divided into regular internodes, each internode bearing a hydrocladial apophysis: 3 caudine nematothecae, one placed in anterior of its base and one pair in base of hydrocladium. Apophyses arranged in right and left sides alternately, not in one plane, placed toward anterior surface. Hydrocladia usually unbranched, but rarely branched, divided into regular internodes, consists of short nematothecate internodes in 0.10-0.11 mm long and long thecate internodes in 0.21-0.23 mm long. Nematothecate internode with one median nematotheca and thecate internode with one hydrotheca and 3 nematothecae: one median inferior, one pair laterals. Nematotheca small, trumpet-shaped, two chambered and immovable. Hydrothecae bowl-shaped, with large aperture, margin undulated and one tongue-like adcauline projection, total length in 0.10-0.12 mm, width of margin in 0.20-0.21 mm, and length of frontal process in 0.06-0.07 mm. Gonotheca large compared to hydrotheca, obconical shaped, wall smooth, margin truncate or with a low and broad neck, total length in 0.42-1.20 mm, width of margin in 0.40-0.63 mm.

Remarks. *Plumularia undulata* resembles *P. filicaulis* in the large aperture of hydrotheca, thick periderm, and very large gonotheca compared to hydrotheca. But it is distinguished from *P. filicaulis* in the position, shape and tongue-like adcauline projection of hydrotheca and the shape of gonotheca.

Distribution. Korea, Japan (Hokkaido).

REFERENCES

- Bonnevie, K., 1898. Zur Systematik der Hydroiden. Zeitsch. Wiss. Zool., **43**: 465-495.
- Briggs, E. A., 1931. Notes on Australian athecate hydroids. Rec. Aust. Mus., **18**: 279-282.
- Broch, H., 1933. Zur Kenntnis der Adriatischen Hydroidenfauna von Split, Arten und Variationen. Skr. norske Vidensk. Akad., mat. nat. Kl., **4**: 1-115.
- Fraser, C. M., 1911. The hydroids of the west coast of North America. With special reference to those of the Vancouver Island region. Bull. Labs. nat. Hist. St. Univ. Ia, **6**: 3-91.
- Fraser, C. M., 1914. Some hydroids of the Vancouver Island region. Trans. R. Soc. Can., sect., **4**, **8**: 99-216.
- Hincks, T., 1868. A history of the British hydroid zoophytes, vol. 1. Text. John Van Voorst Paternoster Row, London: 1-337.
- Hirohito, Emperor of Japan, 1988. The hydroids of Sagami Bay. Bio. Lab. Imp. Household, Tokyo, Japan: 1-110.
- Kamita, T. and T. N. Sato, 1941. Marine fauna at Jinsen (Inchon) Bay, Corea. J. Chosen Nat. hist. Soc., **8**: 1-3.
- Millard, N. A. H., 1975. Monograph on the Hydroidea of South Africa. Ann. S. Afr. Mus., **68**: 1-513.

- Nutting, C. C., 1901. Papers from the Harriman Alaska Expedition. 21. The hydroids. Proc. Wash. Acad. Sci., **3**: 157-216.
- Park, J. H., 1988. Three hydroids (Cnidaria: Hydrozoa) from Ullüngdo and Chejudo, Korea. Kor. J. Syst. Zool., **4**: 57-66.
- Park, J. H., 1990. Systematic study on the marine hydroids (Cnidaria: Hydrozoa) in Korea I. Kor. J. Syst. Zool., **6**: 71-86.
- Park, J. H., 1991. Systematic study on the marine hydroids (Cnidaria: Hydrozoa) in Korea II. The families Sphaerocorynidae, Eudendriidae, Haleciidae and Lafoeidae. Kor. J. Zool., **34**: 541-547.
- Park, J. H., 1992. Zoogeographical distribution of marine hydroids (Cnidaria: Hydrozoa: Hydroida) in Korea. Kor. J. Syst. Zool., **8**(2): 279-300.
- Park, J. H., 1993. Marine hydroids (Cnidaria: Hydrozoa: Hydroida) from Cheju Island, Korea. Kor. J. Syst. Zool., **9**(2): 261-280.
- Park, J. H., 1995. Hydroids (Cnidaria: Hydrozoa: Hydroida) from Chindo Island, Korea. Kor. J. Syst. Zool., **11**(1): 9-17.
- Park, J. H., 1997. Four athecate hydroids from Korean waters. Kor. J. Syst. Zool., **13**(2): 83-92.
- Park, J. H., 1998. Three new records of thecate hydroids from Korean waters. Kor. J. Syst. Zool., **14**(1): 59-66.
- Park, J. H. and B. J. Rho, 1986. A systematic study on the marine hydroids in Korea 9. The families Sertulariidae. Kor. J. Zool., Special Issue no. 1: pp. 52.
- Rho, B. J., 1967. Marine hydroids from the west and south sea of Korea (1). Kor. Cult. Res. Inst. Better Liv., Ewha Womans Univ., **10**: 341-360.
- Rho, B. J., 1969. Studies on the marine hydroids in Korea (2). Kor. Cult. Res. Inst. Better Liv., Ewha Womans Univ., **2**: 161-174.
- Rho, B. J. and S. R. Chang, 1972. Texonomic study on the marine hydroids 3. Marine hydroids from Jeju-Do and Chuja-Kundo. J. Kor. res. Inst. Better Liv., Ewha Womans Univ., **9**: 15-43.
- Rho, B. J. and S. R. Chang, 1974. On the classification and distribution of the marine benthic animals in Korea 1. Hydroids. J. Kor. res. Inst. Better Liv., Ewha Womans Univ., **12**: 133-158.
- Rho, B. J. and J. H. Park, 1979. A systematic study on the marine hydroids in Korea 5. Athecata hydroids. Kor. J. Zool., **22**: 165-174.
- Rho, B. J. and J. H. Park, 1980. A systematic study on the marine hydroids in Korea 6. Thecata. J. Kor. Res. Inst. Better Liv., Ewha Womans Univ., **25**: 15-43.
- Rho, B. J. and J. H. Park, 1983. A systematic study on the marine hydroids in Korea 7. Nine unrecorded species. J. Kor. Res. Inst. Better Liv., Ewha Womans Univ., **31**: 39-56.
- Rho, B. J. and J. H. Park, 1984. A systematic study on the marine hydroids in Korea 8. On two new species belonging to family Plumulariidae. Kor. J. Zool., **27**: 255-263.
- Rho, B. J. and J. H. Park, 1986. A systematic study on the marine hydroids in Korea 9. The family Plumulariidae. J. Kor. Res. Inst. Better Liv., Ewha Womans Univ., **37**: 87-112.
- Stechow, E., 1913. Hydropolipen der Japanischen Ostküste. Abh. d. II. ki. d. k. Ak. D. Wiss. 3, **3**: 1-162.
- Stechow, E., 1919. Zur Kenntnis der Hydroidenfauna des Mittelmeeres, Amerikas und anderer Gebiete. Zool. Jahrb., **42**: 1-172.
- Stechow, E., 1925. Hydroiden der Deutschen Tiefsee-Expedition. Deutsche Tiefsee-Expedition 1898-1899, XVII: 383-546.

- Torrey, H. B., 1902. The hydroids of the Pacific coast of North America. Univ. California Pub., Zool., **1**: 1-104.
- van Beneden, P. J., 1844. Recherches sur l'embryogenie des Tubulaires, et l'histoire naturelle des différents genres de cette famille qui habitent la côte d'Ostende. Mem. Acad. r. Belg., **17**: 1-72. (from Millard, 1975)
- Vannucci, M. and W. J. Rees, 1961. A revision of the genus *Bougainvillia* (Anthomedusae). Bolm Inst. oceanogr. S. Paulo, **11**: 57-100.
- Vervoort, W., 1946. Hydrozoa (C 1) A. Hydropolypen. Fauna Nederl., **14**: 1-336.
- Yamada, M., 1950. The fauna of Akkeshi Bay, 17. Hydroids. J. Fac. Sci. Hokkaido Univ. Ser. 6, Zool., **10**: 1-20.
- Yamada, M., 1954. Species of the genus *Eudendrium* from Japan. Publ. Akkeshi Mar. Biol. Stat., **2**: 1-19.
- Yamada, M., 1958. Hydroids from the Japanese Island Sea, mostly from Matsuyama and its vicinity. J. Fac. Sci. Hokkaido Univ., Ser. 6. Zool., **14**: 51-63.
- Yamada, M., 1959. Hydroida fauna of Japanese and its adjacent waters. Akkeshi Mar. Biol. Stat., **9**: 1-101.

RECEIVED: 3 June 1999

ACCEPTED: 28 August 1999

한국미기록 히드라충류 (자포동물문: 히드라충강) 4종

박 정 희

(수원대학교 자연과학대학 생명과학부)

요 약

1989년 6월부터 1997년 7월까지 우리 나라의 동해에서 채집한 히드라충류 가운데 4종, 고리마디히드라 [*Bimeria annulata* (Nutting, 1901)], 분지보우겐빌히드라 [*Bougainvilla ramosa* (van Beneden, 1844)], 북방산꽃히드라 [*Eudendrium boreale* Yamada, 1954] 그리고 파도깃히드라 [*Plumularia undulata* Yamada, 1950]가 한국미기록종으로 판명되어 재기재하고 보고한다. 이로서 지금까지 밝혀진 한국 해산 히드라충류는 2목, 18과의 134종 및 7아종이 된다.