

Dose-Dependent Inhibitory Effect of Nitric Oxide on Embryo Development

Sea-Hee Park, Bo-Sun Joo*, Hwa-Sook Moon*, Yoon-Yeon Kim**,
Jae-Dong Cho**, Ho-Sung Kang and Han-Do Kim

Department of Molecular Biology, College of Natural Science, Pusan National University,
Pusan 609-735, *Infertility Clinic, Moon Hwa Hospital, Pusan 601-062,
**Cho Jae Dong OB/GY, Chang-Won 641-530, Korea

Nitric Oxide의 농도 의존적인 배발생 억제 효과

부산대학교 분자생물학과, 문화병원 불임클리닉*, 조재동 산부인과의원**

박세희 · 주보선* · 문화숙* · 김윤연** · 조재동** · 강호성 · 김한도

= 국문초록 =

목적: 일산화질소 (nitric oxide; NO)는 생식계를 비롯한 여러 생체내 기관에서 다양하고도 중요한 작용을 하는 것으로 알려져 있으며, 복강액은 난관내강과 연결되어 복강액 내의 세포 성분의 변화는 난관의 미세환경을 변화시켜 수정과 초기 배아 발생에 영향을 줄 수 있다. 본 연구는 배아 발생에 있어서 일산화질소의 역할을 이해하고 복강액 내의 NO 농도 변화가 배아 발생에 미치는 역할을 조사하기 위해 수행되었다.

방법: 과배란시킨 1세대 잡종 암컷 생쥐 (C57BL×CBA/Ca)로부터 1세포기 배아를 얻어 10% synthetic serum substitute가 첨가된 modified human tubal fluid 배양액에서 4일 동안 체외배양하였다 (대조군). 실험을 위해 이러한 배양조건에 sodium nitroprusside (SNP)를 0~1 mM의 다양한 농도로 배양초기부터 첨가하거나, 200 μ M SNP를 2-, 4-, 8-세포기의 각기 다른 배아 시기에 첨가하였으며, 복강경수술을 받는 42명의 여성으로부터 채취한 복강액을 SSS 대신 단백질원으로 사용하여 포배아가 지의 배아 발달을 관찰하였으며, 복강액 내의 NO 농도를 Griess 방법에 의해 측정하였다. 배아의 apoptotic body는 H33334 염색법으로 조사하였으며 배아 발달율은 3회 이상 반복 실험한 결과의 mean±SEM으로 나타내었다.

결과: SNP는 농도에 의존적으로 배발생을 억제하였으나 배아 단계에 대한 특이성은 관찰할 수 없었으며, 특히 100 μ M 이상의 고농도의 SNP는 2-세포기 단계에서 배아 발생을 정지시켰다. 또한 단백질원으로 복강액 이용시 배발생율은 복강액 내의 NO 농도에 따라 현저한 차이가 발견되었으며, 2.5 μ M 이상의 NO를 함유한 복강액에서 배양한 배아의 발생율은 현저하게 감소하였다. cGMP analogue인 8-bromo-cGMP를 배양액에 첨가시 배아 발생에는 변화가 없었으며, SNP에 의해 배발생이 정지된 2-세포기 배아에서 apoptotic body를 발견할 수 없었다.

결론: 이상의 결과로 보아 NO는 고농도에서 배아 발생을 저해하며, 복강액 내의 NO와 같은 성분의 변화는 배아 발생에 유해한 효과를 유발할 것으로 사료된다. 이러한 NO의 배아 발생 억제효과는 cGMP로 중재되는 경로나 apoptosis 유발과는 관계가 없는 것 같다.

Key Words: Nitric oxide, Sodium nitroprusside, Peritoneal fluid, Embryo development

INTRODUCTION

Approximately 14~40% of infertile couples have infertility caused by immunologic disorders which result from the production of anti-sperm antibody in humoral immune response¹ and the soluble products of activated lymphocytes and macrophages in cellular immune response.^{2,3}

Especially, endometriosis which is inflammatory state characterized by the presence and growth of ectopic endometrial tissue outside the uterine cavity, occurs at 20~40% of high incidence in infertile women.⁴ As peritoneal fluid is able to go into the oviductal cavity, changes of cellular and hormonal components within peritoneal fluid may influence the early reproductive processes such as fertilization and early embryo development by modulating the microenvironment that surrounds the embryo.⁵ Peritoneal fluid in women with endometriosis has been reported to increase the number of activated macrophages⁴ and the production of cytokines such as interleukin-1 (IL-1),⁶ interleukin-6 (IL-6)⁷ and tumor necrosis factor- α (TNF- α).^{8,9} These cytokines are known to have adverse effects on sperm cell function, fertilization and embryo development.¹⁰ However, the correct mechanism for this inhibition is unclear.

It is an established fact that these cytokines stimulate macrophages to produce nitric oxide (NO) during conversion of L-arginine to L-citrulline by NO synthase.¹¹ It has been reported that NO plays important roles in various physiological systems. The physiologic actions of NO are regulating smooth muscle relaxation, platelet aggregation and adhesion, cell growth, neurotransmission, apoptosis and immune response.¹² Since the cells of these systems are integral parts of the reproductive organs, NO is thought to play an important role in reproduction. Indeed, many studies have demonstrated that NO regulates pulsatile hypothalamic GnRH secretion,¹³ human granulosa-luteal cell

steroidogenesis,¹⁴ ovulation¹⁵ and penile erection.¹⁶ Therefore, it is suggested that the increased production of NO from the activated macrophage by endometriosis or certain other causes may affect directly or indirectly fertilization and embryo development causing infertility.

NO induces cell death via apoptosis or necrosis. This effects of NO were thought to be solely mediated via activation of 3',5'-cyclic guanosine monophosphate (c-GMP).¹⁷ However, recent studies have demonstrated that NO can also induce biological effects via non-cGMP dependent pathways.^{18,19}

Therefore, this study was performed to understand the effect of NO on embryo development using sodium nitroprusside (SNP), stable NO donor and peritoneal fluids collected from women undergoing laparoscopic operation. NO concentration in PF was measured to determine whether NO level within PF correlates with the embryo development. We also investigated the occurrence of apoptosis following by treatment of SNP and the effect of cGMP analogue on embryo development to understand the mechanism and the pathway for the action of NO.

MATERIALS AND METHODS

1. Materials

The original stock of C57BL/6 \times CBA/Ca hybrid F1 female mice was purchased from Korean Laboratory Animal Center (Korea). PMSG (pregnant mare's serum gonadotropin), hCG (human chorionic gonadotropin), hyaluronidase, mineral oil, sodium nitroprusside, *N*-(1-naphthyl)-ethylendiamine (NED), sulphanilamide, H₃PO₄, NaNO₂, 8-bromo-cGMP, trypan blue, Hoechst 33342 and sodium citrate were purchased from Sigma Chemical Company (St. Louis, MO). Synthetic serum substitute (SSS) was purchased from Irvine Scientific (Santa Ana, CA). Culture dish (60 mm) was purchased from Falcon Inc. (Becton Dickinson, USA). SNP and 8-bromo-cGMP were prepared as a

concentrated stock in serum free modified human tubal fluid (m-HTF) and stored at -20°C before use.

2. Zygote embryo collection and culture

Zygote embryos were obtained from superovulated hybrid F1 female mice (4-6 weeks old). Superovulation was induced by intraperitoneal injections of 5 IU of PMSG followed by 5 IU of hCG after 48 hours. Mating was confirmed by the appearance of a vaginal plug in the next morning. Zygotes were collected from excised oviducts 18 to 20 hours after hCG injection and cultured in m-HTF medium supplemented with 10% SSS at a 37°C incubator in 5% CO₂ for 4 days. Adherent cumulus cells were removed by micropipetting in serum free modified human tubal fluid (m-HTF) medium containing 0.1% hyaluronidase.

3. Treatment of SNP and cGMP-analogue

Zygotes were treated with the various concentration of SNP (0, 100 nM, 10 µM, 50 µM, 100 µM, 200 µM, 1 mM) or 8-bromo-cGMP, an analogue of cGMP (0, 50 µM, 100 µM, 500 µM, 1 mM). And 200 µM SNP was treated at 2-cell, 4-cell and 8-cell embryonic stage to examine the effect of SNP is dependent on SNP treatment stages. Experiments were repeated three or more times and 10-20 embryos were usually used in each experiment.

4. Peritoneal fluid processing

Peritoneal fluid (PF) was obtained from 42 women undergoing laparoscopy for various causes (e.g. endometriosis, myoma) and was centrifuged at 800 g for 10 minutes within 30 minutes after collection. The supernatant was divided into two parts. One part was inactivated complement proteins in a water bath at 56°C for 30 minutes and stored at -20°C to use for embryo culture. The other was immediately stored to assay NO concentration. Before used in culture, an aliquot of PF was filtered (0.22 µm filter, Millex GS, Millipore Corporation,

Bedford, MA) and 1 ml of 10% PF in m-HTF medium was provided for *in vitro* zygote embryo culture. The groups of control were cultured in m-HTF medium containing 10% SSS. Experiments were repeated three times for each PF sample and 10-20 zygote embryos were used in each experiment.

5. Measurement of NO

NO concentration in PF was measured using Griess method.²⁰ Griess assay reagents [Nitroprusside 0.2% *N*-(1-naphthyl)-ethylendiamine (NED) and 2% sulphanilamide in 5% concentrated H₃PO₄] were premixed and then incubated with a peritoneal fluid sample (ratio 1:1) to form a purple azo dye, and absorption was determined at a wavelength of 540 nm by ELISA Reader. NaNO₂ was used to generate standard curve, and NO concentration was calculated from standard curve.

6. Detection of Apoptosis in embryos

To observe the occurrence of apoptosis, normal developing 2-cell embryos, which were not treated with SNP, and 2-cell embryos arrested by treatment of 200 µM SNP were stained with trypan blue and incubated with Hoechst 33342 working solution for 3 min. Hoechst 33342 (Sigma) dye was prepared as stock solution (1 mg/ml) and added to 2.3% sodium citrate in ethanol (10 µl/ml) for working solution. The stained samples were mounted and the characteristics of apoptosis was examined under a fluorescence microscope.

7. Statistical analysis

Statistical analysis was performed by the Student's *t*-test and a *p* value of <0.05 was considered significant.

RESULTS

1. Inhibition of mouse embryo development by SNP treatment

To investigate the effect of nitric oxide on

Table 1. Effect of SNP on mouse embryo development depending on the SNP treatment stages

SNP*	No. of developed embryos to				
	Zygote	2-cell	4-cell	Morula	Blastocyst
control	60	51	51	48	48
zygote	60	48	0	0	0
2-cell	–	60	30	16	0
4-cell	–	–	54	27	0
8-cell	–	–	54	30	0

*200 μ M SNP treated to 2-cell, 4-cell and 8-cell embryonic stage during zygote embryos were cultured to blastocyst stage

mouse embryo development, SNP was treated to mouse zygote culture media with various concentration (0 to 1 mM). The rate of embryo development to blastocyst in the different concentration of SNP is shown in Fig. 1. The addition of SNP inhibited mouse embryo development in a dose-dependent manner. The rates of embryo development were 71% in control, 53% in 0.1 μ M, 48% in 10 μ M, 27% in 50 μ M, 8% in 100 μ M, 0% in 200 μ M and 1 mM. Especially, above 200 μ M of SNP mouse embryo development was completely arrested and degenerated since 2-cell embryonic stage. To determine whether this effect of SNP was treatment stage-specific, 200 μ M SNP was treated at 2-cell, 4-cell and 8-cell embryonic stage. As shown in Table 1, mouse embryo did not develop to blastocyst whenever SNP was treated.

2. The relationships between the level of NO in PF and embryo development

The addition of PF in mouse embryo culture media resulted in the different development rate. We measured the concentration of NO in PF to determine whether this different rate was caused by changes of cellular component within PF. NO concentration was different from each PF sample ranged from 0.79 μ M to 4.4 μ M. PF was divided into 6 groups according to its NO level and the rates of embryo development were represented in Table 2. The




Fig. 1. Effect of SNP on mouse embryo development. SNP was treated with various concentrations (control, 100 nM, 10 μ M, 50 μ M, 100 μ M, 200 μ M, 1 mM) to embryo culture medium. Embryos were cultured from zygote to blastocyst and embryonic development was examined. Development rate was expressed with % of developing to blastocysts from zygotes. Data is mean of three experiments. Values in vertical bar are the number of embryos reached to blastocyst/the total number of embryos provided for experiment. * p <0.05.

rate of embryo development was significantly lower in group V ($26.5 \pm 16.6\%$) and group VI ($29.2 \pm 11.0\%$) compared with control ($62.8 \pm 6.1\%$), which was cultured in medium with 10% SSS (p <0.05). However, the rate of embryo development in 2.5 μ M or less (group I to IV) was similar compared with that of control.

Table 2. Effect of peritoneal fluid on mouse embryo development according to the level of NO in peritoneal fluid

Group (n*)	NO level (μM) (Mean \pm SEM)	No. of embryos	% of developed embryos [†] to			
			2-cell	4-cell	Morula	Blastocyst
	control	120	87.4 \pm 2.9	85.2 \pm 3.3	76.9 \pm 4.5	62.8 \pm 6.1
I (9)	≥ 1.0 (0.84 \pm 0.05)	90	96.6 \pm 2.1	92.7 \pm 3.0	84.9 \pm 2.9	68.3 \pm 8.6
II (7)	1.0~1.5 (1.20 \pm 0.07)	70	100 \pm 0.0	98.2 \pm 1.7	91.9 \pm 4.0	70.0 \pm 8.0
III (5)	1.5~2.0 (1.81 \pm 0.07)	50	95.5 \pm 4.4	95.8 \pm 4.1	93.0 \pm 4.1	91.9 \pm 5.2 ^a
IV (8)	2.0~2.5 (2.15 \pm 0.03)	80	90.2 \pm 3.6	89.9 \pm 4.6	82.7 \pm 4.9	71.2 \pm 5.2
V (5)	2.5~3.0 (2.68 \pm 0.04)	50	85.5 \pm 11.9	70.5 \pm 23.6	40.3 \pm 24.9 ^a	26.5 \pm 16.6 ^a
VI (8)	<3.0 (4.36 \pm 0.05)	80	89.0 \pm 3.3	69.8 \pm 12.3	42.8 \pm 23.6 ^a	29.2 \pm 11.0 ^a

Note: Peritoneal fluid was obtained from 42 women with endometriosis of other causes by laparoscopy

*; n = number of peritoneal fluid sample

[†]; Mean \pm SEM

^a; p<0.05 (versus control)

3. Inhibitory effect of NO on the mouse embryo development via cGMP-dependent or independent pathway

To determine whether this effect of NO on mouse embryo development was mediated by the elevating level of cGMP, 8-bromo-cGMP was added to embryo culture media. As noted in Fig. 2, embryo development was not influenced by 8-bromo-cGMP regardless of the treated concentrations (50 μM to 1 mM). This result suggests that NO may inhibit mouse embryo development through cGMP-independent pathway.

4. Detection of apoptosis in embryo arrested by SNP treatment

To investigate whether the arrest of embryo development following SNP treatment resulted from cell death via NO-mediated apoptosis, we examined the nucleus fragmentation in 200 μM SNP treated 2-cell embryos. Compared with normal developing embryos, the morphological changes of nucleus were not detected in embryos treated with SNP and the size of nucleus was equal to that of control (Fig. 3).




Fig. 2. Effect of cGMP-analogue (8-bromo-cGMP) on mouse embryo development. 8-bromo-cGMP was treated with concentrations of 0 to 1 mM to embryo culture medium. Embryos were cultured from zygote to blastocyst and embryonic development was examined and development rate was expressed with % of developing to blastocysts from zygotes. Data was mean of three experiments. Values in vertical bar are the number of embryos reached to blastocyst/the total number of embryos provided for experiment. *p 0.05.

DISCUSSION

NO can pass easily into the intracellular area and has many different effects on cell metabolism in the biology, physiology, and pathophysiology of reproduction.¹² Because NO is a




Fig. 3. Fluorescence micrographs of 2-cell embryos stained with Hoechst 33342 for detection of apoptotic bodies. Zygote embryos were culture in m-HTF medium without (Fig. 3a) or with (Fig. 3b) 200 μ M SNP. After 24 hours, 2-cell embryos were stained with Hoechst 33342 and observed at $\times 400$ by fluorescence microscope.

free radical, it is degraded easily and has very short life span. Thus, most studies used a more stable nitric oxide donor to elucidate the role of NO *in vitro*. SIN-1, sodium nitroprusside (SNP), SNAP are representative nitric oxide donor. Among them, SNP released only NO *in vitro* cell culture system, while other two donors released superoxide anion or/and hydrogen peroxide as well as NO,²¹ and SNP concentration in cell is known to be in proportion to production of NO.²² Weinberg *et al.*²³ reported that SNP reduced sperm motility and this inhibition was blocked by the NO quencher hemoglobin. This suggests that SNP is more suitable to study the effect of only NO compared with other NO donors.

Our present study showed that mouse embryo development to blastocyst was inhibited dose-dependently by SNP, especially reduced at high concentration of 100 μ M or more. However, this SNP-induced inhibitory effect on embryo development was not stage-specific because embryo development was inhibited regardless of SNP treatment stages. We also found that embryo development was severely arrested

in media containing PF with the higher concentration of NO than in media containing qualified controlled serum. It has been recognized that PF is able to go into the oviductal cavity and may influence the early reproductive processes by modulating the microenvironment that fertilization and early embryo development occur. These results imply that high concentration of NO may adversely affect embryo development, and that changes of component within PF especially such as increased concentration of NO can be closely associated with embryo development.

Previously, many studies have observed that the level of macrophage, cytokines and other cellular components were increased and activated with the extent of pelvic disease and inflammation.^{5,8,9} PF in women with endometriosis adversely affected the cleavage of mouse two-cell embryos²⁴ and contained high concentration of cytokines such as TNF- α and IL-1.¹⁰ However, the correct mechanism for interaction between these cytokines in PF and embryo toxicity has not been elucidated yet. A common pathway of the action of TNF- α on various

cell systems is the induction of an inducible nitric oxide synthase (iNOS), which results in the generation of NO. Thus it is possible that a certain disease such as endometriosis or other factors stimulate the production of cytokines, which subsequently induce the production of NO thereby directly affecting embryo development.

Generally the effects of NO were known to be mediated via activation of soluble guanylate cyclase and guanosine 3',5'-cGMP, which inhibits Ca^{2+} influx and activates its efflux, resulting in a lower intracellular $[Ca^{2+}]$. Decreased intracellular $[Ca^{2+}]$ can inactivate PKC, and thus inhibit PKC-dependent cell proliferation.²⁵ But, the inhibitory effect of NO on mouse embryo development may be not mediated by a cGMP-dependent pathway, since the addition of 8-bromo-cGMP, agent which raises cellular cGMP, did not affect embryo development regardless of treated concentration. This result is supported by recent studies, which suggest that NO can induce its biological effects via non-cGMP-dependent pathways by forming high-affinity-nitroso complexes with a variety of metal compound or binding the heme-containing proteins and iron-sulphur-containing proteins.^{14,18,19}

The effects of NO via these pathway may be dependent on dose and the cell type. One hundred micromole of SNP was shown to inhibit sperm motility,²⁶ while lower concentration of SNP (100 nM or less) was beneficial to the maintenance of post-thaw sperm motion, viability and motility.²⁷ And NO has the double-edged role in inducing apoptosis. High level of NO induced by pathophysiological conditions stimulates apoptosis while the continuous release of endothelial NO inhibits apoptosis.²⁸ We investigated whether the inhibition of embryo development by higher concentration of SNP was caused by inducing apoptosis of embryo, but apoptotic characteristics did not found in embryos arrested by treatment of SNP.

Another possibility we can consider for this

inhibitory effect is the inhibition of embryonic-stage specific protein synthesis by SNP. A lot of stage-specific proteins are necessary for normal embryo development. NO is capable to posttranslationally interfere with heme proteins and enzymes containing iron-sulfur cluster or to modulate enzyme activity by S-nitrosylation.²⁹ Thus, the further study is needed to prove the correct mechanism for the inhibition of embryo development.

In conclusion, high concentration of SNP and PF containing high amounts of NO inhibited early embryo development. This result suggests that *in vivo*, if there is increase of NO level in reproductive systems such as peritoneum and tubal fluid by certain causes, this may correlate with infertility by causing damage to embryo.

REFERENCES

1. Naz RK, Menge AC. Antisperm antibodies. Origin, regulation, and sperm reactivity in human infertility. *Fertil Steril* 1994; 61: 1001-13.
2. Hill JA, Cohen J, Anderson DJ. The effects of lymphokines and monocytes on human sperm fertilizing ability in the zona-free hamster egg penetration test. *Am J Obstet Gynecol* 1989; 160: 1154-9.
3. Jung BJ, Lee SH, Hur M. The efficacy on the immunotherapy with paternal lymphocytes in unexplained infertility. *Korean J Fertil Steril* 1997; 24: 293-300.
4. Ramey JW, Archer DF. Peritoneal fluid: its relevance to the development of endometriosis. *Fertil Steril* 1993; 60: 1-14.
5. Badawy SZA, Cuenca V, Marshall L, Munchback R, Rinas AC, Coble DA. Cellular components in peritoneal fluid in infertile patients with and without endometriosis. *Fertil Steril* 1984; 42: 704-8.
6. Fakh H, Baggett B, Holtz G, Tsanh KY, Lee JC, Williamson HO. Interleukin-1: a possible role in the infertility associated with endometriosis. *Fertil Steril* 1987; 47: 213-7.

7. Harada T, Yoshioka H, Yoshida S, Iwabe T, Onohara Y, Tanikawa M, Terakawa N. Increased interleukine-6 levels in peritoneal fluid of infertile patients with active endometriosis. *Am J Obstet Gynecol* 1997; 176: 593-7.
8. Halme J, Hammond MG, Hulka JK. Retrograde menstruation in healthy women and in patients with endometriosis. *Obstet Gynecol* 1984; 64: 151-4.
9. Eisermann J, Michael JG, Jorge P, Odem RR, Collins JL. Tumor necrosis factor in peritoneal fluid of women undergoing laparoscopic surgery. *Fertil Steril* 1988; 50: 573-9.
10. Taketani Y, Kuo TM, Mizuno M. Comparison of cytokine levels and embryo toxicity in peritoneal fluid in infertile women with untreated or treated endometriosis. *Am J Obstet Gynecol* 1992; 167: 265-70.
11. Bredt DS, Snyder SH. Nitric oxide: A physiologic messenger molecule. *Annu Rev Biochem* 1994; 63: 175-95.
12. Rosselli M, Keller PJ, Dubey RK. Role of nitric oxide in the biology, physiology and pathophysiology of reproduction. *Human Reprod* 1998; 4: 3-24.
13. Moretto M, Lopez FJ, Negro-Vilar A. Nitric oxide regulates luteinizing hormone-releasing hormone secretion. *Endocrinology* 1993; 133: 2399-402.
14. Van Voorhis BJ, Dunn MS, Snyder GD, Weiner CP. Nitric oxide: An autocrine regulator of human granulosa-luteal cell steroidogenesis. *Endocrinology* 1994; 135: 1799-806.
15. Hwsa JS, Preutthipan S, Maguire MP, Chang TS, Wallach EE, Dharmarajan AM. Nitric oxide modulates human chorionic gonadotropin-induced ovulation in the rabbit. *Fertil Steril* 1997; 67: 548-52.
16. Holmquist F, Stief CG, Joncs U. Effect of the nitric oxide synthase inhibitor NG-nitro-L-arginine on the erectile response to cavernous nerve stimulation in the rabbit. *Acta Physiol Scand* 1991; 143: 299-304.
17. Murad F. Regulation of cytosolic guanylyl cyclase by nitric oxide: the NO-cyclic GMP signal transduction system. *Adv Phamacol* 1994; 26: 19-33.
18. Brunswig-Spickenheier B, Mukhopadhyay AK. Stimulation of nitric oxide-cyclic guanosine monophosphate pathway in bovine ovarian theca cells by tumor necrosis factor- α (TNF- α). Is this pathway implicated in the TNF- α -induced inhibition of uterine hormone-stimulated progesterone production. *Biol Reprod* 1997; 57: 700-6.
19. Kroncke KD, Fehsel K, Kolb-Bachofen V. Nitric oxide: cytotoxicity versus cytoprotection; How, Why, When, and Where? *NITRIC OXIDE: Biol Chem* 1997; 1: 107-20.
20. Marzinzig M, Nussler AK, Stadler J, Marzinzig E, Barthlen W. Improved methods to measure end products of nitric oxide in biological fluids: Nitrite, Nitrate, and S-Nitrosothiols. *NITRIC OXIDE: Biol Chem* 1997; 1: 177-89.
21. Ioannidis I, Groot H. Cytotoxicity of nitric oxide in Fu 5 rat hepatoma cells; evidence for cooperative action with hydrogen peroxide. *Biochem J* 1993; 296: 341-5.
22. Kowulak EA, Seth P, Fung HL. Metabolic activation of sodium nitroprusside in vascular smooth muscle. *J Pharmacol Exp Ther* 1992; 262: 916-22.
23. Weinberg JB, Doty E, Bonaventura J, Haney AF. Nitric oxide inhibition of human sperm motility. *Fertil Steril* 1995; 64: 408-13.
24. Morcus RN, Gibbons WE, Findley WE. Effects of peritoneal fluid on *in vitro* cleavage of 2-cell mouse embryos: possible role in infertility associated with endometriosis. *Fertil Steril* 1985; 44: 678-83.
25. Khatib AM, Siegfried G, Quintero M, Mitrovic DR. The mechanism of inhibition of DNA synthesis in articular chondrocytes from young and old rats by nitric oxide. *NITRIC OXIDE: Biol Chem* 1997; 1: 218-25.
26. Rosselli M, Dubey RK, Imthurn B. Effect of nitric oxide on human spermatozoa: evidence that nitric oxide decreases sperm motility and induces sperm toxicity. *Human Reprod* 1995; 10: 1786-90.
27. Hellstrom WJG, Bell M, Wang R. Effect of so-

- dium nitroprusside on sperm motility, viability, and lipid peroxidation. *Fertil Steril* 1994; 61: 1123-8.
28. Dimmeler S, Zeiher AM. Nitric oxide and apoptosis: Another paradigm for the double-edged role of nitric oxide. *NITRIC OXIDE: Biol Chem* 1997; 1: 275-81.
29. Stamler JS, Jaraki O, Osborne J. Nitric oxide circulates in mammalian plasma primarily as a S-nitroso adduct of serum albumin. *Proc Natl Acad Sci* 1992; 89: 7674-6.
-