Cu	ret	Ultras EPMA	onic	Scaler			SEM	
		I.						가
	フ	' 						
1,2)		,		,	가 E	Zander18) EDTA Ca Mg)	•
.,_/.							가 Ca, Mg, P	,
		3),			フト 19 - 21)			
1-6)	•		7,8)	,		, 1979	Wirthlin	22)
		0 47\71		가 .	Ca P			,
	4.0	9 - 17)フト		,		,	-	40 <i>μ</i> m
9 - 15 - 17)	· 14)	,				, I	Barton ²⁵ spersive)	
		,			analysis(EDX)	onorgy an		, 60μπ

Ca P	•	•
	,	2.
Micro - Analysis EPMA	EPMA(Electron Probe)가	, 가7-9 mm
	,	. 0.9%
가 , 가 ,	가 X - ray가 X - ray /	
7),	26 -	
EPMA	, Cohen	
28)	, Hennequin Ca P	100%, 100% 12
, Atil	la ³⁰⁾	12 2
가	·	. 2:1 12 , 1:2 12 48
	,	70 8 - 12 .
가	EPMA ,	2 - 3
II.		(2)
1.		- ,
	가 40 , 가 7 - 9mm 가	. 1.2 - 1.5mm aluminum stub carbon tape
		stub tarbori tape

Tο	h	ما	1	C_{2}	leve	ч

				Table 1. Ca l	evel
4000 4000	1	600,		GC	US
1000 - 1200	sand paper	r 3	Cervical 184.0 ± 5.7		178.5 ± 23.3
			Middle	171.5 ± 3.5	172.5 ± 21.9
9μ m film	(blue color)	3 , $3\mu\mathrm{m}$	Apical	155.5 ± 2.1	163.5 ± 9.2
film(pink cold color) 5	,	μm film(green	(GC : Grace	y curettes US: UI	trasonic scaler)
0.05μm alumir	na powder	silicone			
5	•		JEOL, Jap	oan) 70	
가 1.0	mm 1.4	lmm가			

3 spray air

가

(3) EPMA

EPMA system(JXA - 8900R, JEOL, Japan)

2mm x 2 3 mm

WDS(wavelength dispersive x -

ray spectrometer) Ca, P, Mg, Na

JXA - 8900R Accelerating Voltage: 20.0 kV Dwell Time: 30.0 msec,

Magnification: 120, Probe Diameter (μm): 2,

Probe Current (A): 1.900 - 2.000

(4) **EPMA**

Gold - Palladium (JSM - 840A, III.

1. EPMA

(1) Ca level

Ca level Table 1 Table 2. P level

	GC	US
Cervical	62.5 ± 5.0	66.0 ± 1.4
Middle	60.5 ± 2.1	65.0 ± 1.4
Apical	55.5 ± 0.7	61.5 ± 2.1

(GC : Gracey curettes US : Ultrasonic scaler)

가 4가 level Ca level

Ca level

Ca level (Figure 1 - 6).

Table 3. Mg level

	GC	US
Cervical	4.0 ± 0.0	4.5 ± 0.7
Middle	4.5 ± 0.7	4.5 ± 0.7
Apical	3.5 ± 0.7	4.5 ± 0.7

(GC : Gracey curettes US : Ultrasonic scaler)

(2) P level

Table 4. Na level 가

								·
	GC	US			Calvia	1 18)		
Cervical	1.0 ± 0.0	1.0 ± 0.0	0		, Selvig) 10)		
Middle	1.0 ± 0.0	1.0 ± 0.0	0					
Apical	1.0 ± 0.0	1.0 ± 0.0	0	, Furs	seth 24	1)		
(GC : Gracey	curettes US:UI	trasonic scaleı	r)					,
	P level	Γable 2					,	
4가		level						
,		Co lovel	,		71	,	5)	
		Ca level			가 _	. Darton	3)	
	•				E	PMA		
	С	a level				, Hennequin	29)	EPMA
	(F	igure 7 - 12)).			,		
Ca level .			•		Ca P		N	Мg
(3) Mg le	vel			EPM	A	•		
() 3	Mg level	Table 3	_			ispersive Sp	ectro -	metry)
Ca P	g		level	-	Navele	-		spersive
			10 7 01	•		''g''' 가 가		EDS
, (Figure 13 - 18).				=	DS	21 21	,	ιυς ,
					X -	ray spectrui	m	
(4) Na lev	/el			١٨/	/DS	, 26)	Barto	n ²⁵)
(4) 144 101		Table 4		Cohen	28)	Hennequ		
		i able 4	•	Conen	20)	пеппеци	111 207	
	level ,					,		WDS
(Figure 19 - 24).					-			

2.

EPMA

evaporation evaporation

・ 1μm 가

(Figure 25, 26). alumina powder .

, Ca, P, Na


 $0.05 \mu \mathrm{m}$


Atilla ³⁰⁾	EDMA CEM
Cohen ²⁸⁾	EPMA SEM
Ca P 가 , Ca P level , Ca, P,	1. Ca, P, Mg, Na level . 2. Ca, P, Mg, Na 가
Mg, Na level . O, Leary 31) Coldiron 32)	3. Ca P
ЕРМА	4.
. Ca P Mg 가 Selvig (1962) ¹⁸⁾ , Neider (1972) ¹⁹⁾ , Hals (1977) ²⁰⁾ , Totdal (1985) ²¹⁾ , Cohen (1991) ²⁸⁾	5. Ca, P, Mg 가
, , , , , , , , , , , , , , , , , , , ,	, 가 .
Ca P 가 ,	VI.
가 , ,	 Socransky SS: Microbiology of periodontal disease. Present status and future considerations. J Periodontol 48:497 1977. Van Palenstein Helderman WH:
, 가 , EPMA WDS 가 ·	Microbial etiology of periodontal dis- ease. J Clin Periodontol 8:261 1981. 3. Zappa U, Smith B, Simona C, Graf H, Case D and Kim W; Root Substance Removal by Scaling and Root Planing. J Periodontol 62:750 - 754 1991. 4. Polson AM, Lahenheim S and Hans PJ
	: Cell and fiber attachment to diminer - alized dentin from periodontits - affected root surfaces. J Periodontol


- 57:235 246 1986.
- Polson AM and Hans PJ: Cell and fiber attachment to demineralized dentin: A comparison between normal and periodontitis affected root surfaces. J Clin Periodontol 14: 357 - 365 1987.
- 6. Fukazawa E and Nishimura K : Superficial Cemental Curettage : Its Efficacy in Promoting Improved Cellular Attachment on Human Root Surfaces Previously Damaged by Periodontitis. J Periodontol; 65:168-176 1994.
- Tagge DL, O'Leary TJ and EI-Kafrawy AH: The clinical and histological response of periodontal pockets to root planningand oral hygiene. J Periodontol 46:527 - 533 1975.
- 8. Hughes TP and Caffesse RG: Gingival changes following scaling, root planning, and oral hygiene. A biometric evaluation. J Periodontol 49:245 1978.
- Kerry GJ: Roughness of root surfaces after use of ultrasonic instrument and hand curettes. J Periodontol 38:340 1967.
- Wilkinson RF and Maybury JE: Scanning electron microscopy of the root surface following instrumentation. J Periodontol 44:559 1973.
- Van Volkinburg JW, Green E and Armitage GC: The nature of root surface after curette, cavitron and alphasonic instrumentation. J Periodont Res 11:374 1976.
- Meyer K and Lie T: Root surface roughness in response to periodontal instrumentation studied by combined use of microroughness measurements


- and scanning electron microscopy. J Clin Periodontol 4:77 1977.
- 13. D'Silva IV, Nayak RP, Cherian KM and Mulky MJ: An evaluation of the root topography following periodontal instrumentation: A scanning electron microscopic study. J Periodontol 50:283 1979.
- 14. Garnick JJ and Dent J: A Scanning Electron Micrographical Study of Root Surfaces and Subgingival Bacteria After Hand and Ultrasonic Instrumentation. J Periodontol. 8;441 -447 1989.
- 15. Thornton S and Garnick JJ: Comparison of ultrasonic to hand instruments in the removal of subgingival plaque. J Periodontol 53:35 1982.
- 16.Gellin RG, Miller MC, Javed T, Engler WO and Michkin DJ: The effectiveness of the titan S sonic scaler versus curettes in the removal of subgingival calculus. J Periodontol 57:672 1986.
- Sherman PR, Hutchens LH, Jewson LG, Moriarty JM, Greco GW and McFall WT: The Effectiveness of Subgingival Scaling and Root Planing. I. Clinical Detection of Residual Calculus. J Periodontol61:3 - 8 1990.
- 18. Selvig KA and Zander HA: Chemical analysis and microradiography of cementum and dentin from periodontally diseased human teeth. J Periodontol 33:303 1962.
- 19. Neiders ME, Eick JD, Miller WA and Leitner JW: Electron probe microanalysis of cementum and underlying dentin in young permanent teeth. J Dent Res 51:122 1972.

- 20. Selvig KA and Hals E: Periodontally diseased cementum studied by correlated microradiography, electron probe analysis and electron microscopy. J Periodont Res 12:419 1977.
- 21. Totdal B and Hals E: Electron probe study of human and red deer cementum and root dentin. J Dent Res 93:4 12 1985.
- 22. Wirthlin MR, Pederson ED, Hancock EB, Lambert BL and Leonard EP: The hypermineralization of diseased root surface. J Periodontol 50: 125 127 1979.
- 23. Selvig KA: Biologic changes at the tooth saliva interface in periodontal disease. J Dent Res 48: 846, 1969.
- 24. Furseth R : A study of experimentally exposed and fluoride treated dental cementum in pigs. Acta Odontol.Scand. 28: 305 322.1970.
- Barton NS, Van Swol RL; Periodontally diseased Vs. normal roots as evaluated by scanning electron microscopy and electron probe analysis. J Periodontol. 58:634 - 638 1987.
- 26. X ray compositional microanalysis: EDS and WDS. Available from Http://gea.geo. unioma1.it/microsonda.html
- 27. Electron probe microanalysis laboratory. Available from Http://www.seismo.berke-ley.edu/geology/labs/epma/index.html
- Cohen M, Garnick JJ, Ringle RD, Hans PJ and Thompson WO: Calcium and phosphorus content of root exposed to the oral environment. J Clin Periodontol 19:268 273
 1992
- 29. Hennequin M and Douillard Y: Effects of citric acid treatment on the Ca, P and Mg contents of human dental roots. J Clin Periodontol 22:550 557 1995
- 30. Atilla G and Baylas H: Electron probe analysis of cementum surfaces. J Marmara Univ Dent Fac 2:510 514 1996
- 31. O, Leary TJ and Kafrawy AH: Total cementum removal: A realistic objective? J Periodontol 54:221 226 1983
- 32. Coldiron NB, Yukna RA, Weir J and Caudill RF: A quantitative study of cementum removal with hand curettes. J Periodontol 61:293 299 1990
- Figure 1. Ca Level at Cervical portion (US)
- Figure 2. Ca Level at Cervical portion (GC)
- Figure 3. Ca Level at Middle portion(US)
- Figure 4. Ca Level at Middle portion (GC)
- Figure 5. Ca Level at apical portion (US)
- Figure 6. Ca Level at apical portion (GC)
- Figure 7. P Level at Cervical portion (US)
- Figure 8. P Level at Cervical portion (GC)












Figure 25




Figure 26

- Figure 9. P Level at Middle portion (US)
- Figure 10. P Level at Middle portion (GC)
- Figure 11. P Level at apical portion (US)
- Figure 12. P Level at apical portion (GC)
- Figure 13. Mg Level at Cervical portion (US)
- Figure 14. Mg Level at Cervical portion (GC)
- Figure 15. Mg Level at Middle portion (US)
- Figure 16. Mg Level at Middle portion (GC)
- Figure 17. Mg Level at apical portion (US)
- Figure 18. Mg Level at apical portion (GC)
- Figure 19. Na Level at Cervical portion (US)
- Figure 20. Na Level at Cervical portion (GC)
- Figure 21. Na Level at Middle portion (US)
- Figure 22. Na Level at Middle portion (GC)
- Figure 23. Na Level at apical portion (US)
- Figure 24. Na Level at apical portion (GC)
- Figure 25. SEM (US)
- Figure 26. SEM (GC)

- Abstract -

The Study on the Root Surfaces with SEM and EPMA Following Periodontal Treatment with Curet and Ultrasonic Scaler

Jae - Hyuk Kim, Chong - Yeo Kim, Sung -Bin Lim, Chin - Hyung Chung Department of Periodontology College of Dentistry Dan - Kook University

One of the fundamental causes of periodontal disease is accumulation of bacterial plaque and calculus and most effective method of removing these plaque and cal culus are scaling and root planning using hand curet and ultrasonic scaler. Many studies concerning residual degenerated mineral content after periodontal therapies have been carried out, but some problems about these studies were also known. This research studies mineral concents and distribution of residual root surfaces after perfoming hand curet and ultrasonic scaler on root surfaces of single rooted teeth which were extracted for periodontal reasons. EPMA were used to avoid errors from chemical quantative analysis and in addition SEM observation was also performed.

The results were as follows.

1. No differences were found between curet group andultrasonic scaler group

- in Ca, P, Mg and Na level.
- 2. Concentration level was decreased in the sequence of Ca, P, Mg and Na.
- Ca and P level were decreased as going to apical portion at curet group and ultrasonic scaler group.
- More cementum was removed at cervical portion compared to other portion at curet group and ultrasonic scaler group.
- 5. Ca, P, Mg level was higher in dentin compared to cememtum.

There was no difference in mineral level for Ca, P, Mg and Na between root surfaces treated with hand curet and ultrasonic scaler.