



가  
가  
5  
( , , , 1987).

가  
가  
1  
가  
가  
(1991) 8  
4  
(1996)  
가  
가  
가  
( , 1980).

(Mccloskey & Bluecheck, 1992 :  
Jensen & boback. 1985. :Reeder & Martin. 1987).

가  
( )  
가  
가  
가 ,  
(belief system)  
(practice system)  
( , 1993, 1997). (1993, 1995, 1997, 1998)  
가가  
가  
6가  
(Know - how)  
6가  
’ , ’ ,  
’ , ’ ,  
’ , ’ ,  
’ , ’ ,

**2.**

가  
 , , 18-20 , 2-3  
 3-4 ,  
 3-4 .  
 20%가 3-4  
 1 , (Bryant, overland. 1966  
 : , , , 1987).

(Ladwig, 1990). 가  
 , , ,  
 ( , 1994), , , ,  
 가 (Carr, 1989).

1.

1-4  
 81  
 , 가 ,  
 3 20 6 5 , 81

2.

가 ,  
 1 1 , 2 ,  
 2 pretest  
 SAS ,

Pearson Correlation

1.

가 26-30 가 ,  
 가 , 가 100-199 ,  
 가 . ,  
 가 42.0% , 가 59.3% , 가 40.7% . 58.0% ,  
 가 가 , , 1-3

가 가 .

2. .

2-3 , , ,

가 가 . 17.6 ,

122.9 , 1 69.800 ,

가 ,

가 , , ,

(1 ) ,

34.6%

, 29.6%가

가 가 , 23.5%

(1 ),

(1 )가 , 가 (5 ), (1

), (1 ), (3 ), (4 ), (1 )

가 가 .

3. .

70%

가 ,

tm

2.5 ,

가 ,

가 가 ,

, 가 , 가

0.05 가

가 가 ,

61.7% 가 ,

가 . 28.4%

가 가 , 37%가

가

(23.5%),

가 (3.7%),

(4.9%)

가

가

(1 )가

(1 ),

(1 ),

(3 ),

(4 ),

(1 )

가 가

(1 ),

(5 ),

가

가

,가

가

가

가

가

가

가

가

가

가 1 69,800 가 ,  
 가 가 가  
 가 6 8 12  
 , tm , ,  
 , 6 10  
 , , , ,  
 , , , ,  
 .  
 1) , ,  
 , ,  
 2) 가 가 가  
 가 가 가  
 3) , , ,  
 가 가 가  
 , , ,  
 4) , , ,  
 가 가 가  
 5) , , ,  
 , 가 , , ,  
 , , , ,

(1980). , 19(5), 23.  
 (1991). .  
 (1993). : , 2(2),  
 37-65.  
 (1997). , 36(5), 74-90.  
 (1995).

- , 25(4), 825-836.  
 (1998). : .  
 , 37(3), 27-40.  
 (1997). , 3(1),  
 5-19.  
 (1993). 가 .  
 , (1987). : .  
 Carr-Hill, R., Cambell, D. M., Hall, M. H., & Meredith, A. (1987). Is birth weight determined genetically. *British Medical Journal-Clinical Research*, 295(6600), 687-689.  
 Ladwig, P. W., London, M. L., & Olds, S. B. (1990). *Newborn Nursing*. 2nd, ed. Redwood Addison Welsey. co.  
 McCloskey, J. C., & Bluecheck, G. M. (1992). *Nursing untervention classification*. Mosby co. 387-390.  
 Reeder, S. J., Martin, L. L., & Koniak, D. (1987). *Marternity Nursing*. 392--393.  
 Rutldege, D. L., & Pridham, K. F. (1987). Post partum Mathers perception of competence for infant care. *JOGNN*, May/June 6, 185-194.

**- Abstract -**

key concept : Post-partum Care Center,  
 Post- partum mother, Nursing needs for Post-partum mother

**A Study on the Nursing Needs and  
 Service Satisfactory of Users  
 in the Post-Partum Care Center**

**Jung, Hyun**

These days, as the society has been in the trends of highly industrialized and the family has been downsized, there is remarkably increasing number of women who follow occupation. These changes have made it more difficult for the family to help post-partum mother, which had been performed in side of house.

By the help of social believe that during at least 1 month after childbirth professional nursing program is indispensable for both maternity protection and physical-mental recuperation, now many post-partum care centers for post-partum mother have been in operation. Although these post-partum care center have in use for a long time, no study was performed before this study on the same subject. Data were analyzed using by SAS. The results of study are as follow :

1. The general features of the user of the post-partum care center. The predominant band of user's age is extended from 26 to 30. The users are mainly housewives and

they are in higher level of incomes and educations. As for the feature of delivery methods, they performed the normal spontaneous vaginal delivery method by 58.0% and the Caesarean operation method by 42.0%. As for the sexuality for babies, 59.3% of infants are male, and 40.7% are female. The highest delivery order of users is first and admission after 1-3days delivery is highest.

2. The results for the investigation into the actual condition of the post-partum care center are as follow : About the main reason for entrance of the post-partum care center was found to be the needs for the better nursing programs for recuperation after childbirth. This demands are also supported by their husbands. The average length of stay in the post-partum care center is 17.6 days and the besides promised expense ; powdered milk, milk-suckers, disposable dippers, skin cares, body shape cares, entrophics, injections. The post-partum have private rooms for mother and infants. Over the half of average expense is 229 million won. They are paying accessory fees the post-partum care center have similar step organization : the nurses, the skin carers, the cleaners. Mostly both ways of feeding powdered milk and breast feeding are in use, and mother's milk is preserved in the night time to be given to infant by nurse in charge.

3. The results of the investigation on users' nursing demand to the post-partum care center and satisfaction are as follow : The ranking order of nursing demand of mothers who used the is that ; infant care demands, environmental demands, emotional and mental care demands, education and training demands. As much as 70% of the respondents have dissatisfaction in nursing program, The ranking order of service satisfaction of mothers who used appears to be higher following order ; satisfaction in infant care demand, satisfaction in physical nursing demand, satisfaction in emotional and mental nursing demand, satisfaction in education and training demand, satisfaction in environment nursing demand. The results of pearson correlation. between nursing demand and service satisfaction of mothers who used are found to be relative noticeable in the level of 0.05. only infant care. The 61.7% of the women who used the post-partum care center. are responding that they will reuse the same post-partum care center again.