

Crystallization and Preliminary X-ray Crystallographic Studies of HslU Mutant in *Escherichia coli*

Ji-Joon Song, Cheol Soon Lee,[‡] Yun Sik Kim, Jung Jin Kim, Young-Jun Im, Hyun Hee Kim,[‡] Soon Ji Yoo,[‡] Ihn Sik Seong,[‡] Chin Ha Chung,[‡] and Soo Hyun Eom^{*}

Department of Life Science, Kwangju Institute of Science and Technology, Kwangju 500-712, Korea

[‡]School of Biological Sciences, Seoul National University, Seoul 151-742, Korea

Received May 10, 2000

HslVU is an ATP dependent protease in *E. coli* like proteases La (Lon) and Ti (ClpAP/ClpXP).¹ It is composed of two multimeric components, 19 kDa HslV and 50 kDa HslU proteins.² While HslU itself has an ATPase activity, HslV has a weak peptidase activity so that it slowly degrades certain hydrophobic peptides, such as *N*-carbobenzoxy-Gly-Gly-Leu-7-amino-4-methyl-coumarin and polypeptides such as insulin B-chain and casein.^{3,4} But the proteolytic activity of HslV is increased significantly (up to 150 fold) by associating with HslU in the presence of ATP.³ The primary amino acid sequence of HslV is similar to certain β -type subunits of the 20S proteasomes of archaeobacterium *Thermoplasma acidophilum* with 18% identity.⁵ While β -type subunits of the 20S proteasomes show 72-point symmetry, HslV is a dimer of hexamers with 62 point symmetry. The crystal structure of HslV solved at 3.8 Å resolution shows that in spite of the different symmetry, the folds and the contacts between subunits are conserved, compared with β -type subunits of the 20S proteasomes.⁶ In the case of HslU, it is 50% identical to the ClpX protein of *E. coli* in amino acid sequence. According to the analysis of HslU using electron microscopy, HslUs make ring-shaped forms in the presence of ATP or AMP-PNP (ATP analogue). This ring is composed of 6 or 7 HslU molecules to form hexameric or heptameric rings.⁷ HslU contains two Cys residues, Cys261 and Cys287. It has been suggested that Cys261 is involved in oligomerization and that Cys287 is related to the ATPase function.⁸ In order to reveal the three-dimensional structure, and the mechanism of oligomerization between HslUs, and between HslU and HslV, the HslU_{C261V} was crystallized and studied with X-ray crystallographic method.

Experimental Section

The pGEM-T vector (Promega) carrying the *hslVU* operon (named pGEM-T/HslVU) was constructed as described previously.^{6,9} Site-directed mutations were created by the PCR method, which consists of two sequential PCRs, using pGEM-T/HslVU for HslU mutant as the templates. The primary PCR was carried out using mutagenic primers, which were designed to replace Cys²⁶¹ of HslU with Val, and then the second PCR was performed. After the secondary PCR, the mutated fragments were ligated into pGEM-T vector.

The resulting plasmids were digested with *Nru*I and *Bgl*II, followed by ligation of the restriction fragments into the pGEM-T/HslVU_{C261V} plasmids. The resulting plasmids were transformed into *E. coli* strain XL2 Blue. Substitution of the nucleotide by mutagenesis was confirmed by DNA sequencing.

E. coli cells harboring pGEM-T/HslVU_{C261V} were grown overnight, lysed in 2% (w/v) SDS, and electrophoresed on 13% (w/v) polyacrylamide slab gels containing SDS and 2-mercaptoethanol. Two major bands representing 19 kDa (HslV) and 50 kDa (HslU_{C261V}) polypeptides were detected in cells carrying the recombinant plasmids, but not in cells containing only the vector. To purify HslU_{C261V}, the crude extracts of the *E. coli* cells harboring pGEM-T/HslVU_{C261V} were loaded onto a phosphocellulose column equilibrated with 50 mM KH₂PO₄/K₂HPO₄ buffer (pH 6.5) containing 5 mM MgCl₂, 0.5 mM EDTA, 1 mM DTT, and 10% (v/v) glycerol (referred to as buffer A). After collecting the flow-through fraction, the column was washed extensively with the same buffer, and the proteins bound to the column were eluted with the same buffer containing 0.4 M phosphate. HslU_{C261V} was recovered in the flow-through and HslV in the 0.4 M phosphate eluate. For purification of HslU_{C261V}, the flow-through fraction from the initial phosphocellulose column was dialyzed against buffer containing 20 mM Tris-HCl (pH 7.8), 5 mM MgCl₂, 0.5 mM EDTA, 1 mM DTT, and 10% glycerol (referred to as buffer B) and loaded onto a DEAE-cellulose column equilibrated with the same buffer. After washing the column with buffer B containing 0.15 M NaCl, proteins were eluted with a linear gradient of 0.15-0.3 M NaCl. The fractions containing HslU_{C261V} were pooled, diluted 1 : 1 with buffer B, and applied to a heparin-agarose column equilibrated with the same buffer. Proteins bound to the column were eluted with a linear gradient of 0.2-0.5 M NaCl. The HslU_{C261V} containing fractions were pooled, concentrated by ultrafiltration using a YM30 membrane (Amicon), and chromatographed on a Sephacryl S-300 column equilibrated with buffer B containing 0.1 M NaCl. Then HslU_{C261V} was concentrated.

Two crystal forms that belong to different space groups were grown with hanging drop-vapour diffusion method at room temperature (293 ± 1 K). One form of HslU_{C261V} crystals (Figure 1A) was grown on a siliconized cover slip by equilibrating a mixture containing 1 μL of protein solution (10.6 mg mL⁻¹) containing 2 mM ATP in buffer A and 1 μL reservoir solution (100 mM Tris-HCl pH 8.0, 6-8% PEG 4K,

^{*}Corresponding author. Tel: +82-62-970-2493; Fax: +82-62-970-2484; E-mail: eom@kjist.ac.kr




Figure 1. Orthorhombic (A) and hexagonal (B) crystals of HslU mutant (C261V) from *Escherichia coli*. The maximum dimensions are $0.1 \times 0.1 \times 0.1$ mm.

100-300 mM NaCl, 20 mM $MgCl_2$, 2 mM DTT, and 1 mM EDTA) against 1 ml of reservoir solution. The crystals with maximum dimensions of $0.1 \times 0.1 \times 0.1$ mm grew in one week. X-ray diffraction data were collected on an R-Axis IV image-plate system attached to a Rigaku rotating-anode generator (RU-300) providing $CuK\alpha$ radiation and running at 50 kV and 90 mA with a 0.3 mm focus cup at room temperature (293 ± 1 K).

Preliminary intensity data were collected where the diffraction beyond 5.5 \AA resolution was recorded, and the crystals belong to the primitive orthorhombic space group ($P2_12_12_1$ or $P2_12_12_1$). The unit cell parameters of the crystal were determined to be $a=80.0$, $b=138.1$, and $c=175.0 \text{ \AA}$ with

1933400 \AA^3 unit cell volume, using DENZO.¹⁰ Assuming that asymmetric unit contains three molecules with a molecular mass of 49.6 kDa, V_M value is calculated as $3.25 \text{ \AA}^3 D^{-1}$, resulting in a solvent content of 62.2%. The other form of crystals (Figure 1B) was grown under the same crystallization condition, and experiment was performed in the same manner. The crystal diffracted beyond 6.5 \AA . The unit cell parameters were determined as $a=b=81.0$ and $c=174.6 \text{ \AA}$, and this crystal belongs to primitive hexagonal space group ($P6_1$ or $P6_4$). Assuming that asymmetric unit contains one molecule, V_M value is calculated as $3.33 \text{ \AA}^3 D^{-1}$, resulting in a solvent content of 63.1%.

This work is the initial step toward revealing not only the structure of HslU mutant and its complex with HslV complex but also the mechanism of oligomerization. A search for better-diffracting crystals is continuing by varying the crystallizing conditions.

Acknowledgment. This work was in part supported by grants of the Brain Korea 21 project and Critical Technology 21 project (98-NQ-07-01-A) from the Ministry of Science and Technology, Korea.

References

1. Goldberg, A. L. *Eur. J. Biochem.* **1992**, *203*, 9.
2. Chuang, S. F.; Burland, V.; Plunkett, G.; Daniels, D.; Blattner, F. *Gene* **1993**, *134*, 1.
3. Yoo, S. J.; Seol, J. H.; Shin, D. H.; Rohrwild, M.; Kang, M. S.; Tanaka, K.; Goldberg, A. L.; Chung, C. H. *J. Biol. Chem.* **1996**, *274*, 14035.
4. Seol, S. J.; Yoo, S. J.; Shin, D. H.; Shim, Y. K.; Kang, M. S.; Goldberg, A. L.; Chung, C. H. *Eur. J. Biochem.* **1997**, *247*, 1143.
5. Seemüller, E.; Lupas, A.; Stock, D.; Löwe, J.; Huber, R.; Baumeister, W. *Science* **1995**, *268*, 579.
6. Bochtler, M.; Ditzel, L.; Groll, M.; Huber, R. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 6070.
7. Rohrwild, M.; Coux, O.; Pfeifer, G.; Santarius, U.; Müller, S. A.; Huang, H.-C.; Engel, A.; Baumeister, W.; Goldberg, A. L. *Nat. Struct. Biol.* **1997**, *4*, 133.
8. Yoo, S. J.; Shim, Y. K.; Seong, I. S.; Seol, J. H.; Kang, M. S.; Chung, C. H. *FEBS Lett.* **1997**, *412*, 57.
9. Yoo, S. J.; Kim, H. H.; Shin, D. H.; Lee, C. S.; Seong, I. S.; Seol, J. H.; Shimbara, N.; Tanaka, K.; Chung, C. H. *J. Biol. Chem.* **1998**, *273*, 22929.
10. Otwinowski, Z.; Minor, W. *Methods Enzymol.* **1996**, *276*, 307.