

Impact of Salt Intake on Red and Fallow Deer Production in Australia**

- Review -

Y. J. Ru*, P. C. Glatz and Z. H. Miao

South Australian Research and Development Institute, Livestock Systems, Roseworthy Campus
Roseworthy, South Australia 5371, Australia

ABSTRACT : Southern and south-western Australia is a typical mediterranean environment, characterised by wet, cold winters and dry, hot summers. The evaporation rate varies significantly in summer, resulting in a high salinity of drinking water for grazing animals. In addition, a large amount of land in the cropping areas is affected by salt. Puccinellia, tall wheat grass and saltbushes have been planted to improve the soil condition and to supply feed for grazing animals. Animals grazing these areas often ingest an excessive amount of salt from soil, forage and drinking water which can reduce feed intake, increase the water requirement, depress growth and affect body composition as demonstrated in sheep. While the deer industry has been successfully developed in these regions, the potential impact of excessive salt intake on deer production is unknown. The salt tolerance has been well defined for sheep, cattle and other livestock species, but the variation between animal species, breeds within species, maturity status and grazing environments makes it impossible to apply these values directly to deer. To optimise deer production and effectively use natural resources, it is essential to understand the salt status of grazing deer and the impact of excessive salt intake on growth and reproduction of deer. (*Asian-Aus. J. Anim. Sci.* 2000. Vol. 13, No. 12 : 1779-1787)

Key Words : Salt Tolerance, Pasture, Grazing Deer

INTRODUCTION

Sodium is one of the main macro minerals required by animals, and plays an important role in maintaining osmotic pressure, rumen pH and regulating the volume of fluid in the body (Xu et al., 1994). For optimum growth, reproduction and other physiological functions, adequate amounts of sodium need to be supplied through feeds, drinking water and other sources. NRC (1980) reported that minimum concentrations of sodium in the diet are 0.7-0.9 g/kg DM for sheep, and 0.8-1.2 g/kg DM for cattle, with a significant variation between animal species, breeds within species, and maturity status. There is no data available on the sodium requirement of grazing deer, which is a significant disadvantage for deer producers in Australia. Farmers require information on sodium requirement of deer and the effect of high sodium intake on deer production, especially in southern Australia.

This review was requested by the Deer Program Committee of RIRDC to accompany a full research submission on the effect of salt intake on deer production. The review will focus on i) the environmental conditions in cereal cropping areas and the distribution of red and fallow deer in Australia, and ii) sodium intake and its potential impact on deer production under these environments.

ENVIRONMENT CONDITIONS IN CEREAL CROPPING AREAS IN AUSTRALIA

Southern and south-western Australia is a typical Mediterranean environment. The annual rainfall in this area ranges from 200 mm to 1,000 mm, with 85% of the rain falling in winter. The distribution of annual rainfall exhibits marked variation between years and locations (Bellotti et al., 1992). For example, Geraldton receives 80-85% of annual rainfall in the growing season, but Merredin, Clare, Walpeup and Horsham receive about 66%, 61%, 59% and 58% of the annual rainfall during the growing season, respectively. The temperature ranges from 9°C in winter to above 40°C in summer. Mid-winter temperatures are mild (12-15°C) and mid-winter radiation levels are low, ranging from 9 MJ/m².day to 11 MJ/m².day (Perry, 1989). Evapotranspiration rates vary more in summer than in winter, contributing to the salinity problem in the cropping areas.

In the summer rainfall region, covering north-eastern region from central New South Wales to Queensland, rainfall is higher than in the south and increases in both amount and summer dominance to the north. For example, the average annual rainfall for Dubbo, Narrabri, Dalby, and Biloela are 531, 662, 614, and 705 mm, with 42%, 38%, 29% and 19% falling in the winter (May-October) growing season, respectively. Mid-winter day temperatures range 15-19°C. South eastern Australia is a high rainfall zone, with a significant amount of rainfall in both summer and winter. Mid-winter temperatures are about 10-12°C and radiation levels range from 7 MJ/m².day

* Address reprint request to Y. J. Ru. Tel: +61-8-8303-778 7, Fax: +61-8-8303-7977, E-mail: ru.yingjun@saugov.sa.gov.au.

** This paper has been presented to Australian Rural Industries Research and Development Corporation for the development of a research project.

to 9 MJ/m².day (Perry, 1989). However, the maximum temperatures in these regions can be over 40°C, with 20-40 days per year having a maximum temperature over 35°C (Anon, 2000).

In the cropping region, there is a large amount of land with high salinity. In Western Australia in 1980, 263,000 ha was salt-affected land with a further 2.45 million ha having the potential to become salt-affected (Henschke, 1980; Anon, 1988a). By 1989, 315,000 ha was salt-affected (Malcolm, 1989). Salinity also threatens 385,000 ha in Victoria (Anon, 1989), and 2.2% of the state in New South Wales (Anon, 1988c) and affected 210,900 ha in South Australia (Anon, 1988b). Salt affected soils range from sands to clays and include all grades of soils. Except for the most acidic soils, all soils can grow *Puccinellia ciliata*, *Atriplex* spp., *Halosarcia* spp. and *Maireana brevifolia* or combination of these species (Malcolm, 1989). The Department of Agriculture Western Australia recommended strawberry clover, salt water couch, puccinellia, barley grass, tall wheat grass, saltbushes and others for soils with different salinities (table 1, Malcolm, 1986).

DISTRIBUTION OF RED (CERVUS ELAPHUS) AND FALLOW (DAMA) DEER IN AUSTRALIA

Seventeen species or subspecies of deer were introduced into Australia during the last century by Acclimatisation Societies. Of these, only fallow, red, rusa, sambar, chital and hog remain today (SCA,

1980). While there are no accurate estimates of feral and farmed deer in Australia, the figures published in 1980 by the working party appointed by the Animal Production Committee of the Standing Committee on Agriculture are a reasonable guide.

Feral deer

There is a lack of accurate information on the total population of the feral herd. However, there are about 20000 fallow and 11000 red deer distributed in Australia. The largest populations of fallow deer are in Tasmania, with the rest being in South Australia, Victoria, New South Wales and Queensland. Two major feral populations in New South Wales are located at Rangers Valley in the Glen Innes district and Lake George on the South Tablelands. In South Australia, feral fallow deer are distributed in the Adelaide Hills and in the South-East. In Queensland, fallow deer are confined to the west of Stanthorpe between Pikedale and the Severn River, and the area between Warwick and the Dividing Range (SCA, 1980). Fallow deer are the most widespread and numerous in Australia.

Red deer have adapted to most environments and have adjusted from the sub-tropics to the temperate zone, from sea level to 1800 m, from near-desert conditions to more than 760 cm of rainfall, and from forests to open plains. Red deer are most numerous in the Brisbane River Valley of Queensland, the Mudgee-South Coast area of New South Wales, the Grampians of Victoria and in the Pinjarra region of

Table 1. A guide to selecting salt tolerant forage plants for saltland types in the agricultural areas of Western Australia*

Saltland type and condition		Site severity		
		Mild	Moderate	Severe
Hillside seepage	Soil surface wet in summer	<u>Strawberry clover</u> , Paspalum, Couch, Kikuyu	<u>Salt water couch</u>	<u>Salt water couch</u>
	Soil surface dry in summer	<u>Puccinellia</u> , Barley, Tall wheat grass	<u>Puccinellia</u>	<u>Puccinellia</u>
Saline valley floors	Rainfall >375 mm	<u>Puccinellia</u> , Barley, Tall wheat grass	<u>Puccinellia</u> , Saltbushes**	<u>Samphire</u>
	Rainfall <375 mm	Commonly flooded <u>Saltbushes</u> , Puccinellia, Barley	<u>Saltbushes</u> , Puccinellia, samphire	<u>Samphire</u>
		Seldom flooded <u>Barley</u> , Bluebush, Puccinellia, Saltbushes	<u>Saltbushes</u> , Bulebush, Puccinellia	<u>Saltbushes</u> , Samphire Puccinellia
Dryland salinity	Rainfall <375 mm	Never flooded <u>Barley</u> , Bluebush, Saltbushes	<u>Bluebush</u> , Saltbushes	<u>Bluebush</u> , Saltbushes

* The recommended plant is underlined. Others listed are also capable of reasonable growth.

** Saltbushes are only recommended in areas with less than 500 mm annual rainfall.

*** Source: Malcolm (1986).

Western Australia and the ACT (Anderson, 1984).

Farmed deer

While there are more than 40 recognised species and numerous subspecies of deer through the world, many are not suitable for modern agricultural husbandry and must be kept in an open environment. Farmed deer in Australia are limited to the six existing species, and estimated at 2000 head on 300 farms in 1985 (Mackay, 1985) and 160000 head in 1998 (Mackay, 1998). Of these 40% are red, 50% are fallow with 10% of other species. However, fallow deer are dominant species in New South Wales, Tasmania, Western Australia and South Australia, but Queensland has more red deer (English, 1984). The number of red deer has increased dramatically in South Australia during the last few years.

SALT INTAKE OF GRAZING DEER

Animals can ingest salt from water, pastures, soils (Wilson, 1978) and other mineral supplements, making it difficult to quantify the salt intake under grazing conditions. Salt intake of grazing animals is often influenced by environmental conditions (temperature, rainfall), pasture management (grazing intensity, fertiliser application), and animal factors (maturity stage, selective grazing, species or breed etc.).

Pastures

Under a Mediterranean environment in southern Australia, pastures are dominated by annual species such as medics, subterranean clover, ryegrass and barley grass etc. During the vegetative stage, pastures generally have a low sodium content (Xu et al., 1994), which increases with plant maturation. Ru (1996) reported that sodium content (on dry matter basis) in subterranean clover increased from 0.3-0.5% in August to 0.8-1.6% in November, with a large variation among cultivars. Clover seeds have a low sodium content (0.06-0.11% dry matter). In the cropping areas affected by salt, puccinellia (*Puccinellia ciliata*), tall wheat grass (*Thinopyrum ponticum*) and Chenopodiaceous plants (e.g. *Atriplex*, *Kochia* and *Bassia* spp) have been planted for the improvement of soil conditions due to their low transpiration rate, high efficiency of water use, drought and salinity resistance (Sharma, 1978; Wilson, 1978; Leigh, 1986; Barrett-Lennard, 1999).

Puccinellia which originated from the west coast of Turkey, is a winter active perennial grass suited for saline and waterlogged land. It grows in areas with 300 mm or more annual rainfall and performs better on alkaline soils than on acid soils. This pasture is the key to production of grazing animals on salt affected land in Eyre Peninsula and in the Upper

South East in South Australia. The dry matter digestibility of Puccinellia is over 70% during September and November, and protein content declines from 22% in September to 15% in November. While few grazing data are available, the palatability of this pasture varies between varieties. For example, Rottneest Island prostrate ecotypes is highly unpalatable, whereas the prostrate ecotype from Geraldton is highly palatable (Barrett-Lennard, 1999). Tall wheat grass, originated from Balkans, Asia Minor and southern Russia, is adapted to highly alkaline soils and salt seepage areas. It is less tolerant to salinity than Puccinellia. It grows in warmer months, with a moderate palatability and requires a heavy rotational grazing system. It is only moderately palatable (Nichols, 1998). While it has been reported that on saline soils there are increases in salt uptake and in salt concentrations in the plant materials, and the increased salt concentrations in the shoots of waterlogged plants have adverse effects on plant growth and survival in the longer term (Barrett-Lennard and Nulsen, 1989), there is limited information on the salt intake of animals grazing these pasture species.

Chenopodiaceous plants supply a large amount of forage during drought periods and have remarkable capacity to remain leafy and viable under high radiation and temperature levels (Sharma, 1978). The common feature of *Atriplex* leaves is their vesiculated hairs, which are extremely rich in salt and perform functions such as absorption of water from the atmosphere, water storage, insulation against transpiration and salt secretion (Wood, 1925; Mozafar and Goodin, 1970). These shrubs are grazed by sheep and cattle, and the intake of *A. nummularia* (oldman saltbush) was sufficient to maintain sheep and to allow a small gain in weight. The bodyweight production per unit area was similar for sheep and cattle, but sheep selected more green grass than cattle which ate more dry grass and saltbush (Wilson, 1966a). While proximate analyses of the leaves of these shrubs show that they are high in crude protein and ash during summer, the high ash content up to 30% of the dry matter is a disadvantage, as high salt intakes may be detrimental to grazing animals (Wilson, 1966a; Wilson, 1978).

There is a variation in nutritional quality among species of saltbushes. Leaves of saltbushes have a crude protein content of 16.7-25.2%, crude fat of 1.0-1.6%, crude fibre of 7.8-10.4% and ash of 18.5-27.2% on a dry matter basis. Sodium content ranges 0.2-5.6% on a dry matter basis. Digestible dry matter ranges from 74.5% to 78.8% and digestible energy from 3.2 Mcal/kg to 3.4 Mcal/kg (Khilil et al., 1986). Rehman (1992) also found that among 8 saltbush species, the dry matter digestibility ranged

from 33% to 62%, and nitrogen digestibility varied from 51% to 81%. Crude protein content in dry matter varies from 10% to 20% and ash content from 13% to 23%. There was no correlation between preference of sheep and the content of crude protein, ash content or sodium content of saltbush samples. The sodium content in saltbush is much higher than the 0.06% level recommended for beef cattle (NRC, 1976); the level of 0.5% recommended for goats (NRC, 1981) and 0.04-0.1% recommended for sheep (NRC, 1975). The digestible energy content is higher than those required to meet the needs of dry (2.2 Mcal/kg) and lactating (2.5 Mcal/kg) cows (NRC, 1976) and can meet the requirement of sheep and goats (NRC, 1975, 1981). However, the nutritional value of these bushes for deer and the impact of sodium content on deer production are not clear.

Water

Surface water supplies are not regarded as a major source of minerals for the grazing animal. However, evaporation, which is more significant in summer, from streams and lakes results in a concentration of the dissolved solids. Land management strategies also affects the salinity of surface water available for grazing animals. For instance, the replacement of deep-rooted trees by shallow rooting grasses in the clearing of native forest for agriculture in south-west Western Australia increased stream salinity. During the irrigation season, drought periods following prolonged floods, the concentration of salts may increase significantly (Anon, 1976). In the field situation, evaporation from troughs in summer can cause an increase of 10% in concentration of salts even though the troughs are cleaned regularly (Wilson, 1978). Groundwater is more saline than surface water from being in contact with soil and rock for longer periods and has been concentrated by plants through transpiration. Judson and McFarlane (1998) indicated that the quality of underground water is variable and significant quantities of Na, Ca and Mg salt may be consumed by grazing animals. Bore water in the Northern regions of Australia usually contain less than 5000 mg/litre of total dissolved salts whereas in the southern parts of the country concentrations are in excess of 10000 mg/litre of total dissolved salts. This concentration is near the maximum tolerable concentration for mature non-lactating livestock (Judson and McFarlane, 1998).

Holmes and Waterhouse (1983) reported typical concentrations of Na, Ca and Mg in groundwater from selected sites in the south-east of South Australia. The data indicate that water with the higher cation concentration could provide about 20-100% of Na and about 5-20% of the daily Ca and Mg needs of the mature animals. Total salinity is higher in summer

than in winter (table 2). However, in many places, especially in north-eastern and central Australia, the salinity of ground water is unknown (Anon, 1976).

Soils

Livestock cannot avoid ingesting soil during grazing. Factors contributing to an increase in the ingestion of soil are high stocking rates, areas of poor drainage, poor pasture growth, feeding root crops or grain supplements in mud or dust, and when stock are grazing stubbles and subterranean clover pasture where grain and clover burr are buried. On short pastures, cattle can ingest over 100g soil/kg herbage DM (Dewes, 1996) and sheep up to 300 g soil/kg herbage DM (Thornton, 1983). Under normal grazing conditions, ingested soil can be a source of more minerals than ingested pasture (Healy, 1973). No studies have been undertaken to determine the intake of soil by deer while grazing.

IMPACT OF SALT INTAKE ON ANIMAL PRODUCTION

The salt intake has to match the requirement of deer for optimum production. It has been shown that sodium deficiency influences pasture selection and appetite, and increases appetite for salt or salty foods (Blaxter et al., 1974). Research conducted at the Rowett Research Institute also indicated that when supplementing sodium with salt blocks on pastures with low sodium content, stags licked salt blocks and hinds sometimes drank the urine of their sucking calves to increase their sodium intakes. The ratio of sodium and potassium is a valid index of sodium status in deer as in sheep (Blaxter et al., 1974). While there is little data available demonstrating the effect of

Table 2. Representative analyses of water in the South-East of South Australia

Source	Main cations (mg/litre)			Total salinity (mg/litre)
	Na	Ca	Mg	
Unconfined aquifer, Mt Gambier	81	113	18	690
Coastal spring, Ewens Ponds	46	73	20	387
Confined aquifer, Mt Gambier water supply	115	71	30	612
Unconfined aquifer, Padthaway	453	109	63	1,678
Groundwater drain				
July	192	65	38	855
December	333	45	58	1,224

(Holmes and Waterhouse, 1983)

sodium deficiency on deer production, research with other livestock species shows that growth rate, feed utilisation efficiency and milk yield can be depressed if sodium is limiting (Xu et al., 1994).

Although there are no reports on the impact of excessive sodium intake on the performance of grazing deer, the effect of high sodium intake on water intake, feed intake and consequently animal production of other livestock species such as sheep has been studied comprehensively.

Water intake

The water intake from forages of grazing animals is dependent on ambient temperature, forage moisture and forage mineral content. Inglis (1985) reported the normal water intake of animals (table 3), but the type of diet has a significant influence on water requirement (Inglis, 1985; Wilson and Hindley, 1968). For example, water intake of red deer varies from 0.27 litre/kg DM intake for a grass diet to 3.82 litre/kg DM on concentrated pellets (Bonengel, 1969). Fallow deer require less water per unit of feed intake (Denholm, 1984).

Ingestion of excess sodium by stock may lead to a higher water requirement and a decreased tolerance to saline drinking water, especially in the salt affected cropping areas. In these regions saltbush and bluebush may comprise part of the available feed and stock are dependent wholly or partially on salty water. It was evident that when grazing saltbush, water intake of sheep increases. The maximum water intakes varied from 4 litre/day in the summer of a good season to 12 litre/day during a drought (Wilson, 1978).

Table 3. Normal water intakes of animals

Animal	Daily consumption (litres/day)
Merino ewe	Up to 3.7 in summer; 1.6 average over a year
Beef cow	Up to 53 in summer; 35 average over a year
Dairy cattle	70
Horses	45
Pigs	
Sow and litter	23-27
Sow (pregnant)	11-14
Grower	
23 kg	3-5
45 kg	7
68 kg	9
90 kg	11
Poultry (100 birds)	
Laying hens	33
Broilers	30 (higher now)

(Inglis, 1985)

Rumen digestion

Over 90% of the ingested sodium is excreted in the urine (Wilson, 1966b), and the kidney maintains fluid and electrolyte balance in animals drinking salty water, if sufficient fresh water is available. However, the digestive system is unable to cope with a large volume of saline water ingested to maintain fluid balance (Wilson and Dudzinski, 1973), and can lead to a higher osmotic pressure and a lower microbial population and activity in the rumen as found in sheep (Potter et al., 1972). The degree of change in osmotic pressure, microbial population and activity depends on the type of ration. For example, sheep have a higher tolerance to saline water when fed on lucerne chaff than on oaten chaff (Wilson and Dudzinski, 1973). There is no quantitative information on the effects of high sodium intake on the digestion in the rumen, but Elam (1961) recorded an increase in the rate of passage of feed through the gut when salt was added to the diet.

Feed intake and animal production

While there is no data on the feed intake and growth rate of deer at different levels of salt intake, studies with other animal species shows that increasing salt intake often leads to a higher water requirement and a reduction of feed intake, especially in hot weather, on saline vegetation or on a poor quality ration. Wilson (1966b) reported that the addition of 0.9 or 1.2% sodium chloride to the drinking water for sheep fed saltbush diets caused a marked decrease in intake and a substantial loss of body weight, with a small decrease in digestibility. Generally, total salt content of 1% in drinking water will not influence sheep performance. Saline water about 1.5% of total salts may not affect animal production if the environmental conditions are good. However, over 2% total salts in the drinking water is detrimental to production and survival of sheep (Wilson, 1978).

The response of animals to salt intake is dependent on the age of animals. For adult sheep, 1.6% of total salt in water significantly depresses weight gain in summer, but for lambs, 1.3% of total salt causes a reduction in body weight gain and wool growth, and increases the incidence of diarrhoea. Even water containing 1.0% of total salts causes some reduction in weight gain and wool growth of lambs (Wilson, 1975). Potter and McIntosh (1974) also reported that drinking water containing 1.3% sodium chloride increases lamb mortality at birth, particularly for twins. Sheep on saline water have higher body water content and their carcasses contain less fat, but more water and protein than sheep on fresh water (Walker et al., 1971). This might indicate that weight change, without determination of body composition, is probably not adequate for assessing the effect of saline water on

animal production.

SALT TOLERANCE OF DEER

Salt tolerance has been interpreted as the absence of a depression in the intake of a certain ration, and is well defined for most livestock species except for deer. Under grazing condition, animals may ingest excessive salt through feed, drinking water, and ingestion of soil (Howell, 1996). Excessive intake of sodium is one of the more commonly encountered problems and often causes loss of appetite, reduced milk production and reduced growth (Xu et al., 1994). Water is one of major sources of salt for grazing animals. Wilson (1978) reported that sheep can tolerate 1.5% of sodium chloride in their drinking water, and Peirce (1959) recommended 1.3% as a safe maximum concentration, although 2.0% could be tolerated (Wilson, 1966b). However, Peirce (1957, 1962, 1966, 1968a, 1968b) found that water containing 1.0-1.3% soluble salts reduced appetite and caused some deaths in adult sheep, and sometimes reduced the number of lambs born and their growth rate, although saline water (1.3% NaCl) had no effect on the concentrations of sodium, potassium, calcium or chloride in the blood plasma (Peirce, 1959, 1963). Australian Water Resources Council recommended that the ground water might be suitable for some stock if the content of total dissolved solid is 3000-7000 mg/litre. If the content of total dissolved solid is over 14000 mg/litre, this water cannot be used by any stock (Anon, 1976). Inglis (1985) summarised the salinity tolerance of livestock and poultry based on previous research (see table 4). The salinity tolerance can vary with environmental conditions, type of feed, or physiological stages (see table 5, Anon, 1969). Sheep and cattle grazing green feed can tolerate higher concentrations of salt in water than the same stock on dry feed. Stock grazing on saltbush are less tolerant than those grazing other types of pastures. Pregnant, lactating or young animals

have a poorer tolerance to saline water than dry mature animals (Inglis, 1985). Peirce (1957, 1962, 1966, 1968a, 1968b) and Wilson (1975) also reported that ewes and lambs are less tolerant of saline water than the non-pregnant and older sheep, and older sheep or sheep with previous experience of saline water are more tolerant than young and inexperienced sheep. Therefore, the safe values for salinity of 1.2% on grassland and 0.8% on saltbush were recommended by Wilson (1975), but these values should be further reduced for ewes and lambs, if lambs are born in summer.

There are large differences in salinity tolerance between animal species or breeds within species (NRC, 1980; Inglis, 1985; Judson and McFarlane, 1998). NRC (1980) reported that the maximum tolerable concentration were 35 and 16-35 g/kg DM for sheep and cattle, respectively. For sheep different tolerances are observed when fed on different rations or levels of intake due to individual differences in their ability to excrete or tolerate sodium chloride (Wilson, 1966b).

POTENTIAL DEER FARMING IN THE ARID AND SEMI-ARID AREAS

In the arid and semi arid regions which cover 70 to 75% of the continent (Ffolliott et al., 1995), with only 2-3% of the population (Mitchell, 1979), rainfall is variable and of low intensity. Typically the land is flat and only wide, shallow storage of water is possible. The high evaporation rates result from high radiation, high temperature and low humidity, especially in summer. Water quality, especially high concentrations of salts present some of the greatest limitations for the use of water by livestock (Anon, 1976; Goodspeed and Winkworth, 1978).

The development of deer farming in the arid and semi-arid areas of Australia offers diversification opportunities for existing farmers as well as new farmers. It has been well understood that deer are very adaptable and can be farmed in most parts of Australia (Mackay, 1998). The competition between cattle and deer for feed resources is minimal due to little dietary overlap between the two species (Ffolliott et al., 1995). More importantly, deer are more efficient converters of fodder to meat than cattle or sheep, have a dressing percentage of around 60% which is significantly better than either sheep or cattle (Anderson, 1984). Blaxter (1974) also reported that the first class meat obtained from a lean young deer carcass is 33% of its empty body weight. In sheep of a similar size the amount of first class meat obtained is only 18% of empty body weight. The feed conversion ratio was 2.8-3.8 which was half that usually found for cattle and better than that obtained in intensive lamb production. Drew and Greer (1977)

Table 4. Salinity tolerance of livestock and poultry (mg total soluble salts/litre water)

Animal	Maximum concentration for health growth	Maximum concentration to maintain conditions	Maximum concentration tolerated
Sheep	6000	13000	Depends on type of feed
Beef cattle	4000	5000	10000
Dairy cattle	3000	4000	6000
Horses	4000	6000	7000
Pigs	2000	3000	4000
Poultry	2000	3000	3500

Source: Inglis (1985).

Table 5. Upper limits used by State Authorities for total salts in water (ppm) for livestock

Class of livestock	New South Wales	Victoria	Queensland	South Australia	Western Australia	Northern Territory
Poultry	4,000	3,500	3,500	3,000 ^a	3,000	
Pigs	4,000	4,500	5,500	3,000 ^a	4,500	
Horses	7,000	6,000	5,500 ^b	7,000	6,500	6,000
Cattle						
Dairy	10,000	6,000	5,500 ^b	7,000	7,000	6,000
Beef	10,000	7,000	8,500 ^c	10,000	10,000	10,000
Sheep						
Lamb, Weaners		4,500			10,000	
Ewes in milk		6,000			10,000	
Adult, on dry feed	14,000	7,000 ^e	14,500	13,000	13,000	12,000
Adult, on green grass		15,000	18,500	18,000	18,500 ^d	15,000

^a 4,000 ppm if on salt-free rations, 1,500 ppm if on high salt diet.

(Anon, 1969)

^b 7,000 ppm if on green feed.

^c 10,000 ppm if on green feed.

^d for short period.

^e If unaccustomed to saline water; if accustomed, up to 15,000 ppm.

estimated that pasture dry matter intakes for every kg carcass gain were 9.5 kg for red deer and 30 kg for lamb. Wilson (1984) also confirmed the potential advantage of deer meat production over traditional beef production and found that deer carcasses have little fat and most is subcutaneous, intermuscular and retroperitoneal.

The vegetation of large areas of arid and semi-arid Australia is dominated by species of the genera *Atriplex* (saltbushes) and *Kochia* (bluebushes). These shrubs are invaluable feed resources in summer for grazing sheep and cattle. While the nutritional value of these bushes are evaluated by a number of researchers, the high salt concentration in leaves is a disadvantage to animal production. High salt intake of grazing animals from bushes and salty water in these regions, especially over drought periods, causes a higher water requirement, which may limit the expansion of beef industry in these regions (Wilson, 1979). Thus the lower daily water requirement of deer in comparison with beef cattle offers a great opportunity to use this land for deer farming. However, farmers need information on salt tolerance of deer to optimise the production system and eliminate animal ethics concerns.

CONCLUSION

Animals require sodium for growth, reproduction and other physiological functions. However, in southern Australia or in the salt-affected areas, excessive intake of sodium often occurs through drinking water, ingestion of salty feed and soil, especially in dry summers when temperature and evaporation rates are high. The impact of excessive

sodium intake on feed intake, growth rate and water requirement has been well documented for most livestock species except for deer. While the salt tolerance levels of grazing animals have been defined, especially for sheep, there is no data available for grazing deer. More importantly, the tolerance level is influenced by species, breeds within species, physiological stages of animal, feed type, feeding level, and environmental condition. Therefore the data obtained from sheep and cattle cannot simply be applied to deer. To optimise deer production in southern Australia and to successfully develop the deer industry in the dry areas or salt-affected areas the following research is required;

- i) basic information of sodium status of deer in these regions particularly from an animal ethics point of view,
- ii) an understanding of the impact of excessive sodium intake on deer production,
- iii) tolerance levels of different deer species at various physiological stages.

ACKNOWLEDGEMENT

The authors would like to acknowledge Mr. Warwick Hack and Mr. Chris Tuckwell for their helpful advice during the preparation of this manuscript.

REFERENCES

- Anderson, R. 1984. Deer farming in Australia. In Deer Refresher Course. The Post-graduate Committee in Veterinary Science, The University of Sydney. pp.

- 55-103.
- Anon. 1969. Quality aspects of farm water supplies. Victoria Irrigation Research and Advisory Services Committee. The Department of National Development, Canberra.
- Anon. 1976. Review of Australia's Water Resources. Department of National Resources, Australian Water Resources Council. Australian Government Publishing Service Canberra.
- Anon. 1988a. Report on salinity in Western Australia, Discussion paper, Parliament of Western Australia, Legislative Council. p. 72.
- Anon. 1988b. Submission to the House of Representatives Standing Committee on Environment, Recreation and the Arts. Inquiry into Land Degradation. South Australian Government.
- Anon. 1988c. Land degradation survey of New South Wales 1987-88. Soil Conservation Service of New South Wales p. 32.
- Anon. 1989. Salt action : joint action. Victoria's strategy for managing land and water action. Government of Victoria. p. 54
- Anon. 2000. Climate averages. Bureau of Meteorology. <http://www.bom.gov.au/climate/averages/tables>.
- Barren-Lennard, E. G. 1999. Puccinellia - for productive saltland pastures. Agriculture Western Australia Farmnote 1/99.
- Barrett-Lennard, E. G. and R. A. Nulsen. 1989. Dryland soil salinity-cure, containment or catastrophe? Proc. of the 5th Aust. Agron. Conf. Western Australia: Australian Society of Agronomy. pp. 212-220.
- Bellotti, W., W. Collins and A. Moore. 1992. The mediterranean environments. In: Management for Wool Quality in Mediterranean Environments (Ed. P. T. Doyle, J. A. Fortune and N. R. Adams). Department of Agriculture Western Australia, Perth, Australia. pp. 50-59.
- Blaxter, K. L., R. N. B. Kay, G. A. M. Sharman, J. M. M. Cunningham and W. J. Hamilton. 1974. Farming the red deer. Department of Agriculture and Fisheries for Scotland, Edinburgh, Her Majesty's Stationery Office.
- Bonengel, H. 1969. Feeding trials with growing red deer. Theses, Tierarztl. Fak. Univ. German. p. 117.
- Denholm, L. J. 1984. The nutrition of farmed deer. In: Deer Refresher Course. The Post-graduate Committee in Veterinary Science, The University of Sydney. pp. 662-707.
- Dewes, H. F. 1996. The rate of soil ingestion by dairy cows and the effect on available copper, calcium, sodium and magnesium. N. Z. Vet. J. 44:199-200.
- Drew, K. R. and G. J. Greer. 1977. Venison production and carcass composition of red deer. N. Z. J. Agric. Sci. 11:187.
- Elam, C. J. 1961. Effect of high salt intake on digestibility in cattle. J. Anim. Sci. 20:931.
- English, A. W. 1984. The Australian Deer Farming Industry-progress and prospects. In: Deer Refresher Course. The Post-graduate Committee in Veterinary Science, The University of Sydney. pp. 533-540.
- Ffolliott, P. F., K. N. Brooks, H. M. Gregersen and L. A. Lundgren. 1995. Dryland Forestry Planning and management. John Wiley & Sons, New York. p. 22.
- Goodspeed, M. J. and R. E. Winkworth. 1978. Fate and effect of runoff-with particular reference to arid and semi-arid plain lands. In: Studies of the Australian Arid Zone (Ed. K. M. W. Howes). CSIRO, Australia, p. 53.
- Healy, W. B. 1973. Nutritional aspects of soil ingestion by grazing animals. In: Chemistry and Biochemistry of Herbage (Ed. G. W. Buller and R. W. Bailey). Academic Press: New York, USA. pp. 567-588.
- Henschke, C. J. 1980. Saltland in statistics the 1979 saltland survey. J. Agric. Western Aust. 4:116-119.
- Holmes, J. W. and J. D. Waterhouse. 1983. Hydrology. In: Natural History of the South East (Ed. M. J. Tyler, C. R. Twidale, J. K. Ling and J. W. Holmes). Occasional Publication of the Royal Society of South Australia Inc., Graphic Service Pty Ltd, Northfield. pp. 49-59.
- Howell, J. M. 1996. Toxicities and excessive intakes of minerals. In: Detection and Treatment of Mineral Nutrition Problems in Grazing Sheep (Ed. D. G. Masters and C. L. White). ACIAR Monograph No. 37. Watson Ferguson & Co., Brisbane, Australia. pp. 95-117.
- Inglis, S. 1985. Livestock water supplies. Fact sheet 82/77. Department of Agriculture, South Australia.
- Judson, G. J. and I. D. McFarlane. 1998. Mineral disorders in grazing livestock and the usefulness of soil and plant analysis in the assessment of these disorders. Aust. J. Exp. Agric. 38:707-723.
- Khalil, J. K., W. N. Sawaya and S. Z. Hyder. 1986. Nutrient composition of *Atriplex* leaves grown in Saudi Arabia. J. Range Manage. 39:104-107.
- Leigh, J. H. 1986. Forage value and utilisation of chenopod dominated shrubland. Reclamation and Revegetation Res. 5:387-402.
- Mackay, B. I. 1985. Report on study tour of New Zealand Deer Farming Industry. NSW Department of Agriculture. Orange, Australia.
- Mackay, B. I. 1998. Deer Farming. In: The New Rural Industries (Ed. K. W. Hyde). RIRDC, Canberra, Australia. pp. 32-39.
- Malcolm, C. V. 1986. Western Australian Department of Agriculture. Farmnote 32/86. Perth Western Australia, Australia.
- Malcolm, C. V. 1989. Pasture production from saltland. Proc. 5th Aust. Agron. Conf. Western Australia, Australia. pp. 67-73.
- Mitchell, A. S. 1979. Alternative enterprises in arid Northern Territory. 2nd Biennial Conference Working Papers. Australian Rangeland Society, Adelaide, Australia. pp. 4-1, 4-25.
- Mozafar, A. and J. R. Goodin. 1970. Vesiculated hairs: a mechanism for salt tolerance in *Atriplex halimus* L. Plant physiol. 45:62-5.
- Nichols, C. 1998. Establishing and managing tall wheat grass in saline soils for productivity. Agriculture Notes (AG0707), Natural Resources and Environment, Victoria, Australia.
- NRC. 1975. Nutrient requirements of domestic animals. No. 5. Nutrient requirements of sheep. National Research Council. Washington, DC.
- NRC. 1976. Nutrient requirements of domestic animals. No. 4. Nutrient requirements of beef cattle. National Research Council. Washington, DC.
- NRC. 1980. Mineral tolerance of domestic animals. National Research Council. Washington, DC., USA.
- NRC. 1981. Nutrient requirements of domestic animals. No.

15. Nutrient requirements of goats. National Research Council. Washington, DC.
- Peirce, A. W. 1957. Studies on salt tolerance of sheep. I. The tolerance of sheep for sodium chloride in the drinking water. *Aust. J. Agric. Res.* 8:711-722.
- Peirce, A. W. 1959. Studies on salt tolerance of sheep. II. The tolerance of sheep for mixtures of sodium chloride and magnesium chloride in the drinking water. *Aust. J. Agric. Res.* 10:725-735.
- Peirce, A. W. 1962. Studies on salt tolerance of sheep. IV. The tolerance of sheep for mixtures of sodium chloride and calcium chloride in the drinking water. *Aust. J. Agric. Res.* 13:479-486.
- Peirce, A. W. 1963. Studies on salt tolerance of sheep. V. The tolerance of sheep for mixtures of sodium chloride, sodium carbonate, and sodium bicarbonate in the drinking water. *Aust. J. Agric. Res.* 14:815-823.
- Peirce, A. W. 1966. Studies on salt tolerance of sheep. VI. The tolerance of wethers in pens for drinking waters of the types obtained from underground sources in Australia. *Aust. J. Agric. Res.* 17:209-218.
- Peirce, A. W. 1968a. Studies on salt tolerance of sheep. VII. The tolerance of ewes and their lambs in pens for drinking waters of the types obtained from underground sources in Australia. *Aust. J. Agric. Res.* 19:577-587.
- Peirce, A. W. 1968b. Studies on salt tolerance of sheep. VIII. The tolerance of grazing ewes and their lambs for drinking waters of the types obtained from underground sources in Australia. *Aust. J. Agric. Res.* 19:589-595.
- Perry, M. W. 1989. Farming systems of southern Australia. *Proc. the 5th Aust. Agron. Conf., Western Australia, Australia.* pp. 67-73.
- Potter, B. J. and G. H. McIntosh. 1974. Effect of salt water ingestion on pregnancy in the ewe and on lamb survival. *Aust. J. Agric. Res.* 25:909-917.
- Potter, B. J., D. J. Walker and W. W. Forrest. 1972. Changes in intraruminal function of sheep when drinking saline water. *Br. J. Nutr.* 27:75.
- Rehman, A. 1992. Factors affecting preference of sheep for saltbush (*Atriplex spp.*). Masters Thesis, University of Western Australia, Australia.
- Ru, Y. J. 1996. The influence of cultivar variation on the potential production of swards of subterranean clover when utilised by grazing animals. Ph.D. Thesis. University of Adelaide, Australia.
- SCA. 1980. Deer production in Australia. Canberra, Australia. pp. 1-2.
- Sharma, M. L. 1978. Water use by Chenopod shrublands. In: *Studies of the Australian Arid Zone* (Ed. K. M. W. Howes). CSIRO, Australia. p. 139.
- Thornton, I. 1983. Soil-plant-animal interactions in relation to the incidence of trace element disorders in grazing livestock. In: *Trace Elements in Animal Production and Veterinary Practice* (Ed. N. F. Suttle, R. G. Gunn, W. M. Allen, K. A. Linklater and G. Wiener). Occasional Publication of the British Society of Animal Production No. 7. Haddington, UK. pp. 39-49.
- Walker, D. J., B. J. Potter and G. B. Jones. 1971. Modification of carcass characteristics in sheep maintained on a saline water regime. *Aust. J. Exp. Agric. Anim. Husb.* 11:14-17.
- Wilson, A. D. 1966a. The value of *Atriplex* (saltbush) and *Kochia* (bluebush) species as food for sheep. *Aust. J. Agric. Res.* 17:147-153.
- Wilson, A. D. 1966b. The intake and excretion of sodium by sheep fed on species of *Atriplex* (saltbush) and *Kochia* (bluebush). *Aust. J. Agric. Res.* 17:155-163.
- Wilson, A. D. 1975. Influence of water salinity on sheep performance while grazing on natural grassland and saltbush pastures. *Aust. J. Exp. Agric. Anim. Husb.* 15:760.
- Wilson, A. D. 1978. Water requirements of sheep. In: *Studies of the Australian Arid Zone* (Ed. K. M. W. Howes). CSIRO, Australia. p. 179.
- Wilson, A. D. 1979. Cattle and sheep production on saltbush. 2nd Biennial Conference Working Papers, Australian Rangeland Society, Adelaide, Australia. pp 3-11, 3-18.
- Wilson, A. D. and M. L. Dudzinski. 1973. Influence of the concentration and volume of saline water on the food intake of sheep, and on their excretion of sodium and water in urine and faeces. *Aust. J. Agric. Res.* 24:245-256.
- Wilson, A. D. and N. L. Hindley. 1968. Comparison of the voluntary water intakes of some Australian sheep. *Field Station Records* 7, 25 Division of Plant Industry. CSIRO, Australia.
- Wilson, P. R. 1984. Nutrition and reproduction of farmed deer. In: *Deer Refresher Course. The Post-graduate Committee in Veterinary Science, The University of Sydney, Australia.* pp. 95-103.
- Wood, J. G. 1925. The selective absorption of chlorine ions and absorption of water by the leaves of the genus *Atriplex*. *Aust. J. Exp. Biol. Med. Sci.* 2:45-56.
- Xu, Z. Y., S. X. Yang and S. Yang. 1994. *Animal Nutrition*. Chinese Agricultural Publishing Co., Beijing, China. p. 44.