

ON A SUBCLASS OF PRESTALIKE FUNCTIONS WITH NEGATIVE COEFFICIENTS

S. K. LEE AND S. B. JOSHI

ABSTRACT. Motivated by recent work of Uralegaddi and Sarangi[12], we aim at presenting here system study of novel subclass $R_\alpha[\mu, \beta, \xi]$ of prestarlike functions. Further using operators of fractional calculus, we have obtained distortion theorem for $R_\alpha[\mu, \beta, \xi]$. Lastly the extreme points of $R_\alpha[\mu, \beta, \xi]$ are obtained.

1. Introduction.

Let A denote the class of functions of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

that are analytic in the unit disc $U = \{z : |z| < 1\}$, let S denote the subclass of A consisting of analytic and univalent functions $f(z)$ in the unit disc U . Further T denote subclass of A consisting of functions $f(z)$ of the form

$$(1.2) \quad f(z) = z - \sum_{n=2}^{\infty} a_n z^n, \quad a_n \geq 0.$$

Schild[8] studied a subclass of S consisting of polynomials having $|z| = 1$ as radius univalence. Subsequently, Silverman[10] proved useful results for the subclasses $S^*(\alpha)$ and $C(\alpha)$ of S , where $S^*(\alpha)$ and $C(\alpha)$ denote respectively, the subclasses of starlike functions of order α and convex functions of order α , $0 \leq \alpha < 1$. We note that $S^*(\alpha)$ was introduced by Robertson[5].

Received June 8, 2000.

1991 Mathematics Subject Classification: Primary 30C45.

Key words and phrases: Fractional calculus, univalent function, distortion theorem, extreme points.

The function

$$(1.3) \quad S_\alpha(z) = z(1-z)^{-2(1-\alpha)}$$

is the well known extremal function for the class $S^*(\alpha)$. Letting,

$$(1.4) \quad C(\alpha, n) = \frac{\prod_{k=2}^n (k-2\alpha)}{(n-1)!}, \quad n = 2, 3, \dots$$

$S_\alpha(z)$ can be written in the form

$$(1.5) \quad S_\alpha(z) = z + \sum_{n=2}^{\infty} C(\alpha, n)z^n.$$

We note that $C(\alpha, n)$ is decreasing in α and satisfies

$$(1.6) \quad \lim_{n \rightarrow \infty} C(\alpha, n) = \begin{cases} \infty & \alpha < \frac{1}{2} \\ 1 & \alpha = \frac{1}{2} \\ 0 & \alpha > \frac{1}{2}. \end{cases}$$

Let $(f * g)(z)$ denote the convolution or Hadamard product of $f(z)$ given by (1.1) and $g(z)$ given by

$$g(z) = z + \sum_{n=2}^{\infty} b_n z^n,$$

then

$$(1.8) \quad (f * g)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n.$$

Let $R_\alpha(\mu, \beta, \xi)$ denote the class of prestarlike functions, that satisfies the condition

$$(1.9) \quad \left| \frac{\frac{zh'(z)}{h(z)} - 1}{2\xi\left(\frac{zh'(z)}{h(z)} - \mu\right) - \left(\frac{zh'(z)}{h(z)} - 1\right)} \right| < \beta$$

where, $h(z) = f * S_\alpha(z)$, $0 < \beta \leq 1$, $0 \leq \mu < 1$, $1/2 < \xi \leq 1$. The class of α -prestarlike functions was introduced by Ruscheweyh[7] and later on rather extensively studied by Silverman and Silvia[9], Owa and Ahuja[4] and Uralegaddi and Sarangi[12].

Let

$$(1.10) \quad R_\alpha[\mu, \beta, \xi] = R_\alpha(\mu, \beta, \xi) \cap T.$$

Our main tool in the present paper is the following, which can be easily proved, the details are omitted.

LEMMA 1. Let $f(z)$ be defined by (1.2), then $f(z)$ is in the class $R_\alpha[\mu, \beta, \xi]$ if and only if

$$\sum_{n=2}^{\infty} ((n - 1) + \beta(2\xi(n - \mu) - (n - 1))) C(\alpha, n)a_n \leq 2\beta\xi(1 - \mu).$$

The result is sharp.

2. Distortion Theorems Involving Fractional Calculus

In this section, we shall prove distortion theorems for functions belonging to the class $R_\alpha[\mu, \beta, \xi]$. Each of these would involve operators of fractional calculus which are defined as follows (cf. e.g [2, 3, 6, 11]).

DEFINITION 1. The fractional integral of order λ is defined by

$$(2.1) \quad D_z^{-\lambda} f(z) = \frac{1}{\Gamma(\lambda)} \int_0^z \frac{f(\zeta)}{(z - \zeta)^{1-\lambda}} d\zeta$$

where $\lambda > 0$, $f(z)$ is an analytic function in a simply connected region of the z -plane containing the origin, and the multiplicity of $(z - \zeta)^{\lambda-1}$ is removed by requiring $\log(z - \zeta)$ to be real when $z - \zeta > 0$.

DEFINITION 2. The fractional derivative of order λ is defined by

$$(2.2) \quad D_z^\lambda f(z) = \frac{1}{\Gamma(1 - \lambda)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z - \zeta)^\lambda} d\zeta,$$

where $0 \leq \lambda < 1$, $f(z)$ is an analytic function in a simply connected region of the z -plane containing the origin and the multiplicity of $(z - \zeta)^{-\lambda}$ is removed as in Definition 1.

DEFINITION 3. Under the hypothesis of Definition 2, the fractional derivative of order $n + \lambda$ is defined by

$$(2.3) \quad D_z^{n+\lambda} f(z) = \frac{d^n}{dz^n} D_z^\lambda f(z),$$

where $0 \leq \lambda < 1$, $n \in N \cup \{0\}$, $N = \{1, 2, \dots\}$.

THEOREM 1. Let $f(z)$ given by (1.2) be in the class $R_\alpha[\mu, \beta, \xi]$. Then

$$(2.4) \quad |D_z^{-\lambda} f(z)| \geq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left(1 - \frac{\beta\xi(1-\mu)}{(1+\beta(4\xi-2\xi\mu-1))(1-\alpha)} |z| \right)$$

and

$$(2.5) \quad |D_z^{-\lambda} f(z)| \leq \frac{|z|^{1+\lambda}}{\Gamma(2+\lambda)} \left(1 + \frac{\beta\xi(1-\mu)}{(1+\beta(4\xi-2\xi\mu-1))(1-\alpha)} |z| \right)$$

for $\lambda > 0$, $z \in U$. The bounds are sharp.

Proof. Let

$$(2.6) \quad \begin{aligned} F(z) &= \Gamma(2+\lambda) z^{-\lambda} D_z^{-\lambda} f(z) \\ &= z - \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)} a_n z^n \end{aligned}$$

for $\lambda > 0$. We note that

$$(2.7) \quad 0 < \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)} < n$$

for $\lambda > 0$, $n \geq 2$, and that $C(\alpha, n+1) \geq C(\alpha, n)$, for $0 \leq \alpha < 1/2$, and $n \geq 2$. Consequently, by using Lemma 1, we have

$$\begin{aligned} |F(z)| &\geq |z| - |z|^2 \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)} a_n \\ &\geq |z| - \frac{\beta\xi(1-\mu)}{(1+\beta(4\xi-2\xi\mu-1))(1-\alpha)} |z|^2 \end{aligned}$$

which implies (2.4), and

$$\begin{aligned} |F(z)| &\leq |z| + |z|^2 \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2+\lambda)}{\Gamma(n+1+\lambda)} a_n \\ &\leq |z| + \frac{\xi\beta(1-\mu)}{(1+\beta(4\xi-2\xi\mu-1))(1-\alpha)} |z|^2 \end{aligned}$$

which gives (2.5).

The result is sharp for the function $f(z)$ given by

$$D_z^{-\lambda} f(z) = \frac{z^{1+\lambda}}{\Gamma(2+\lambda)} \left(1 - \frac{\beta\xi(1-\mu)}{(1+\beta(4\xi-2\xi\mu-1))(1-\alpha)} z \right).$$

□

COROLLARY 1. *Let the functions $f(z)$ be defined by (1.2) is in the class $R_\alpha[\mu, \beta, \xi]$, with $0 \leq \alpha \leq 1/2$, $1/2 < \xi \leq 1$, $0 < \beta \leq 1$ and $0 \leq \mu < 1$. Then $D_z^{-\lambda}f(z)$ is included in a disc with center at origin and radius r_1 given by*

$$(2.8) \quad r_1 = \frac{1}{\Gamma(2 + \lambda)} \left(1 + \frac{\beta\xi(1 - \mu)}{(1 + \beta(4\xi - 2\xi\mu - 1))(1 - \alpha)} \right),$$

where $\lambda > 0$.

THEOREM 2. *Let the functions $f(z)$ given by (1.2) be in the class $R_\alpha[\mu, \beta, \xi]$. Then*

$$(2.9) \quad |D_z^\lambda f(z)| \geq \frac{|z|^{1-\lambda}}{\Gamma(2 - \lambda)} \left(1 - \frac{2\xi\beta(1 - \mu)}{(1 + \beta(4\xi - 4\xi\mu - 1))(1 - \alpha)} |z| \right)$$

and

$$(2.10) \quad |D_z^\lambda f(z)| \leq \frac{|z|^{1-\lambda}}{\Gamma(2 - \lambda)} \left(1 + \frac{2\xi\beta(1 - \mu)}{(1 + \beta(4\xi - 4\xi\mu - 1))(1 - \alpha)} |z| \right)$$

for $0 \leq \lambda < 1$, $z \in U$. The bounds are sharp.

Proof. Let

$$\begin{aligned} G(z) &= \Gamma(2 - \lambda)z^\lambda D_z^\lambda f(z) \\ &= z - \sum_{n=2}^{\infty} \frac{\Gamma(n + 1)\Gamma(2 - \lambda)}{\Gamma(n + 1 - \lambda)} a_n z^n \end{aligned}$$

for $0 \leq \lambda < 1$. By using Lemma 1, we observe that

$$\begin{aligned} (2.11) \quad & \frac{1}{2} (1 + \beta(4\xi - 4\xi\mu - 1)) C(\alpha, 2) \sum_{n=2}^{\infty} n a_n \\ & \leq \sum_{n=2}^{\infty} ((n - 1) + \beta(2\xi(n - \mu) - (n - 1))) C(\alpha, n) a_n \\ & \leq 2\beta\xi(1 - \mu), \end{aligned}$$

which implies that

$$(2.12) \quad \sum_{n=2}^{\infty} na_n \leq \frac{2\xi\beta(1-\mu)}{(1+\beta(4\xi-4\xi\mu-1))(1-\alpha)}.$$

Further, we note that $1 < \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} < n$ for $0 \leq \lambda < 1$, $n \geq 2$. Hence we have

$$\begin{aligned} |G(z)| &\geq |z| - |z|^2 \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} a_n \\ &\geq |z| - |z|^2 \sum_{n=2}^{\infty} na_n \\ &\geq |z| - \frac{2\xi\beta(1-\mu)}{(1+\beta(4\xi-4\xi\mu-1))(1-\alpha)} |z|^2 \end{aligned}$$

which proves (2.9), and

$$\begin{aligned} |G(z)| &\leq |z| + |z|^2 \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} a_n \\ &\leq |z| + |z|^2 \sum_{n=2}^{\infty} na_n \\ &\leq |z| + \frac{2\xi\beta(1-\mu)}{(1+\beta(4\xi-4\xi\mu-1))(1-\alpha)} |z|^2 \end{aligned}$$

which gives (2.10).

Finally, The bound of (2.9) and (2.10) are sharp, extremal function being

$$D_z^\lambda f(z) = \frac{z^{1-\lambda}}{\Gamma(2-\lambda)} \left(1 - \frac{2\xi\beta(1-\mu)}{(1+\beta(4\xi-4\xi\mu-1))(1-\alpha)} \right).$$

□

COROLLARY 2. *Let the function $f(z)$ given by (1.2) be in the class $R_\alpha[\mu, \beta, \xi]$. Then $D_z^\lambda f(z)$ is included in the disc with center at origin and radius r_2 given by,*

$$(2.13) \quad r_2 = \frac{1}{\Gamma(2-\lambda)} \left(1 + \frac{2\xi\beta(1-\mu)}{(1+\beta(4\xi-4\xi\mu-1))(1-\alpha)} \right),$$

where $0 \leq \lambda < 1$.

Finally, we obtain extreme points of $R_\alpha[\mu, \beta, \xi]$ by the routine calculation.

THEOREM 3. *Let*

$$f_1(z) = z$$

and

$$(2.14) \quad f_n(z) = z - \frac{2\beta\xi(1-\mu)}{((n-1) + \beta(2\xi(n-\mu) - (n-1)))C(\alpha, n)} z^n,$$

$n = 2, 3, \dots$. Then $f \in R_\alpha[\mu, \beta, \xi]$ if and only if it can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \lambda_n f_n(z),$$

where $\lambda_n \geq 0$, $\sum_{n=1}^{\infty} \lambda_n = 1$.

References

1. O. P. Ahuja and H. Silverman, *Convolutions of prestarlike functions*, Internat. J. Math. and Math. Sci. **6** (1983), 59–68.
2. A. Erdelyi, W. Magnus, F. Oberbettinger and F. G. Tricomi, *Tables of Integral Transforms Vol II*, McGraw Hill Book Company, New York, London and Toronto, 1954.
3. S. Owa, *On distortion theorems I*, Kyungpook Math. J. **18** (1978), 53–59.
4. S. Owa and O. P. Ahuja, *An application of the fractional calculus*, Math. Japon. **30** (1985), 947–955.
5. M. S. Robertson, *On the theory of univalent functions*, Ann. Math. **37** (1936), 374–408.
6. B. Ross, *A brief history and exposition of the fundamental theory of fractional calculus*, Fractional Calculus and its Applications (Ed. B. Ross), Springer-Verlag, Berlin, Heidelberg and New York (1975), 1–36.

7. St Ruscheweyh, *Linear operator between classes of prestarlike functions*, Comm. Math. Helv. **52** (1977), 497–509.
8. A. Schild, *On starlike functions of order α* , Amer. J. Math. **87** (1965), 65–70.
9. H. Silverman and E. M. Silvia, *Prestarlike functions with negative coefficients*, Internat J. Math. and Math. Sci. **2** (1979), 427–439.
10. H. M. Silverman, *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc. **51** (1975), 109–116.
11. H. M. Srivastava and S. Owa (Editors), *Univalent functions, Fractional Calculus and their Applications*, Halsted Press (Ellis Harwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1989.
12. B. A. Uralegaddi and S. M. Sarangi, *Certain generalization of prestarlike functions with negative coefficients*, Ganita **34** (1983), 99–105.

Department of Mathematics
Gyeongsang National University
Jinju 660-701, Korea
E-mail: sklee@nongae.gsnu.ac.kr

Department of Mathematics
Walchand College of Engineering
Sangli-416 415, India
E-mail: joshisb@hotmail.com